

I am not robot!

Intensive care society respiratory guidelines

The FICM/ICS Guideline Development Group have used GRADE methodology to make the following recommendations for the management of adult patients with acute respiratory distress syndrome (ARDS). The British Thoracic Society supports the recommendations in this guideline. Where mechanical ventilation is required, the use of low tidal volumes (< 6 ml/kg ideal body weight) and airway pressures (plateau pressure < 30 cmH2O) was recommended. For patients with moderate/severe ARDS (PF ratio < 20kPa), prone positioning was recommended for at least 12 hours per day. devibi By contrast, high frequency oscillation is not recommended and it is suggested that inhaled nitric oxide is not used. The use of a conservative fluid management strategy was suggested for all patients, whereas mechanical ventilation with high positive end-expiratory pressure (PEEP) and the use of the neuro-muscular blocking agent cisatracurium for 48 hours was suggested for ARDS patients with PF ratios less than or equal to 27 and 20 kPa respectively. Extra-corporeal membrane oxygenation (ECMO) was suggested as an adjunct to protective mechanical ventilation for patients with very severe ARDS. In the absence of adequate evidence, research recommendations were made for the use of corticosteroids and extra-corporeal carbon dioxide removal (ECCOR). cinahitoyinoyu Variation in Early Management Practices in Moderate-to-Severe ARDS in the United States: The Severe ARDS: Generating Evidence Study.



In the absence of adequate evidence, research recommendations were made for the use of corticosteroids and extra-corporeal carbon dioxide removal (ECCOR).



The use of a conservative fluid management strategy was suggested for all patients, whereas mechanical ventilation with high positive end-expiratory pressure (PEEP) and the use of the neuro-muscular blocking agent cisatracurium for 48 hours was suggested for ARDS patients with PF ratios less than or equal to 27 and 20 kPa respectively. Extra-corporeal membrane oxygenation (ECMO) was suggested as an adjunct to protective mechanical ventilation for patients with very severe ARDS. In the absence of adequate evidence, research recommendations were made for the use of corticosteroids and extra-corporeal carbon dioxide removal (ECCOR). Variation in Early Management Practices in Moderate-to-Severe ARDS in the United States: The Severe ARDS: Generating Evidence Study.



Qadir N, Bartz RR, Cooter ML, Hough CL, Lanspa MJ, Banner-Goodspeed VM, Chen JT, Giovanni S, Gomaa D, Sjoding MW, Hajizadeh N, Komisarow J, Duggal A, Khanna AK, Kashyap R, Khan A, Chang SY, Tonna JE, Anderson HL 3rd, Liebler JM, Mosier JM, Morris PE, Genthon A, Louh IK, Tidswell M, Stephens RS, Esper AM, Dries DJ, Martinez A, Schreyer KE, Bender W, Tiwari A, Guru PK, Hanna S, Gong MN, Park PK; Severe ARDS: Generating Evidence (SAGE) Study Investigators; Society of Critical Care Medicine's Discovery Network.



Qadir N, Bartz RR, Cooter ML, Hough CL, Lanspa MJ, Banner-Goodspeed VM, Chen JT, Giovanni S, Gomaa D, Sjoding MW, Hajizadeh N, Komisarow J, Duggal A, Khanna AK, Kashyap R, Khan A, Chang SY, Tonna JE, Anderson HL 3rd, Liebler JM, Mosier JM, Morris PE, Genthon A, Louh IK, Tidswell M, Stephens RS, Esper AM, Dries DJ, Martinez A, Schreyer KE, Bender W, Tiwari A, Guru PK, Hanna S, Gong MN, Park PK; Severe ARDS: Generating Evidence (SAGE) Study Investigators; Society of Critical Care Medicine's Discovery Network.
Qadir N, et al. Chest.



Chest. vafozusetudafo 2021 Oct;160(4):1304-1315. doi: 10.1016/j.chest.2021.05.047. Epub 2021 Jun 4. Chest. 2021. PMID: 34089739 Free PMC article. Membership Guidance Learning Research Thriving at Work SOA24 Policy and News About ICU Donate Join Login Since 1990, ESICM has been committed to producing guidelines and consensus statements from colleagues within the society as well as in collaboration with other international societies to provide the best quality evidence for clinicians at the bedside. These manuscripts have been published in the journal Intensive Care Medicine and other leading journals (endorsement from ESICM does not imply automatic acceptance for publication in ICM journal).
If you wish to propose guidelines or consensus statements, please write to research@esicm.org and we will forward your request to the Chair of the Guidelines Group. At the end of 2020, the ESICM Executive Committee decided to initiate the development of guidelines. The society is fortunate enough to benefit from the expertise of renowned specialists and leaders in a substantial variety of topics in intensive care. Therefore it was decided to make a selection and to begin with four guidelines on ultrasound, antimicrobial, fluids and acute respiratory distress syndrome. under development: Guideline about antimicrobial stewardship for severe infections and sepsis. The co-leaders are Jan De Waele (Ghent University Hospital, Belgium) and Garyphallia Poulakou (National and Kapodistrian University of Athens, Greece). Development in progress. Guideline about fluid therapy in the ICU. mavapi The co-leaders are Daniel De Backer (CHIREC, Brussels, Belgium) and Morten Hyllander Møller (University Hospital Rigshospitalet, Copenhagen, Denmark). Development in progress. ESICM Consensus Conference on shock diagnosis and haemodynamic monitoring in the ICU. The co-chairs are Michel Chew (University of Lund, Sweden) and Xavier Monnet (University of Paris Saclay, France). Transfusion Guideline part III – reversal of anticoagulants in the bleeding critically ill (A. Vlaar, Amsterdam, The Netherlands). Weaning from Mechanical ventilation (joint guideline with ERS) (co-chairs are John Laffey, Nicole Juffermans, Martin Dres, Paolo Navalesi) ESICM Consensus Guidelines on Palliative and End of Life Care in ICU patients. The co-chairs are Elie Azoulay (University Hospital St-Louis, Paris, France), Jozef Kesecioglu (University Medical Centre, Utrecht, The Netherlands), Katerina Rusinova (General University Hospital, Prague, Czechia). 2023 Giacomo Grasselli, Carolyn S. Calfee, Luigi Camporota, Daniele Poole et al. ESICM guidelines on acute respiratory distress syndrome: definition, phenotyping and respiratory support strategies. Intensive Care Med (2023). . Published in open access on 16 June 2023. 2022 Sandroni, C., Nolan, J.P., Andersen, L.W. et al. ERC-ESICM guidelines on temperature control after cardiac arrest in adults. Intensive Care Med (2022). . Published on 28 January 2022. 2021 Robba, C., Wong, A., Poole, D. et al. lulutotitutivu Basic ultrasound head-to-toe skills for intensivists in the general and neuro intensive care unit population: consensus and expert recommendations of the European Society of Intensive Care Medicine Intensive Care Med 47, 1347–1367 (2021). doi: 10.1007/s00134-021-06486-z. Free PMC article. Paul M et al., European Society of Clinical Microbiology and Infectious Diseases (ESCMID) guidelines for the treatment of infections caused by multidrug-resistant Gram-negative bacilli (endorsed by European society of intensive care medicine), Clinical Microbiology and Infection, Transfusion strategies in non-bleeding critically ill adults: a clinical practice guideline from the European Society of Intensive Care Medicine. Vlaar AP, Oczkowski S, de Bruin S, Wijnberge M, Antonelli M, Aubron C, Aries P, Duranteau J, Juffermans NP, Meier J, Murphy GJ, Abbasciano R, Muller M, Shah A, Perner A, Rygaard S, Walsh TS, Guyatt G, Dionne JC, Cecconi M.Intensive Care Med. 2020 Apr;46(4):673-696. fuduni doi: 10.1007/s00134-019-05884-8. Epub 2020 Jan 7. PMID: 31912207. pedaxo Free PMC article. 2020 Noninvasive respiratory support in the hypoxaemic peri-operative/periprocedural patient: a joint ESA/ESICM guideline. detixiyehuvugi Intensive Care Med.(2020). Surviving Sepsis Campaign International Guidelines for the Management of Septic Shock and Sepsis-Associated Organ Dysfunction in Children. Intensive Care Med 46, 10–67 (2020). Executive Summary. Updated nomenclature of delirium and acute encephalopathy: statement of ten Societies. hohoxupifoxo Intensive Care Med. (2020). degoleyivununuwu Antimicrobial Therapeutic Drug Monitoring (TDM) in Critically Ill Adult Patients – a position paper. Intensive Care Med. 2020 Jun;46(6): 1127-1153 Antimicrobial de-escalation in critically ill patients: a position statement from a task force of the European Society of Intensive Care Medicine (ESICM) and European Society of Clinical Microbiology and Infectious Diseases (ESCMID) Critically Ill Patients Study Group (ESGCIP). Intensive Care Medicine Volume 46, pages245–265 (2020). yepolumako (PMID:31781835) Fluid therapy in neurointensive care patients: ESICM consensus and clinical practice recommendations. Intensive Care Med. 2018 Apr;44(4):449-463. Guidelines for the diagnosis and management of critical illness-related corticosteroid insufficiency (CIRCI) in critically ill patients (Part II). Crit Care Med. 2018 Apr;44(4):474-477.
Second consensus on the assessment of sublingual microcirculation in critically ill patients: results from a task force of the European Society of Intensive Care Medicine. Intensive Care Med. 2018 March 2018, Volume 44, Issue 3, pages 281–299 2017 Guidelines for the diagnosis and management of critical illness-related corticosteroid insufficiency (CIRCI) in critically ill patients (Part I). Intensive Care Med.
2017 December 2017, Volume 43, Issue 12, pages 1751–1763 Recommendations for mechanical ventilation of critically ill children from the Paediatric Mechanical Ventilation Consensus Conference (PEMVECC).Intensive Care Med. 2017 Dec;43(12):1764-1780 International ERS/ESICM/ESCMID/ALAT guidelines for the management of hospital-acquired pneumonia and ventilator-associated pneumonia: Guidelines for the management of hospital-acquired pneumonia (HAP)/ventilator-associated pneumonia (VAP). Eur Respir J. 2017 Sep 10;50(3). An Official American Thoracic Society/European Society of Intensive Care Medicine/Society of Critical Care Medicine Clinical Practice Guideline: Mechanical Ventilation in Adult Patients with Acute Respiratory Distress Syndrome. Am J Respir Crit Care Med. 2017 May 1;195(9):1253-1263. Erratum Early enteral nutrition in critically ill patients Clinical practice guidelines. Intensive Care Medicine. 2017;43(3):380-398. Surviving Sepsis Campaign: International Guidelines for Management of Sepsis and Septic Shock: 2016. Intensive Care Med. 2017 Mar;43(3):304-377. 2016 An Official ATS/AACN/ACCP/ESICM/SCCM Policy Statement: Responding to Requests for Potentially Inappropriate Treatments in Intensive Care Units. Am J Respir Crit Care Med. 2015 Jun 1;191(11):1318-30. Ventilatory support of patients with sepsis or septic shock in resource-limited settings. Intensive Care Medicine. 2016;42:100-103. 2015 2014 Consensus on circulatory shock and haemodynamic monitoring. Task force of the European Society of Intensive Care Medicine. Intensive Care Medicine, Volume 40, Issue 12 / December, 2014 Consensus Summary Statement of the International Multidisciplinary Consensus Conference on Multimodality Monitoring in Neurocritical Care. A statement for healthcare professionals from the Neurocritical Care Society and the European Society of Intensive Care Medicine. Intensive Care Medicine, Volume 40, Issue 9 / September, 2014 Clinical practice guideline on diagnosis and treatment of hyponatraemia. Intensive Care Medicine, Volume 40, Issue 3 / March, 2014 2013 Surviving Sepsis Campaign: International guidelines for management of severe sepsis and septic shock, 2012. Intensive Care Medicine – Jan 2013 2012 Mortality after surgery in Europe: a 7 day cohort study.Pearse RM. European Surgical Outcomes Study. (EuSOS) group for the Trials groups of the European Society of Intensive Care Medicine and the European Society of Anaesthesiology. Lancet. 2012 Sep 22;380(9847):1059-65. doi: 10.1016/S0140-6736(12)61148-9. Acute Respiratory Distress Syndrome – The Berlin Definition. The ARDS Definition Task Force. JAMA. 2012;307(23):2526-2533. Prospectively defined indicators to improve the safety and quality of care for critically ill patients: a report from the Task Force on Safety and Quality of the European Society of Intensive Care Medicine (ESICM). Intensive Care Med. 2012 Jan 26 [Epub ahead of print] 2011 Recommendations on basic requirements for intensive care units: structural and organisational aspects. Intensive Care Med. 2011 Oct;37(10):1575-87. Epub 2011 Sep 15 International expert statement on training standards for critical care ultrasonography Expert Round Table on Ultrasound in ICU. Intensive Care Med. 2011 Jul;37(7):1077-83. Epub 2011 May 26 Recommendations on basic requirements for intensive care units: structural and organisational aspects. Intensive Care Medicine, June 2011 2010 An Official ATS/ERS/ESICM/SCCM/SRLF Statement: Prevention and Management of Acute Renal Failure in the ICU Patient: an international consensus conference in intensive care medicine.ATS/ERS/ESICM/SCCM/SRLF Ad Hoc Committee on Acute Renal Failure. Am J Respir Crit Care Med.
2010 May 15;181(10):1128-55 2008 Surviving Sepsis Campaign: International guidelines for management of severe sepsis and septic shock: 2008. Intensive Care Medicine Vol 34:17-60 Erratum ESC guidelines for the diagnosis and treatment of acute and chronic heart failure 2008: the Task Force for the diagnosis and treatment of acute and chronic heart failure 2008 of the European Society of Cardiology. Developed in collaboration with the Heart Failure Association of the ESC (HFA) and endorsed by the European Society of Intensive Care Medicine (ESICM).ESC Committee for Practice Guidelines (CPG). Eur J Heart Fail. 2008 Oct;10(10):933-89. Epub 2008 Sep 16. Review. No abstract available. Erratum in: Eur J Heart Fail. 2010 Apr;12(4):416.
Dosage error in article text. Eur J Heart Fail. 2009 Jan;11(1):110 2007 Management of bleeding following major trauma: a European guideline. Critical Care 2007, 11:R17 The European Society of Intensive Care Medicine (ESICM) and the Surviving Sepsis Campaign (SSC).Intensive Care Med. 2007 Mar;33(3):423-5. Epub 2007 Feb 16. 2005 Executive summary of the guidelines on the diagnosis and treatment of acute heart failure. Rev Esp Cardiol. 2005 Apr;58(4):389-429.
Review. Original language-Spanish. No abstract available The European Union Directive on Clinical Research: present status of implementation in EU member states' legislations with regard to the incompetent patient.Intensive Care Medicine Vol 31(3): 476-479, 2005. 2004 Surviving Sepsis Campaign guidelines for management of severe sepsis and septic shock.Intensive Care Medicine. Vol 30(5): 536-555, 2004. The Surviving Sepsis Campaign guidelines for the management of severe sepsis and septic shock: Background, recommendations, and discussion from an evidence-based review. A supplement to Critical Care Medicine Vol 32 (11): S445-S597, 2004. 2001 2001 SCCM/ESICM/ACCP/ATS/SIS: International Sepsis Definitions Conference. (ESICM Section on Sepsis) Intensive Care Medicine. Vol 29(4): 530-538, 2003. European Task Force on ventilator-associated pneumonia (VAP). (ERS/ESICM/ESCMID/ESA) Eur Respir J. 2001 May;17(5):1034-45 PMID: 11488306 1999 Recommendations for intra-hospital transport of the severely head injured patient.
ESICM Working group on Neurosurgical intensive care. Intensive Care Medicine. Vol 25(12): 1441-1443, 1999 1998 Guidelines for the pre-hospital care of patients with severe head injuries. ESICM Working group on Neurosurgical intensive care. Intensive Care Medicine. Vol 24(11): 1221-1225, 1998 1997 Informed consent for research purposes in intensive care patients in Europe Part II. Intensive Care Medicine. Vol 23(4): 435-439, 1997. 1996 The SOFA (Sepsis-related Organ Failure Assessment) score to describe organ dysfunction/failure. (ESICM Working group on Sepsis-related problems) Intensive Care Med Vol 22(7): 707-710, 1996 Guidelines for a training programme in intensive care medicine. Intensive Care Medicine.Vol 22(2): 166-172, 1996 1994 Guidelines for the utilisation of intensive care units. Intensive Care Medicine.Vol 20 (2): 163-164, 1994 Guidelines for training in intensive care medicine. Intensive Care Medicine. Vol 20(1): 80-81, 1994 Ferguson ND, Fan E, Camporota L et al (2012) The Berlin definition of ARDS: an expanded rationale, justification, and supplementary material. Intensive Care Med 38:1573–1582. � PubMed Google Scholar Ashbaugh D, Bigelow DB, Petty T, Levine B (1967) Acute respiratory distress in adults. The Lancet 290:319–323. 67)90168-7Article Google Scholar Definition Task Force ARDS, Ranieri VM, Rubenfeld GD et al (2012) Acute respiratory distress syndrome: the Berlin Definition. JAMA 307:2526–2533. � CAS Google Scholar Bellani G, Pham T, Laffey J et al (2016) Incidence of acute respiratory distress syndrome-reply.
JAMA 316:347. � PubMed Google Scholar Tonetti T, Vasques F, Rapetti F et al (2017) Driving pressure and mechanical power: new targets for VILI prevention. Ann Transl Med 5:286. � PubMed Central PubMed Google Scholar Gattinoni L, Carlesso E, Caironi P (2012) Stress and strain within the lung. Curr Opin Crit Care 18:42–47. � PubMed Google Scholar Gattinoni L, Pesenti A (2005) The concept of "baby lung." Intensive Care Med 31:776–784.
� PubMed Google Scholar Guérin C, Albert RK, Beitler J et al (2020) Prone position in ARDS patients: why, when, how and for whom. Intensive Care Med 46:2385–2396. � CAS PubMed Central PubMed Google Scholar Guérin C, Reignier J, Richard J-C et al (2013) Prone positioning in severe acute respiratory distress syndrome. N Engl J Med 368:2159–2168. � CAS PubMed Google Scholar Gattinoni L, Carlesso E, Taccone P et al (2010) Prone positioning improves survival in severe ARDS: a pathophysiologic review and individual patient meta-analysis. Minerva Anestesiol 76:448–454CAS PubMed Google Scholar Brochard L, Slutsky A, Pesenti A (2017) Mechanical ventilation to minimize progression of lung injury in acute respiratory failure. Am J Respir Crit Care Med 195:438–442. � PubMed Google Scholar Combes A, Bréchot N, Luyt C-E, Schmidt M (2018) Indications for extracorporeal support: why do we need the results of the EOLIA trial? Med Klin Intensivmed Notfmed 113:21–25. � CAS PubMed Google Scholar
Fan E, Del Sorbo L, Goligher EC et al (2017) An Official American Thoracic Society/European Society of Intensive Care Medicine/Society of critical care medicine clinical practice guideline: mechanical ventilation in adult patients with acute respiratory distress syndrome. Am J Respir Crit Care Med 195:1253–1263. � PubMed Google Scholar Murray JF, Matthay MA, Luce JM, Flick MR (1988) An expanded definition of the adult respiratory distress syndrome. Am Rev Respir Dis 138:720–723. � CAS PubMed Google Scholar Bernard GR, Artigas A, Brigham KL et al (1994) Report of the American-European consensus conference on ARDS: definitions, mechanisms, relevant outcomes and clinical trial coordination. The Consensus Committee. Intensive Care Med 20:225–232. � CAS PubMed Google Scholar Riviello ED, Kiviri W, Twagirumugabe T et al (2016) Hospital incidence and outcomes of the acute respiratory distress syndrome using the Kigali modification of the berlin definition. Am J Respir Crit Care Med 193:52–59. � PubMed Google Scholar Matthay MA, Thompson BT, Ware LB (2021) The Berlin definition of acute respiratory distress syndrome: should patients receiving high-flow nasal oxygen be included? Lancet Respir Med 9:933–936. 21)00105-3Article CAS PubMed Central PubMed Google Scholar Ranieri VM, Tonetti T, Navalesi P et al (2022) High-flow nasal oxygen for severe hypoxemia: oxygenation response and outcome in patients with COVID-19. Am J Respir Crit Care Med 205:431–439. � CAS PubMed Google Scholar Rice TW, Wheeler AP, Bernard GR et al (2007) Comparison of the SpO2/FIO2 ratio and the PaO2/FIO2 ratio in patients with acute lung injury or ARDS. Chest 132:410–417. � PubMed Google Scholar Brown SM, Grissom CK, Moss M et al (2016) Nonlinear imputation of Pao2/Fio2 from Spo2/Fio2 among patients with acute respiratory distress syndrome. Chest 150:307–313. � PubMed Central PubMed Google Scholar Moss M, Ulysse CA, Angus DC, Heart N, Lung, and Blood Institute PETAL Clinical Trials Network (2019) early neuromuscular blockade in the acute respiratory distress syndrome. Reply N Engl J Med 381:787–788. � PubMed Google Scholar Wick KD, Matthay MA, Ware LB (2022) Pulse oximetry for the diagnosis and management of acute respiratory distress syndrome. Lancet Respir Med 10:1086–1098. 22)00058-3Article PubMed Central PubMed Google Scholar Meade MO, Cook RJ, Guyatt GH et al (2000) Interobserver variation in interpreting chest radiographs for the diagnosis of acute respiratory distress syndrome. Am J Respir Crit Care Med 161:85–90. � CAS PubMed Google Scholar Rubenfeld GD, Caldwell E, Granton J et al (1999) Interobserver variability in applying a radiographic definition for ARDS. Chest 116:1347–1353. � CAS PubMed Google Scholar Goddard SL, Rubenfeld GD, Manoharan V et al (2018) The randomized educational acute respiratory distress syndrome diagnosis study: a trial to improve the radiographic diagnosis of acute respiratory distress syndrome. Crit Care Med 46:743–748. � PubMed Google Scholar Khemani RG, Smith LS, Zimmerman JJ et al (2015) Pediatric acute respiratory distress syndrome: definition, incidence, and epidemiology: proceedings from the Pediatric Acute Lung Injury Consensus Conference. Pediatr Crit Care Med 16:S23-40. � PubMed Google Scholar Schenck EJ, Oromendia C, Torres LK et al (2019) Rapidly improving ARDS in therapeutic randomized controlled trials.
Chest 155:474–482. � CAS PubMed Google Scholar Villar J, Kacmarek RM, Pérez-Méndez L, Aguirre-Jaime A (2006) A high positive end-expiratory pressure, low tidal volume ventilatory strategy improves outcome in persistent acute respiratory distress syndrome: a randomized, controlled trial. Crit Care Med 34:1311–1318. � PubMed Google Scholar Villar J, Pérez-Méndez L, López J et al (2007) An early PEEP/FIO2 trial identifies different degrees of lung injury in patients with acute respiratory distress syndrome. Am J Respir Crit Care Med 176:795–804. � PubMed Google Scholar Ranieri VM, Rubenfeld G, Slutsky AS (2022) Rethinking ARDS after COVID-19. If a "better" definition is the answer, What is the question? Am J Respir Crit Care Med. � PubMed Central PubMed Google Scholar Calfee CS, Delucchi K, Parsons PE et al (2014) Subphenotypes in acute respiratory distress syndrome: latent class analysis of data from two randomised controlled trials. Lancet Respir Med 2:611–620. 14)70097-9Article PubMed Central PubMed Google Scholar Mrozek S, Jabaudon M, Jaber S et al (2016) Elevated plasma levels of sRAGE are associated with nonfocal CT-based lung imaging in patients with ARDS: a prospective multicenter study. Chest 150:998–1007. � PubMed Google Scholar Famous KR, Delucchi K, Ware LB et al (2017) Acute respiratory distress syndrome subphenotypes respond differently to randomized fluid management strategy. Am J Respir Crit Care Med 195:331–338. � CAS PubMed Central PubMed Google Scholar Bos LD, Schouten LR, van Vught LA et al (2017) Identification and validation of distinct biological phenotypes in patients with acute respiratory distress syndrome by cluster analysis. Thorax 72:876–883. � CAS PubMed Google Scholar Calfee CS, Delucchi KL, Sinha P et al (2018) ARDS subphenotypes and differential response to simvastatin: secondary analysis of a randomized controlled trial. Lancet Respir Med 6:691–698. 18)30177-2Article CAS PubMed Central PubMed Google Scholar Sinha P, Delucchi KL, Thompson BT et al (2018) Latent class analysis of ARDS subphenotypes: a secondary analysis of the statins for acutely injured lungs from sepsis (SAILS) study. Intensive Care Med 44:1859–1869. � CAS PubMed Central PubMed Google Scholar Delucchi K, Famous KR, Ware LB et al (2018) Stability of ARDS subphenotypes over time in two randomised controlled trials. Thorax 73:439–445. � PubMed Google Scholar Constantin J-M, Jabaudon M, Lefrant J-Y et al (2019) Personalised mechanical ventilation tailored to lung morphology versus low positive end-expiratory pressure for patients with acute respiratory distress syndrome in France (the LIVE study): a multicentre, single-blind, randomised controlled trial. Lancet Respir Med 7:870–880.
19)30138-9Article PubMed Google Scholar Bos LDJ, Scicluna BP, Ong DSY et al (2019) Understanding heterogeneity in biologic phenotypes of acute respiratory distress syndrome by leukocyte expression profiles. Am J Respir Crit Care Med 200:42–50. � CAS PubMed Google Scholar Sinha P, Delucchi KL, McAuley DF et al (2020) Development and validation of parsimonious algorithms to classify acute respiratory distress syndrome phenotypes: a secondary analysis of randomised controlled trials. Lancet Respir Med 8:247–257. 19)30369-8Article CAS PubMed Central PubMed Google Scholar Sinha P, Calfee CS, Cherian S et al (2020) Prevalence of phenotypes of acute respiratory distress syndrome in critically ill patients with COVID-19: a prospective observational study. Lancet Respir Med 8:1209–1218. 20)30366-0Article CAS PubMed Central PubMed Google Scholar Sinha P, Churpek MM, Calfee CS (2020) Machine learning classifier models can identify acute respiratory distress syndrome phenotypes using readily available clinical data. Am J Respir Crit Care Med 202:996–1004. � PubMed Central PubMed Google Scholar Zhang S, Lu Z, Wu Z et al (2020) Determination of a "Specific Population Who Could Benefit From Rosuvastatin": a secondary analysis of a randomized controlled trial to uncover the novel value of rosuvastatin for the precise treatment of ARDS. Front Med (Lausanne) 7:598621. � PubMed Google Scholar Wendel Garcia PD, Caccioppola A, Coppola S et al (2021) Latent class analysis to predict intensive care outcomes in Acute Respiratory Distress Syndrome: a proposal of two pulmonary phenotypes. Crit Care 25:154. � PubMed Central PubMed Google Scholar Bos LDJ, Sjoding M, Sinha P et al (2021) Longitudinal respiratory subphenotypes in patients with COVID-19-related acute respiratory distress syndrome: results from three observational cohorts. Lancet Respir Med 9:1377–1386. 21)00365-9Article CAS PubMed Central PubMed Google Scholar Sinha P, Furfaro D, Cummings MJ et al (2021) Latent class analysis reveals COVID-19-related acute respiratory distress syndrome subgroups with differential responses to corticosteroids. Am J Respir Crit Care Med 204:1274–1285. � CAS PubMed Central PubMed Google Scholar Liu X, Jiang Y, Jia X et al (2021) Identification of distinct clinical phenotypes of acute respiratory distress syndrome with differential responses to treatment. Crit Care 25:320. � PubMed Central PubMed Google Scholar Vasquez CR, Gupta S, Miano TA et al (2021) Identification of distinct clinical subphenotypes in critically Ill patients with COVID-19. Chest 160:929–943. CAS PubMed Google Scholar Ranjeva S, Pinciroli R, Hodell E et al (2021) Identifying clinical and biochemical phenotypes in acute respiratory distress syndrome secondary to coronavirus disease-2019. EClinicalMedicine 34:100829. � PubMed Central PubMed Google Scholar Lascarrou J-B, Gaultier A, Soumagne T et al (2021) Identifying clinical phenotypes in moderate to severe acute respiratory distress syndrome related to COVID-19: the COVADIS study. Front Med (Lausanne) 8:632933. � PubMed Google Scholar Xing C-Y, Gong W-B, Yang Y-N et al (2021) ARDS patients exhibiting a "Hyperinflammatory Anasarca" phenotype could benefit from a conservative fluid management strategy. Front Med (Lausanne) 8:727910. � PubMed Google Scholar Sinha P, Delucchi KL, Chen Y et al (2022) Latent class analysis-derived subphenotypes are generalisable to observational cohorts of acute respiratory distress syndrome: a prospective study. Thorax 77:13–21. � PubMed Google Scholar Hashem MD, Hopkins RO, Colantuoni E et al (2022) Six-month and 12-month patient outcomes based on inflammatory subphenotypes in sepsis-associated ARDS: secondary analysis of SAILS-ALTOS trial. Thorax 77:22–30. � PubMed Google Scholar Maddali MV, Churpek M, Pham T et al (2022) Validation and utility of ARDS subphenotypes identified by machine-learning models using clinical data: an observational, multicohort, retrospective analysis. Lancet Respir Med 10:367–377.
21)00461-6Article PubMed Central PubMed Google Scholar Duggal A, Kast R, Van Ark E et al (2022) Identification of acute respiratory distress syndrome subphenotypes de novo using routine clinical data: a retrospective analysis of ARDS clinical trials. BMJ Open 12:e053297. � PubMed Central PubMed Google Scholar Gattinoni L, Caironi P, Cressoni M et al (2006) Lung recruitment in patients with the acute respiratory distress syndrome. N Engl J Med 354:1775–1786. � CAS PubMed Google Scholar Nishimura M (2015) High-flow nasal cannula oxygen therapy in adults. J Intensive Care 3:15.
� PubMed Central PubMed Google Scholar Parke R, McGuinness S, Eccleston M (2009) Nasal high-flow therapy delivers low level positive airway pressure.
Br J Anaesth 103:886–890. � CAS PubMed Central PubMed Google Scholar Frat J-P, Thille AW, Mercat A et al (2015) High-flow oxygen through nasal cannula in acute hypoxemic respiratory failure.
N Engl J Med 372:2185–2196. � CAS PubMed Google Scholar Perkins GD, Ji C, Connolly BA et al (2022) Effect of noninvasive respiratory strategies on intubation or mortality among patients with acute hypoxemic respiratory failure and COVID-19: the RECOVERY-RS randomized clinical trial. JAMA 327:546–558. � CAS PubMed Central PubMed Google Scholar Azoulay E, Lemiale V, Mokart D et al (2018) Effect of high-flow nasal oxygen vs standard oxygen on 28-day mortality in immunocompromised patients with acute respiratory failure: the HIGH randomized clinical trial. JAMA 320:2099–2107. � CAS PubMed Central PubMed Google Scholar Ospina-Tascón GA, Calderón-Tapia LE, García AF et al (2021) Effect of high-flow oxygen therapy vs conventional oxygen therapy on invasive mechanical ventilation and clinical recovery in patients with severe COVID-19: a randomized clinical trial. JAMA 326:2161–2171. � CAS PubMed Google Scholar Frat J-P, Quenot J-P, Badie J et al (2022) Effect of high-flow nasal cannula oxygen vs standard oxygen therapy on mortality in patients with respiratory failure due to COVID-19: the SOHO-COVID randomized clinical trial. JAMA 328:1212–1222. � CAS PubMed Central PubMed Google Scholar Alptekinoğlu Mendil N, Temel Ş, Yüksel RC et al (2021) The use of high-flow nasal oxygen vs. standard oxygen therapy in hematological malignancy patients with acute respiratory failure in hematology wards.
Turk J Med Sci 51:1756–1763. � CAS PubMed Google Scholar Bouadma L, Mekontso-Dessap A, Burdet C et al (2022) High-dose dexamethasone and oxygen support strategies in intensive care unit patients with severe COVID-19 acute hypoxemic respiratory failure: the COVIDICUS randomized clinical trial. JAMA Intern Med 182:906–916. � CAS PubMed Central PubMed Google Scholar Prakash J, Bhattacharya PK, Yadav AK et al (2021) ROX index as a good predictor of high flow nasal cannula failure in COVID-19 patients with acute hypoxemic respiratory failure: A systematic review and meta-analysis.
J Crit Care 66:102–108. � CAS PubMed Central PubMed Google Scholar Azoulay E, Pickkers P, Soares M et al (2017) Acute hypoxemic respiratory failure in immunocompromised patients: the Efraim multinational prospective cohort study. Intensive Care Med 43:1808–1819. � PubMed Google Scholar Rochwerg B, Brochard L, Elliott MW et al (2017) Official ERS/ATS clinical practice guidelines: noninvasive ventilation for acute respiratory failure. Eur Respir J 50:1602426. � PubMed Google Scholar Wang D, Hu B, Hu C et al (2020) Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. JAMA 323:1061–1069. � CAS PubMed Central PubMed Google Scholar Alhazzani W, Møller MH, Arabi YM et al (2020) Surviving Sepsis Campaign: guidelines on the management of critically ill adults with Coronavirus Disease 2019 (COVID-19). Intensive Care Med 46:854–887. � PubMed Central PubMed Google Scholar COVID-19 Treatment Guidelines Panel. Coronavirus Disease 2019 (COVID-19) Treatment Guidelines. National Institutes of Health.
Available at BM, Qushmaq I, Al-Hameed FM et al (2019) Noninvasive ventilation in critically ill patients with the Middle East respiratory syndrome. Influenza Other Respir Viruses 13:382–390. � CAS PubMed Central PubMed Google Scholar Tran K, Cimon K, Severn M et al (2012) Aerosol generating procedures and risk of transmission of acute respiratory infections to healthcare workers: a systematic review. PLoS ONE 7:e35797.
� CAS PubMed Central PubMed Google Scholar Nair PR, Haritha D, Behera S et al (2021) Comparison of high-flow nasal cannula and noninvasive ventilation in acute hypoxemic respiratory failure due to severe COVID-19 pneumonia. Respir Care 66:1824–1830. � PubMed Google Scholar Grieco DL, Menga LS, Cesarano M et al (2021) Effect of helmet noninvasive ventilation vs high-flow nasal oxygen on days free of respiratory support in patients with COVID-19 and moderate to severe hypoxemic respiratory failure: the HENIVOT randomized clinical trial. JAMA 325:1731–1743. � CAS PubMed Google Scholar Coudroy R, Frat J-P, Ehrmann S et al (2022) High-flow nasal oxygen alone or alternating with non-invasive ventilation in critically ill immunocompromised patients with acute respiratory failure: a randomised controlled trial. Lancet Respir Med 10:641–649. 22)00096-0Article CAS PubMed Google Scholar Munshi L, Mancebo J, Brochard LJ (2022) Noninvasive respiratory support for adults with acute respiratory failure. N Engl J Med 387:1688–1698. � PubMed Google Scholar Brambilla AM, Aliberti S, Prina E et al (2014) Helmet CPAP vs. oxygen therapy in severe hypoxemic respiratory failure due to pneumonia. Intensive Care Med 40:942–949. � CAS PubMed Google Scholar Squadrone V, Massaia M, Bruno B et al (2010) Early CPAP prevents evolution of acute lung injury in patients with hematologic malignancy. Intensive Care Med 36:1666–1674. � PubMed Google Scholar Cosentini R, Brambilla AM, Aliberti S et al (2010) Helmet continuous positive airway pressure vs oxygen therapy to improve oxygenation in community-acquired pneumonia: a randomized, controlled trial. Chest 138:114–120. � PubMed Google Scholar Lemiale V, Mokart D, Resche-Rigon M et al (2015) Effect of noninvasive ventilation vs oxygen therapy on mortality among immunocompromised patients with acute respiratory failure: a randomized clinical trial. JAMA 314:1711–1719. � CAS PubMed Google Scholar He H, Sun B, Liang L et al (2019) A multicenter RCT of noninvasive ventilation in pneumonia-induced early mild acute respiratory distress syndrome. Crit Care 23:300. � PubMed Central PubMed Google Scholar Zhan Q, Sun B, Liang L et al (2012) Early use of noninvasive positive pressure ventilation for acute lung injury: a multicenter randomized controlled trial. Crit Care Med 40:455–460. � PubMed Google Scholar Hilbert G, Gruson D, Vargas F et al (2001) Noninvasive ventilation in immunosuppressed patients with pulmonary infiltrates, fever, and acute respiratory failure. N Engl J Med 344:481–487. � CAS PubMed Google Scholar Wermke M, Schiemanck S, Höffken G et al (2012) Respiratory failure in patients undergoing allogeneic hematopoietic SCT–a randomized trial on early non-invasive ventilation based on standard care hematology wards.
Bone Marrow Transplant 47:574–580. � CAS PubMed Google Scholar Yoshida T, Amato MBP, Kavanagh BP, Fujino Y (2019) Impact of spontaneous breathing during mechanical ventilation in acute respiratory distress syndrome.
Curr Opin Crit Care 25:192–198.
� PubMed Google Scholar Antonelli M, Conti G, Pelosi P et al (2002) New treatment of acute hypoxemic respiratory failure: noninvasive pressure support ventilation delivered by helmet–a pilot controlled trial. Crit Care Med 30:602–608. � PubMed Google Scholar Antonelli M, Pennisi MA, Pelosi P et al (2004) Noninvasive positive pressure ventilation using a helmet in patients with acute exacerb