

Effective Treatment for Covid-19. Buy and keep them in Stock.
Do your Research also. Talk to a doctor about how to take these medicines.

Take Following Medicines as soon as symptoms of Covid-19 are there.

For Adults* without Severe Comorbid Conditions:

Antivirals to Control Covid-19 Virus:

1. Tablet Ivermectin - 1 sheet.

Dosage: Total Dosage per Day – 12 mg for 60 KG weight.
Adjust to your weight
Period: **Day 1 to Until Total Recovery**
When to Take: Morning 0 –Lunch 1 - Evening 0.
Take After Food.

2. Tablet. Fabiflu 400 mg – 1 box (17 tablets)

Dosage: Total Dosage per Day – 1600 mg
Period: **Day 1 to 6**
When to Take: Morning 2 –Lunch 0 - Evening 2
Take After Food

3. MNB's Cellular Armor - Ultimate Immunity Booster

Buy www.hfnlife.com or www.amazon.in

Dosage: **Mix 80 ml in 1000 ml water**
Period: **Day 1- Until Total Recovery**
When to Take: 5 times, 200 ml each time
Take 30 minutes Before Food

4. Tablets Vitamin C, D, & Zinc Tablets. Dosage per day

Dosage: Vitamin C (2000 mg per day)
Vitamin D3 (10000 I.U per day)
Zinc (60 mg) per day
Period: **Day 1- Until Total Recovery + 10 Days**
When to Take: Take After Food

If symptoms of Cough, Difficulty in Breathing, Cold etc.
Antibacterial to Control lung Infection:

5. MNB's Cellular Armor - Respiratory Relief Liquid.
Buy at www.hfnlife.com or www.amazon.in

Dosage: **Mix 100 ml in 100 ml of water** in the steamer
Period: **Day 1 - Until Total Recovery**
When to Take: Steam 4-6 times day

6. Tablet Doxycycline 100 mg 1 sheet (Day 2 to 4, 1-0-1).

Dosage: **Total Dosage per Day – 100 mg**
Period: **Day 1 of Lung Related Issues**
When to Take: Morning 1 –Lunch 0 - Evening 0
Take After Food

If Fever is there

7. Tablet Dolo 650 (only when fever is there)

If Phlegm is there

8. Syrup. Jvtuss LS 1 bottle. Only when cough with phlegm (mucus)

Dosage: Total Dosage per Day – 15 ml
Period: **Until Phlegm is gone**
When to Take: Morning 5 ml –Lunch 5 ml - Evening 5 ml
Take With or without Food

Or

Mucinac 600mg

Dosage: Total Dosage per Day – 600 mg
Period: **Until Phlegm is gone**
When to Take: Add the tablet in half glass of water and allow it to dissolve completely, and drink 1 hour before food.

If Breathing Difficulty and Oxygen Saturation is Falling down

Steroids (these should NOT be taken unless prescribed by a doctor, not to be taken beyond 5 days. Excessive use will lead to prolonged health risks and also Mucormycosis issues.)

9. Tablet Dexamethasone or Tablet Medrol or Tablet Goutnill – 1 sheet

Dosage: Total Dosage per Day – 4 mg
Period: **When Severe Lung Related Issues are present**
When to Take: Morning 1 –Lunch 0 - Evening 1
Take After Food

Allergy Medicines (Antihistamines)

10. Tablet Peritol Tablet (Cyproheptadine) - 1 sheet

Dosage:	Total Dosage per Day – 12 mg
Period:	From Day 8 – Until Lung Related Issues resolved
When to Take:	Morning 1 –Lunch 1 - Evening 1 Take After Food

* Exclusion Criteria:

- Pregnancy
- Patients with pre-existing terminal condition with life expectancy < 6 months
- Patient with clinically frailty according to a score of the clinical frailty scale equal or superior to 7
- Patients with pre-existing severe lung disease requiring home oxygen therapy.
- Patients with pre-existing severe hepatic cirrhosis (Grade C according to the Child-Pugh classification)
- Patients with pre-existing kidney failure (GFR strictly less than 15ml/min/1.73m² according The KDIGO classification) or requiring renal replacement therapy.
- Patients with pre-existing angle-closure glaucoma
- Patient with symptomatic prostatic hypertrophy or bladder neck obstruction
- Patient with history of seizure disorder
- Patient with history of adverse reaction to antihistamines or to Cyproheptadine
- Patients taking routinely SSRI or monoamine oxidase inhibitor therapy.
- Patients presenting severe hepatic cytolysis with ALP >5 ULN at the time of study inclusion.