

GENERAL FEED CHART

The fertilizer dosage (low or high) is chosen based on the soil type and nutrient content, measured by the Electrical Conductivity (EC). In soils with a high EC, a low dosage is recommended to avoid over-fertilization. As a preventive measure, it is suggested to apply nutrient irrigation for three cycles followed by one with only water at a balanced pH in the fourth cycle. This guideline may vary depending on the specific nutrient concentration in the soil. Continuous root flushes will help prevent any nutrient excess.

	VEG WEEK 1	VEG WEEK 2	VEG WEEK 3	VEG WEEK 4	FLOWER WEEK 1	FLOWER WEEK 2	FLOWER WEEK 3	FLOWER WEEK 4	FLOWER WEEK 5	FLOWER WEEK 6	FLOWER WEEK 7	FLOWER WEEK 8	FLOWER WEEK 9
BASE	5 ML/G	7 ML/G	7 ML/G	7 ML/G	5 ML/G	7 ML/G	7 ML/G	7 ML/G	7 ML/G	7 ML/G	7 ML/G	5 ML/G	5 ML/G
GROW	10 ML/G	14 ML/G	14 ML/G	14 ML/G	-	-	-	-	-	-	-	-	-
BLOOM	-	-	-	-	10 ML/G	14 ML/G	14 ML/G	14 ML/G	14 ML/G	14 ML/G	14 ML/G	10 ML/G	10 ML/G
GORILLAGROW	FOLIAR	FOLIAR	FOLIAR	FOLIAR	-	-	-	-	-	-	-	-	-
CALIMAGICIAN	0.5 GR/G	0.5 GR/G	0.5 GR/G	-	-	-	-	-	-	-	-	-	-
BASILIC	4 ML/G	4 ML/G	4 ML/G	4 ML/G	4 ML/G	4 ML/G	4 ML/G	4 ML/G	4 ML/G	4 ML/G	-	-	-
SEAGOLD	-	-	4 ML/G	4 ML/G	4 ML/G	4 ML/G	4 ML/G	4 ML/G	4 ML/G	4 ML/G	4 ML/G	4 ML/G	4 ML/G
CANDY-KOLA	-	-	-	1 GR/G	1 GR/G	1 GR/G	1 GR/G	1 GR/G	1 GR/G	1 GR/G	1 GR/G	-	-
DRAGONBUD	-	-	-	-	FOLIAR	FOLIAR	FOLIAR	1 GR/G	1.5 GR/G	2 GR/G	2.5 GR/G	-	-

NOTE: If you want to deliver a nutrient solution with lower electroconductivity, you can apply the Basilic product foliarly.

Precautions: Wear eye/face protection and avoid eating, drinking, or smoking during product use. After handling, ensure thorough handwashing.

First Aid: In case of eye contact, rinse cautiously with water, remove contact lenses if applicable, and seek medical attention if irritation persists. If swallowed, rinse mouth and consult a doctor if feeling unwell.

*Dosage is grams per gallon.

*Mix a new nutrient solution weekly.

*Recommended pH: 5.8 -6.5

*Flush recommended in the last week before harvest.

*Wear eye face protection.

WWW.HARVESTCONUTRIENTS.COM



FOLIAR FEED CHART

A proper foliar application is done 1 to 3 times per week, with a complete week of rest between applications. During the vegetative stage, it's important to cover the entire plant, including stems and leaves. Similarly, in the early flowering stages, it's advisable to apply throughout the entire plant. After the third week of flowering, it's recommended to avoid direct application on the flowers and focus on stems and leaves. It's best to refrain from foliar applications after the sixth or seventh week of flowering.

	VEG WEEK 1	VEG WEEK 2	VEG WEEK 3	VEG WEEK 4	FLOWER WEEK 1	FLOWER WEEK 2	FLOWER WEEK 3	FLOWER WEEK 4	FLOWER WEEK 5	FLOWER WEEK 6	FLOWER WEEK 7	FLOWER WEEK 8	FLOWER WEEK 9
GORILLAGROW	1-1.5 GR/G	1-1.5 GR/G	1.5-3.5 GR/G	1.5-3.5 GR/G	-	-	-	-	-	-	-	-	-
DRAGONBUD	-	-	-	-	1.5 GR/G	1.5 GR/G	2.5 GR/G	-	-	-	-	-	-
SEAGOLD	4 ML/G	4 ML/G	4 ML/G	4 ML/G	4 ML/G	4 ML/G	4 ML/G	4 ML/G	-	-	-	-	-
BASILIC	3 ML/G	3 ML/G	3 ML/G	3 ML/G	3 ML/G	3 ML/G	3 ML/G	3 ML/G	-	-	-	-	-

Avoid using with basic metal sulfates, strongly alkaline materials (such as hydrated lime and lime sulfur), or sprays containing oils.

A compatibility test is recommended if the desired combination has not been previously used. Apply the spray solution to a small test area before widespread application to determine any potential undesirable phytotoxic effects.

First Aid: In case of eye contact, rinse cautiously with water, remove contact lenses if applicable, and seek medical attention if irritation persists. If swallowed, rinse mouth and consult a doctor if feeling unwell.

Precautions: Wear eye/face protection and avoid eating, drinking, or smoking during product use. After handling, ensure thorough handwashing.

*Dosage is grams per gallon. *Mix a new nutrient solution weekly. *Recommended pH: 5.8 -6.5

*Flush recommended in the last week before harvest. *Wear eye face protection.

WWW.HARVESTCONUTRIENTS.COM

