How do I know if my dog has **Blastomycosis?**

Has your dog been exposed to regions of the country **where blasto frequently occurs?**



United States: common areas highlighted in orange

Has your dog been in environments that may harbor fungal growth?



Examples include:

-]. Soil
- 2. Excavated land
- 3. Riverways



Any concerns?

VISIT OUR WEBSITE: https://miravistavets.com/

Blastomycosis

a fungal infection caused by breathing in spores from the fungus Blastomyces

in **Dogs**







Common signs


Check for signs

- weight loss
- fever
- skin or eye lesions
- seizures
- trouble breathingharsh cough
- lameness
- IOI I ICI IC22
- depression

Consider susceptibility

 most common in large, young, sporting breeds including:
Coonhounds, Pointers, Weimaraners, Dobermans and Retrievers

*but also occurs in all breeds



if this sounds familiar...





Visit your vet

Your vet may recommend

- labwork
- X-rays
- biopsy



Diagnosis

Receive results

- accurate
- fast



• within 24-68 hours of receipt of specimen



Treatment

Duration: months to years

one or more antifungal medications

possible ongoing xrays and lab tests

keep physical activity low to reduce stress

Pronosis: excellent with early diagnosis and treatment

^{*}Signs can be evident in a period of days to months after first contact with disease