

# Musculoskeletal Pearls

(and Pet Peeves)



Dennis K-Borna, M.D.  
Family & Sports Medicine, Kaiser Fontana

No disclosures to report


PRIMARY CARE HAWAII CONFERENCE

April 1-5, 2024

Kauai, Hawaii



# X-ray Basics




- Can't describe a fracture without at least an AP and lateral
- Consider oblique view if defect can only be seen in one view
- Consider comparison view when dealing with growth plate injuries
- Weightbearing views can be more helpful when evaluating joint spaces
- Order knee x-rays under order (KP Health Connect)

73562BC




# Sprain vs Strain



## Sprain


- Involves injury of a ligament
- Typically a result of a joint being taken beyond its normal range of motion
- Repetitive injuries are uncommon
- Often worse acutely, but improve over time
- Initial treatment with rest/immobilization, followed by range of motion
- Typically graded 1-3



## Strain


- Involves injury of a muscle or tendon
- Can arise from overuse\*
- May not be as severe initially, but can last multiple months
- Treatment with rest and activity modifications, followed by passive and eccentric stretching, and then strengthening

# Sprain vs Strain



## Sprain


- Involves injury of a ligament
- Typically a result of a joint being taken beyond its normal range of motion
- Repetitive injuries are uncommon
- Often worse acutely, but improve over time
- Initial treatment with rest/immobilization, followed by range of motion
- Typically graded 1-3



## Strain

- Involves injury of a muscle or tendon
- Can arise from overuse\*
- May not be as severe initially, but can last multiple months
- Treatment with rest and activity modifications, followed by passive and eccentric stretching, and then strengthening

# Injury Treatment




- Bring tissues together for healing

- Achilles/calf strain

- Shorten the gastroc-Achilles complex


- Heel lift to the height that resolves pain



- Finger joint sprains

- Biceps/triceps


- Quads/Hamstrings



- Immobilize the joint above and below a fracture

- When in doubt, splint and bring them back in a week


# Pediatric Injury Treatment



- Ligamentous injuries are rare in children because the ligaments are stronger than the growth plate
- Be careful about diagnosing a “sprain” in young kids
- Soft tissue injuries heal rapidly in kids
  - If a “sprain” or “strain” is still painful after a week, strongly suspect a fracture
- When in doubt, splint and bring them back in a week


# Growth Plate Fractures

## Salter-Harris Classification



# Growth Plate Fractures


## Salter-Harris Classification







# Growth Plate Fractures

Salter-Harris Classification




# Tendinosis



normal tendon

tendinopathy

# Tendinosis




normal tendon



tendinopathy


# Eccentric Stretching Exercises



- The hallmark of effective therapy for tendinosis and strains
- Tension in the musculotendinous unit as it is stretched




# NSAIDs




- While many NSAIDs are similar, there are unique differences among them that can be tailored to treating specific issues
- Anti-inflammatory effects at higher doses (prescription)
- Some have stronger anti-inflammatory properties, and some have more analgesic effects
  - GI side effects increased with COX-1 inhibition
- A short burst or a course of an NSAID can be helpful depending on the acuity and severity of the condition
  - “3 ibuprofen 3 times a day for 3 days”
  - Long-acting NSAID x 7-10 days

# NSAIDs




NSAID	Anti-inflammatory Effects	Analgesic Effects	GI Side Effects
→ Ibuprofen*	Moderate to High	Moderate to High	Moderate to High
→ Naproxen*	Moderate to High	Moderate to High	Moderate to High
→ Diclofenac	High	High	High
→ Indomethacin	Very High	Moderate to High	High
→ Ketorolac	High	High	High
Celecoxib	Moderate to High	Moderate to High	Lower
→ Meloxicam	Moderate to High	Moderate to High	Moderate
Piroxicam	High	High	High
→ Etodolac	Moderate to High	Moderate to High	Moderate
Sulindac	Moderate to High	Moderate to High	Moderate
↗ Nabumetone	Moderate	Moderate	Lower
↘ Oxaprozin	Moderate	Moderate	Moderate
Salsalate	Moderate	Moderate	Lower


# Acetaminophen vs NSAIDs



- Although not for inflammation, acetaminophen can improve pain symptoms “close to” NSAIDs
  - NNT **4.6** (acetaminophen) vs **3** (ibuprofen)
- For milder chronic pain, regular use of acetaminophen with NSAIDs for flare-ups may be appropriate
  - NNT for acetaminophen + ibuprofen **1.6**
- While rare, hepatotoxicity with regular daily use is lower for acetaminophen than for NSAIDs
  - Diclofenac and sulindac can carry relatively increased hepatic risks
  - For pain control in cirrhotic patients, a lower daily dose of acetaminophen (2g) is regarded as safer than NSAIDs



# Ketorolac (Toradol)




- Can be a very useful medication for acute pain/inflammation
- Can be especially effective for renal colic
- Recent studies have shown minimal differences between 15mg, 30mg, and 60mg IM doses






# Skeletal Muscle Relaxants




- Widely prescribed, but with limited evidence on efficacy
- Can have high adverse effect profile (CNS effects)
- Often used along with NSAIDS for many musculoskeletal issues
  - Acute back pain
  - Neck pain
  - Myofascial pain conditions
  - 🤔 Strains
  - 🤔 Tendinitis
  - Insomnia




# Skeletal “Muscle Relaxants”




Cyclobenzaprine



Amitriptyline




Nortriptyline



“Flexeril 10mg tid”

# Skeletal Muscle Relaxants



- **Antispasticity agents:** work on the spinal cord or directly on the skeletal muscles to improve muscle hypertonicity and involuntary spasms
- **Antispasmodic agents:** decrease muscle spasms through changes in CNS conduction






Figure 1 Classification of Skeletal Muscle Relaxants

# Skeletal Muscle Relaxants



Medication	Spasticity	Pain	Sedation	Duration
→ Cyclobenzaprine	Moderate	High	High	4-6 hours
→ Baclofen	High	Moderate	Moderate to high	4-8 hours
Tizanidine	Moderate to High	Moderate	Moderate to high	4-6 hours
→ Methocarbamol	Low to moderate	Low to moderate	Low to moderate	3-6 hours
Diazepam	Moderate to High	Low to moderate	High	6-12 hours
Metaxalone	Low to moderate	Low to moderate	Low to moderate	4-6 hours
Carisoprodol	Low to moderate	Low to moderate	High	4-6 hours

# Back



## Anterior Back

- Discs
- Vertebral bodies




## Posterior Back

- Central canal
- Facet joints
- Pars interarticularis
- Paraspinous muscles\*
- Sacroiliac joints



# Back



## Anterior Back



## Posterior Back

- Worse with sitting/leaning forward


- Worse with standing/leaning backward

- Work on anterior stretching and posterior strengthening

- Work on anterior strengthening and posterior stretching



# .pispasm



For muscle spasms of the neck and back, there are several things that are important to help relieve your symptoms sooner rather than later:

1. Medications -- this includes anti-inflammatory pain medications with or without muscle relaxers
2. Heat to the affected area -- usually for about 30 minutes or so
3. Stretching exercises -- these should be done slowly, and should involve stretching AWAY from the area that hurts
4. Massage -- start gently, and then gradually increase the pressure and the size of the massage circles.
5. Drink plenty of fluids -- dehydrated muscles tend to be more spastic than well-hydrated muscles.

The best way to approach these treatments is in the following order:

Take the medication(s) and wait an hour.


Then apply the heat for 30 minutes.

Then do the stretching exercises and the massaging. This gives you a better chance to stretch out and relax the affected muscles.

Avoid excessive bedrest and sedentary positions, as well as highly exertional activities, heavy lifting, or frequent bending or other exacerbating activities.




# Prednisone




- Corticosteroids for acute injuries can inhibit the healing process and are usually **not** recommended
- Systemic corticosteroids for acute back pain
  - No clear benefit for non-radicular low back pain or spinal stenosis
  - May be helpful for acute radiculopathy only
  - Do not reduce likelihood of surgery for slipped/bulging disc
- Systemic corticosteroids for carpal tunnel syndrome
  - Not recommended
- Systemic corticosteroids can be helpful for acute gout and Bell's palsy
- Multiple joint pains; diagnostic for presumed rheumatologic issue




# Ankle Injuries




- For acute ankle injuries, pay close attention to the lateral view



# Ankle -- Mortise View




# Ankle Sprains




- Don't forget to check the Achilles
- Most causes of recurrent ankle issues are related to instability
- Stable ankle sprains:
  1. RICE
  2. ROM exercises
    - “the alphabet”
  3. Strengthening exercises
    - towel or theraband exercises
  - \*4. Weightbearing strengthening/proprioceptive exercises
    - “tiptoes of one foot for 15 seconds”

# Ankle Stirrup Splint



Bad Sprain or  
Ankle Fracture

# Posterior Splint



Nondisplaced  
stable foot  
fractures or  
foot sprains

# Cam Walker (Cam Boot)




Stable foot or  
ankle injury

# Braces

- Ankle sleeve
  - Mild support, compression for swelling
- Lace-up ankle brace
  - Provides additional support to an unstable ankle upon return to activities
  - Similar support to ankle taping
  - Can result in chronic instability/weakness if worn excessively, especially without appropriate rehab
- Heel lifts
  - Achilles tendinitis or calf strains



# Shoulder




- Many diagnoses strongly correlate with certain ages
  - Under 18: growth plate (epiphysitis)
  - 18-25: shoulder instability
  - 20-30: AC joint pain\* (if pain is superior)
  - 35-50: shoulder impingement, rotator cuff syndrome
  - 40-60: small to moderate rotator cuff tears
  - 60+: large rotator cuff tears
  - 45-65: frozen shoulder






# Shoulder -- Epiphysitis



- Proximal humeral epiphysitis very common in young throwers
- Pain in the proximal shoulder, especially with throwing or resisted activities
- Exam can be similar to rotator cuff injury
- X-rays helpful to confirm open growth plate
  - Often normal, but may show growth plate widening (epiphysiolysis)

# Shoulder -- Epiphysitis




Normal shoulder




Abnormal shoulder  
widening of the growth plate

# Shoulder -- Epiphysitis



- Treatment is rest from throwing activities and from exacerbating activities for 1-3 months
- Encourage pain-free activities
- Be careful about ordering physical therapy!
- Counsel about “random sharp pains”
- Re-check at 4-6 week intervals
- Gradual return to throwing once exam is completely pain-free

# Shoulder -- Instability



<https://www.dubaisportsorthopaedics.com>

- The glenohumeral joint is inherently unstable


- Stabilizers:

- Ligamentous capsule
- Labrum
- Rotator cuff


- The rotator cuff acts like a suction cup to help stabilize the joint

- Weak cuff means more shifting movement in the joint

# Shoulder -- Instability




## Internal rotator strengthening exercise




© Healthwise, Incorporated

## External rotator strengthening exercise




© Healthwise, Incorporated

## Scapular exercise: Retraction




© Healthwise, Incorporated


# Shoulder – AC Joint



- Involved in many shoulder movements, so AC injury or pain will yield multiple “positive” exam findings
  - Check for AC tenderness
  - Cross-body extension
  - Scarf test
  - Pain at the ends of abduction
  - Pain with shoulder compression



# Shoulder – AC Joint




● AC injury (“separated shoulder”)




● AC arthritis



● AC pain is common in young people who do a lot of bench press and shoulder press (military press)




# Shoulder – Rotator Cuff Syndrome/Impingement



- The rotator cuff undergoes age-related degeneration
- Pain is often proximal lateral
  - “It’s never the deltoid”
  - Can radiate to the elbow, but not past it
- Similar to instability, RC strengthening is needed
  - Other considerations:
    - Degree of cuff pathology
    - Pain limitations
- Scapula mechanics are important









# Shoulder – Rotator Cuff Tears

- Large rotator cuff tears compromise the ability of the glenohumeral joint to articulate properly
- Deltoid pulls the humeral head superiorly
  - “high-riding humerus”




# Shoulder – Adhesive Capsulitis



- Early frozen shoulder can mimic a bad impingement syndrome
- Time course is 1-3 years (average 18 months)
  - Freezing stage
  - Frozen stage
  - Thawing stage
- More common in females, diabetics, and thyroid patients
  - Peak age: mid-50s
- Look at restricted passive external rotation for diagnosis
- Aggressive ROM therapy early on is important to help shorten the course


# Knee



- Acute knee injury with pop, immediate swelling, and an inability to finish playing is ACL tear 80% of the time
- For acute knee injuries, limit to weightbearing as tolerated only
- Check ability to do a straight leg raise to ensure that extensor mechanism is intact
- Knee immobilizer
  - Keeps the knee in extension
  - OK for extensor mechanism injury
  - For other injuries, ace wrap or compression brace with crutches may be better
  - Can promote weakness, quad atrophy within a few days



# Knee – Patellofemoral Syndrome



- “Runner’s knee”
- Most common cause of anterior knee pain
- Caused by irritation of the patellofemoral compartment
  - Dysfunctional patellar tracking
  - Malalignment
  - Trauma
- Chondromalacia patella is often used interchangeably, but it technically also involves a wearing down of the cartilaginous surfaces




# Patellofemoral Pain Syndrome Treatment


- Strengthening the anteromedial knee
- Strengthening the hip
- Stretching of the IT band
- Arch supports to help with malalignment
- Patella stabilizing brace
- Surgery (rare)




# Patellofemoral Pain Syndrome Treatment



# Patellofemoral Pain Syndrome Treatment







# Pain Threshold vs Pain Tolerance

## ● Pain threshold:

- The minimum level at which a stimulus is considered painful
- Some psychologic components, but more **physiologically** influenced
  - Genetics
  - Age
  - Exercise
  - Opiates

## ● Pain tolerance:

- The maximum level of pain that can be endured
- **Psychologic** factors can have a greater influence
  - Attitudes
  - Depression/anxiety

# Thank you!



# References

Abril L, Zamora C, Cordero M, Williams AR, Friedman BW. The Relative Efficacy of Seven Skeletal Muscle Relaxants. An Analysis of Data From Randomized Studies. *J Emerg Med*. 2022 Apr;62(4):455-461. doi: 10.1016/j.jemermed.2021.09.025. Epub 2022 Jan 20. PMID: 35067395.

Alghamdi Y, Morya R, Bahathiq D, Bokhari A, Alaboud A, Abdulhamid A, Ghaddaf A, Jamjoom M. Comparison of acetaminophen, ketamine, or ketorolac versus morphine in the treatment of acute renal colic: A network meta-analysis. *The American Journal of Emergency Medicine*, Volume 73, 2023, Pages 187-196, ISSN 0735-6757.

Chou R, Peterson K, Helfand M. Comparative efficacy and safety of skeletal muscle relaxants for spasticity and musculoskeletal conditions: a systematic review. *Journal of Pain and Symptom Management*, Volume 28, Issue 2, 2004, Pages 140-175, ISSN 0885-3924.

Chou R, Pinto RZ, Fu R, Lowe RA, Henschke N, McAuley JH, Dana T. Systemic corticosteroids for radicular and non-radicular low back pain. *Cochrane Database of Systematic Reviews* 2022, Issue 10. Art. No.: CD012450. DOI: 10.1002/14651858.CD012450.pub2.

Click Physiotherapy. Heel lifts for Achilles Tendonitis – evidence based or fad? Retrieved February 4, 2024 from <https://www.clickphysiotherapy.com.au/heel-lifts-for-achilles-tendinopathy-evidence-based-or-fad/>

Coghill, R. C., & Eisenach, J. (2003). Individual Differences in Pain Sensitivity: Implications for Treatment Decisions. *Anesthesiology*, 98(6), 1312-1314.

Dvorin EL, Ebell MH. Short-Term Systemic Corticosteroids: Appropriate Use in Primary Care. *Am Fam Physician*. 2020 Jan 15;101(2):89-94. PMID: 31939645.

# References

Motov S, Yasavolian M, Likourezos A, Pushkar I, Hossain R, Drapkin J, Cohen V, Filk N, Smith A, Huang F, Rockoff B, Homel P, Fromm C. Comparison of Intravenous Ketorolac at Three Single-Dose Regimens for Treating Acute Pain in the Emergency Department: A Randomized Controlled Trial. *Ann Emerg Med*. 2017 Aug;70(2):177-184. doi: 10.1016/j.annemergmed.2016.10.014. Epub 2016 Dec 16. PMID: 27993418.

Nathaniel J. Turner, Drew A. Long, Joseph R. Bongiorno, Timothy P. Katoski, Lisa M. Jin, John Paul Horsch, Brian J. Ahern. Comparing two doses of intramuscular ketorolac for treatment of acute musculoskeletal pain in a military emergency department, *The American Journal of Emergency Medicine*, Volume 50, 2021, Pages 142-147, ISSN 0735-6757.

Nichols, Andrew W MD. Complications Associated With the Use of Corticosteroids in the Treatment of Athletic Injuries. *Clinical Journal of Sport Medicine* 15(5):p E370, September 2005. | DOI: 10.1097/01.jsm.0000179233.17885.18

Sachs CJ. Oral analgesics for acute nonspecific pain. *Am Fam Physician*. 2005 Mar 1;71(5):913-8. PMID: 15768621.

See S, Ginzburg R. Choosing a skeletal muscle relaxant. *Am Fam Physician*. 2008 Aug 1;78(3):365-70. PMID: 18711953.

Witenko C, Moorman-Li R, Motycka C, Duane K, Hincapie-Castillo J, Leonard P, Valaer C. Considerations for the appropriate use of skeletal muscle relaxants for the management of acute low back pain. *P T*. 2014 Jun;39(6):427-35. PMID: 25050056; PMCID: PMC4103716.

Xu, Y. and Murrell, G.A., 2008. The basic science of tendinopathy. *Clinical orthopaedics and related research*, 466, pp.1528-1538.