

A tennis racket is shown in a close-up, slightly blurred view, with a green tennis ball resting on the strings. The racket's frame is blue and yellow. The background is a mix of green and blue, suggesting an outdoor setting like a tennis court.

The Masters Athlete

Tricks to Keep Your Aging Patients Moving

Jordan D. Metzl, MD

Hospital for Special Surgery


- I have no conflicts of interest to disclose

Today's Lecture

- Demographics
- Preparing for Exercise in the Masters Athlete
- Orthopedic Issues for the Masters Athlete
- Treatment Plans for the Masters Athlete in Your Practice







GARATACHEA ET AL.

REJUVENATION RESEARCH
Volume 18, Number 1, 2015
© Mary Ann Liebert, Inc.
DOI: 10.1089/rej.2014.1623


Who Is the Masters Athlete?

- A master athlete is an athlete who competes in a particular sport or discipline at a high level of proficiency, usually after the **age of 35!** (What?!)



USA Trends

6.9% of the population in 2000 to a projected 19.3% by 2050





Population Research Bureau, 2022

Sports With Masters Divisions



- National Senior Games > 10,000
- Cross Fit
- Ball/Racquet sports
- Running/Triathlon
- Spartan
- Strongman
- Boxing
- Body Building

Aging and Elite-Level Sports Performance



Overall Improvement in Athletic Performance

Comparison of 1896 Olympic winning times in running events and current Masters records that surpass those winning times (from ESPN and World Masters Records)

Running events	1896 Olympic winning time (from the first Olympic games in Athens)	Current age-group records that surpass 1896 Olympic times and age at which these records were achieved
100 m (s)	12.0	11.7 (61 years)
200 m (s)	22.2	22.1 (46 years)
400 m (s)	54.2	53.9 (63 years)
800 m (min:s)	2:11.0	2:10.4 (60 years)
1500 m (min:s)	4:33:2	4:27:7 (60 years)
Marathon (h:min:s)	2:58:50	2:54:5 (73 years)

Preparing the Masters Athlete for Sport



Recommendations for Preparticipation Screening and the Assessment of Cardiovascular Disease in Masters Athletes

An Advisory for Healthcare Professionals From the Working Groups of the World Heart Federation, the International Federation of Sports Medicine, and the American Heart Association Committee on Exercise, Cardiac Rehabilitation, and Prevention

Barry J. Maron, Claudio Gil S. Araújo, Paul D. Thompson, Gerald F. Fletcher, Antonio Bayés de Luna, Jerome L. Fleg, Antonio Pelliccia, Gary J. Balady, Francesco Furlanello, Steven P. Van Camp, Roberto Elosua, Bernard R. Chaitman and Terry L. Bazzarre

- Risk for sudden cardiac death in the range of **1:15,000 joggers** per year or **1:50,000** participants in marathons, with a marked predominance of **deaths in men**
- Young high school and college-aged athletes has been calculated to be \approx 1:200 000 to 1:300 000 per academic year
- ECG for athletes > 45
- ECG + Exercise Testing with symptoms or - Men > 45 and Women > 55 with: hypercholesterolemia, dyslipidemia (total cholesterol >200 mg/dL; elevated low-density lipoprotein [LDL] cholesterol [$>$ 130 mg/dL]; systemic hypertension (systolic blood pressure $>$ 140 mm Hg or diastolic pressure $>$ 90 mm Hg); current or recent cigarette smoking; diabetes mellitus (fasting plasma glucose \geq 126 mg/dL or treatment with insulin or oral hypoglycemics); history of myocardial infarction or sudden cardiac death in a first-degree relative $<$ 60 years old

Cardiovascular Risk Assessment in the Older Athlete

- In adult athletes ≥ 35 years old, CAD is the primary cause of major adverse cardiovascular events.
- A CAC scan quantifies the burden of calcified atherosclerotic plaque and provides important prognostic value.
- A comprehensive risk assessment for CVD in older athletes includes a review of conventional risk markers, physical examination, 12-lead ECG, and CAC score.
- 10-year risk of ASCVD should be estimated for adult athletes using a validated risk calculator such as the ACC/AHA, Astro-CHARM, or MESA calculators.
- CAC scoring or risk-enhancing factors such as hsCRP and Lp(a) may be used to reclassify or revise the risk assessment.
- Studies of endurance athletes not only demonstrate higher prevalence of CAC compared with nonathletes but also suggest coronary plaque composition in athletes is more stable and may not confer the same risk.
- Lifestyle interventions are recommended to optimize cardiovascular health.
- Blood pressure should be optimally controlled to reduce CVD risk.
- Statin therapy is recommended if 10-year risk of ASCVD is $\geq 10\%$ and considered if risk is $\geq 7.5\%$.
- Low-dose aspirin 75-100 mg daily can be considered for athletes with high ASCVD risk and low bleeding risk.
- Stress testing is recommended for athletes with symptoms concerning for ischemic heart disease.
- Stress testing may be considered for asymptomatic athletes with high ASCVD risk (eg, markedly elevated CAC score ≥ 400).
- Echocardiography or cardiac MRI is recommended for athletes with physical examination or ECG abnormalities concerning for valvular heart disease or cardiomyopathy.

Preparing for Competition: Masters Athlete Screening Study

- 5-year prospective screening study (Canada)
- Checking for CVD and MACE (major adverse cardiac events)
- 798 athletes, all sports, screened annually
- Anthropometrics, blood pressure, resting electrocardiogram, modified American Heart Association 14-element recommendations, cardiovascular event questionnaire, physical examination
- Abnormal screen according to the European Association of Cardiovascular Prevention and Canadian Cardiology Society Guidelines underwent further evaluations




Preparing for Competition: Masters Athlete Screening Study

- 798 masters athletes (62.7% male, 54.6±9.5 years) were screened; 91 (11.4%) of the cohort were found to have CVD. CAD was the most common diagnosis (69.2%)
- During the following four years, there were an additional 89 CVD diagnoses with an incidence rate of 3.58/100, 4.14/100, 3.74/100, 1.19/100, for years two to five, respectively.
- A total of 10 MACE occurred (two cardiovascular deaths, five myocardial infarctions and three cerebrovascular accidents). All events occurred in male athletes (63.6±12.5 years).
- Yearly cardiovascular screening of masters athletes identified ~3 new diagnoses per 100 athletes per year. Ten MACE occurred despite yearly screening and high CV fitness of masters athletes.




MSK Screening for Masters Athletes

- Skeletal muscle is 40% of body mass
- Sarcopenia has both functional and metabolic implications
- Resistance training increases muscle mass, strength, and fiber hypertrophy
- Mitochondrial density and maximal oxygen uptake increases with endurance training




Age-related Sarcopenia


- The average human will lose 40% of their muscle mass between the ages of 30-60.
- After 60, sarcopenia progresses at 40% loss/decade




Modified WHO/HPS, Geneva 2000




Age-Related Sarcopenia




40-year-old triathlete



74-year-old sedentary man




70-year-old triathlete




Strength Loss with Aging

- Loss of muscle fibers occurs with aging beyond 50 (Falkner, 2009)
- Repetitive strength training encourages muscle hypertrophy at any age (Pearson, 2022)
- Type II muscle fiber loss > Type I with aging (Khadeserian, 1998)




DOI: 10.1097/JSM.0b013e3181845f1c, 1998

Muscle Fiber Types

	Type I fibers	Type II a fibers	Type II x fibers	Type II b fibers
Contraction time	Slow	Moderately Fast	Fast	Very fast
Size of motor neuron	Small	Medium	Large	Very large
Resistance to fatigue	High	Fairly high	Intermediate	Low
Activity Used for	Aerobic	Long-term anaerobic	Short-term anaerobic	Short-term anaerobic
Maximum duration of use	Hours	<30 minutes	<5 minutes	<1 minute
Power produced	Low	Medium	High	Very high
Mitochondrial density	High	High	Medium	Low
Capillary density	High	Intermediate	Low	Low
Oxidative capacity	High	High	Intermediate	Low
Glycolytic capacity	Low	High	High	High
Major storage fuel	Triglycerides	Creatine phosphate, glycogen	Creatine phosphate, glycogen	Creatine phosphate, glycogen
Myosin heavy chain, human genes	MYH7	MYH2	MYH1	MYH4 

Common MSK Issues in the Masters Athlete

- Muscle/Tendon
 - Tendonitis
 - Tendonosis
 - Partial Tear
 - Rupture
- Ligament
- Bone
 - Osteoarthritis
 - Bone Insufficiency



Principles for Assessment of MSK Injury in Masters Athlete

- Evaluate locally, think globally
- Make sure you aren't creating a bigger problem (rest=rust)
- Beware MRI findings
- Consider cause and prevention strategies



Don't Miss the Forrest for the Trees

**What is Your
Goal and
Role in
Primary
Care?**





Physical Activity Guidelines 2023




Fitness Resources

- Cardio (2-3x/week)
- Strength (2-3x/week)
- Balance (2-3x/week)
 - In-Person
 - Virtual



What About Exercise Intensity?

Energy Systems





Name: _____ Date: _____

Aerobic Activity

Type: Walk Run Swim Bike Other _____

Frequency (days/week): 2 3 4 5 6 7

Intensity: Light Moderate Vigorous
(A Casual Walk) (A Brisk Walk) (Jogging or Running)

Time (minutes/day): 10 20 30 60 60 or more

Steps/day: 2,500 5,000 7,500 10,000 10,000 or more

Strength Training

- Muscle strengthening should be done at least two days per week
- Exercise should be done to strengthen all major muscle groups: legs, hips, back, chest, abdomen, shoulder, arms
- For each exercise, 8-12 repetitions should be completed
- Examples: push-ups, pull-ups, sit-ups, and lunges

Physician Signature: _____



Physiologic Capacity and Aging


Oxygen consumption and its determinants at maximal exercise in endurance exercise-trained men

	Young men (28 years)	Older men (60 years)	Age-related change (%)
Oxygen consumption ($\text{ml kg}^{-1} \text{min}^{-1}$)	68.2	49.4	28
Cardiac output (l min^{-1})	27.0	21.7	20
Stroke volume (ml beat^{-1})	147	132	10
Heart rate (beats min^{-1})	184	165	10
a-v O_2 difference (ml (100 ml)^{-1})	16.7	15.2	8

The data were compiled from four studies in which values for all of the variables were reported in groups of young and older groups ([Grimby *et al.* 1966](#); [Hagberg *et al.* 1985](#); [Rivera *et al.* 1989](#); [Ogawa *et al.* 1992](#)).

Aging and Physiologic Capacity

Exercise economy: the steady-state oxygen consumption while exercising at a specific submaximal exercise intensity below the lactate threshold




The ability to sustain a high fraction of one's maximal oxygen consumption during submaximal exercise

Maximal oxygen consumption, the upper limit of energy production

Lactate Threshold




- The point during exercise of increasing intensity at which lactate levels begin to rise disproportionately
- Occurs at 50% of VO_2Max in untrained people
- Can be increased to 85-90% with training
- Maximum sustainable workload





Raising Lactate Threshold

- Accumulate volume at maximal steady state as well as high intensity intervals
- **Push it up**
 - Tempo workouts
 - “comfortably hard”
- **Pull it up**
 - High intensity Intervals



Maximum VO₂

- VO₂ Max = *the maximum amount of oxygen able to be used during exhaustive exercise*



$$VO_2 = CO \times (a-vO_2)$$

Maximum VO_2 Values

- 30-39 year old, average
 - Male = 42ml/kg/min
 - Female = 37 ml/kg/min
- 40-49 year old, average
 - Male = 40ml/kg/min
 - Female = 34ml/kg/min
- 50-59 year old, average
 - Male = 37ml/kg/min
 - Female = 31ml/kg/min
- 70-79 year old, average
 - Male = 31 ml/kg/min
 - Female = 28 ml/kg/min
- Heart Transplant candidates
 - ≤ 14 ml/kg/min



Elite endurance athletes
60-80ml/kg/min

What is HIIT?



0	Rest
1	Really Easy
2	Easy
3	Moderate
4	Sort of Hard
5	Hard
6	
7	Really Hard
8	
9	Really, Really, Hard

HIIT = 90-95% of Peak Heart Rate for short bursts of activity (5-60 seconds) [ACSM – 2019]

- Peak Heart Rate – 220 – age
- *or*
- $211 - 0.64 \times \text{Age}$

Example: $(211 - (.64 \times 50)) = 179$

Max HR = $179 \times .9 = 161$

Who can HIIT It?

- All ages
- All shapes and sizes
- All athletes have different physical and metabolic limitations
- Can evolve over time




HIIT in Older Adults


- HIIT increases VO_2 max in subjects > 65 y/o (Marriot 2021)
- HIIT maintains cardiopulmonary fitness in subjects 45-65 (Alzar 2022)
- HIIT maintains functional movement in subjects > 65 (Stern 2023)
- HIIT more effective than moderate intensity exercise in adults > 65 (Marriot 2021)



Heart Rate & Lactate vs. Power (Watts)




Medical Benefits of HIIT




Cellular Response to Intensity Training

- 12 week study
- 36 men and 36 women
- Young = 18-30
- Old = 65-80
- Group 1 - High Intensity Biking
- Group 2- Strength Training
- Group 3 – Strength + Interval Training
- Muscle biopsies pre/post – (assessed lean muscle mass + insulin sensitivity)




Cellular Benefits of HIIT




- HIIT and combined training improved aerobic capacity and skeletal muscle mitochondrial respiration
- High-intensity interval training improved age-related decline in muscle mitochondria
- HIIT reversed many age-related differences in of mitochondrial proteins in concert with increased mitochondrial protein synthesis

Oxidative Metabolism across Exercise Intensities




adapted, modified from Tucker, Ross, and Dugas *The Runner's Body* (2009) Rodale

Moderate Intensity “Zone 2”




adapted, modified from Tucker, Ross, and Dugas *The Runner's Body* (2009) Rodale

Vigorous Intensity



adapted, modified from Tucker, Ross, and Dugas *The Runner's Body* (2009) Rodale

High Intensity





Friends

Different Intensities → Different Adaptations

Light → Moderate → Vigorous → Maximal



“Manipulation of different exercise training variables promotes specific & diverse mitochondrial adaptations”

Granata C, Jamnick NA, Bishop DJ. Training-Induced Changes in Mitochondrial Content and Respiratory Function in Human Skeletal Muscle. *Sports Med.* 2018 Aug;48(8):1809-1828. doi: 10.1007/s40279-018-0936-y. PMID: 29934848.

Exercise Summary – The Masters Athlete

- Daily Movement
- Strength 3x/week
- Cardio 3x/week
- Balance 2-3x/week
- Consistency





Compliance



Fitness Compliance: Overview

- **Demographic and Biological Factors**
(Men>Women, Thin>Obese)
- **Self-Esteem** (Higher SE, Higher Compliance)
- **Social and Cultural Support, Incentive**
- **Physical Environment**



What Motivates You?

1. External Motivation
2. Internal Motivation

What Makes People Fitness Compliant?

Fun

Community

Results

Goals



Factors Generating Internal Motivation



- Community
- Competency
- Results
- Enjoyment

Coon D. *Gateways to Mind and Behavior*, 2010

Enjoyment During Exercise Mediates Adherence

- 45 random participants, avg age 46 y/o, women>men 6:1
- The fun group – enjoyment focused training:
 - 1) Group>Solo
 - 2) Group decision making>dictating
 - 3) Positive feedback = achievement
 - 4) Intensity regulation
 - 5) Workout announced in advance (transparency)
 - 6) Training diversity (different forms)

Enjoyment During Exercise Mediates Adherence

- Control Group – goal to improve fitness
- Emphasis on fitness, no social cues
- Parameters:
 - 1) cardiorespiratory exercise
 - 2) resistance exercise
 - 3) flexibility exercise
 - 4) neuromotor exercise

Enjoyment During Exercise Mediates Adherence

Table 1. Means and standard deviations for PACES and exercise frequency.

		<i>M</i>	<i>SD</i>	<i>t</i>	<i>df</i>	<i>p</i>
1. Session PACES	IG	68.9	4.8	-0.9	39	0.37
	CG	70.2	4.5			
4. Session PACES	IG	71.1	1.9	6.9	39	<0.01
	CG	66.2	2.6			
8. Session PACES	IG	71.3	1.8	6.7	24.0	<0.01
	CG	65.8	3.0			
Exercise Frequency	IG	7.9	0.3	2.6	17.0	0.02
	CG	7.2	1.3			

Note: *M* = Mean; *SD* = standard deviation; *t* = value of the t-distribution; *df* = degrees of freedom; *p* = probability value; PACES = Physical Activity Enjoyment Scale; IG = intervention group; CG = control group.

The Fun Factor



What Makes People Fitness Compliant?

Fun

Community

Results


Goals



The Pleasure and Displeasure People Feel When they Exercise at Different Intensities



Decennial Update and Progress towards a Tripartite Rationale for Exercise Intensity Prescription

Panteleimon Ekkekakis,¹ Gaynor Parfitt² and Steven J. Petruzzello³



The Communal Effect of Exercise Adherence





Aral. Nature, 2017

What Makes People Fitness Compliant?

Fun

Community

Results

Goals



The Optimal Fitness Program

- Cardio
- Strength
- Flexibility/Balance

- *Exercise Snacks*



What Makes People Fitness Compliant?

Fun

Community

Results

Goals



Goals **ERVEZA**
PACIFICO
CLARA



Clinical Recommendations

SORT: Strength of Recommendation Taxonomy

A: consistent, good-quality patient-oriented evidence

B: inconsistent or limited-quality patient-oriented evidence

C: consensus, disease-oriented evidence, usual practice, expert opinion, or case series

Clinical Recommendation	SORT Evidence Rating
Preparticipation screening for cardiovascular disease should include a standard 12-lead electrocardiogram (ECG) for all masters athletes older than 40 years. ECG exercise testing should be performed for older athletes with 1 or more cardiac risk factors, including diabetes, a family history of myocardial infarction, or dyslipidemia, among others. ¹⁷	B
Adults and masters athletes alike should engage in moderate to vigorous aerobic exercise for 20- to 30-minute sessions 3 to 5 days each week, as well as resistance, flexibility, and balance training. ¹¹	B
Generally, masters athletes should avoid returning to high-impact sports after total joint arthroplasty but may return to low-impact sports. ^{14,33} In the setting of hip resurfacing, however, return to high-impact activities may be permissible. ²¹	B
Treatment for musculoskeletal injuries in the masters athlete should include rehabilitation and analgesia, such as nonsteroidal anti-inflammatory drugs (NSAIDs), when appropriate. ^{10,19} NSAIDs should be used with caution given the gastrointestinal and cardiovascular risks. ¹⁹	B

Training Hacks for the Masters Athlete

BY Simon Ward

Aging is inevitable, but it doesn't have to be the end of your athletic career. Here are some training adjustments to consider as you age.



Training Peaks, 2023

HSS

Lift Heavy

Muscle mass starts to decline in the mid- to late thirties. It occurs at a rate of around 3-5% per decade and speeds up once we hit the mid- to late 50s, to about 7-8% per decade. This mainly affects the fast-twitch fibers, which is why an athlete tends to lose power and speed rather than endurance.

Sadly, no amount of swimming, cycling or running will help preserve fast-twitch fibers because endurance activities generally don't provide sufficient load for that type of muscle adaptation. Instead of doubling down on the long slow distance (LSD), Dr. Stacy Sims suggests LHS (“lifting heavy sh**t”) as you age to preserve muscle mass. This means exercises like deadlifts, squats, leg presses, and seated or bent-over rows with a heavy (for you) load and a low number of repetitions. Dr. Sims' work (and her excellent book ROAR) focuses on female athletes of all ages, but the concepts apply to men as well.

Many aging athletes feel that lifting in such a way may actually cause them injury, and of course, if you have a poor lifting technique, then that might happen. But if you have good technique, then heavy work in the gym can have a number of wonderful benefits. Maximal strength can improve, and with it, power. Heavy weights can also lead to improved resilience in connective tissues, which will be a big help in the battle to stay injury-free. At the very least, one can slow down that loss of muscle mass—and in some circumstances, you may even gain new muscle.

Maintain Range of Motion

With age, our joints tighten up, and the range of movement (ROM) around a joint or series of joints can be reduced. This has implications for all three triathlon disciplines, but especially swimming and running. If velocity is measured by limb frequency and length of stroke or stride, a loss of ROM (coupled with that loss of speed/power mentioned earlier) ultimately results in a slowdown.

Decreased mobility also increases your risk of injury. Any athlete should aim to avoid an injury, but it's even more important in your older years, as any lost fitness is harder to regain. To increase your mobility, Dr. Kelly Starrett recommends doing at least 15 minutes of mobility work for every 60 minutes of training each week. For example, if you aim for 10 hours of swimming, cycling, and running, then your minimum would be 2.5 hours.

Don't Skimp on Intensity

As you age, long, slow distances (even for long-distance triathletes) should become less of a priority, and high-intensity training must occupy the forefront of your mind. Like with heavy lifting, this can seem counterintuitive to older athletes, but trust me, it does work, and if you are healthy and injury-free, then you have absolutely nothing to fear.

To retain or build in regular high-intensity interval workouts, aim to work at an RPE of 9-10 for around 10% of your total weekly duration. Of course, this will depend upon your training history and attention to other recovery factors.

You may want to be cautious about fast running, especially if you have a history of calf or Achilles problems, but on the bike and in the pool, there should be nothing to hold you back. Work at the same percentage above FTP or CSS that a younger athlete might. Quality is the key, so don't be shy about taking a longer recovery interval between repetitions if you feel you need it.

Listen to Your Body


You have gained some athletic wisdom over the years, so you should use it! Recovery between workouts also requires adjustment with age. Maybe you could have done a track workout and a Vo2 max bike workout in 48 hours in your thirties, but you might need 72 or even 96 hours as you enter your 60s. Mountain bike hall-of-famer and XTERRA athlete Ned Overend once said, “I can still do what I always did, it just takes me longer to get it done.”

The bottom line is that you have to be more compassionate to your body. You’ve been using it hard for 30, 40, or even 50 years, and it has served you well. Now is the time to repay that service with some kindness. Look after those aching joints, tired muscles and well-used heart.

Listen to your body and if you’re ever in doubt about a workout or your health on that day, take it easy, or rest. As an older, wiser athlete you have hopefully learned to control your ego a bit and can be comfortable finding the path for long-term consistency.

At some point, we all must accept that we are no longer going to get faster or more powerful—but avoiding the slowdown as much as possible can be just as exciting a goal! By lifting heavy weights, maintaining intensity, and looking after your mobility and recovery, you’ll be healthy and performing well into your later years.

91



Thank You



Questions and
Discussion

References

- McTiernan A, Friedenreich CM, Katzmarzyk PT, et al. Physical activity in cancer prevention and survival: A systematic review. *Medicine and Science in Sports and Exercise* 2019; 51(6):1252-1261.
- Rezendes LFM, Sá TH, Markozannes G, et al. Physical activity and cancer: an umbrella review of the literature including 22 major anatomical sites and 770 000 cancer cases. *British Journal of Sports Medicine* 2018; 52(13):826-833.
- Patel AV, Friedenreich CM, Moore SC, et al. American College of Sports Medicine Roundtable Report on physical activity, sedentary behavior, and cancer prevention and control. *Medicine and Science in Sports and Exercise* 2019; 51(11):2391-2402.
- Keimling M, Behrens G, Schmid D, Jochem C, Leitzmann MF. The association between physical activity and bladder cancer: systematic review and meta-analysis. *British Journal of Cancer* 2014; 110(7):1862-1870.
- Moore SC, Lee IM, Weiderpass E, et al. Association of leisure-time physical activity with risk of 26 types of cancer in 1.44 million adults. *JAMA Internal Medicine* 2016; 176(6):816-825.
- Pizot C, Boniol M, Mullie P, et al. Physical activity, hormone replacement therapy and breast cancer risk: A meta-analysis of prospective studies. *European Journal of Cancer* 2016; 52:138-154.
- Hardefeldt PJ, Penninkilampi R, Edirimanne S, Eslick GD. Physical activity and weight loss reduce the risk of breast cancer: A meta-analysis of 139 prospective and retrospective studies. *Clinical Breast Cancer* 2018; 18(4):e601-e612.
- Eliassen AH, Hankinson SE, Rosner B, Holmes MD, Willett WC. Physical activity and risk of breast cancer among postmenopausal women. *Archives of Internal Medicine* 2010; 170(19):1758-1764.
- Fournier A, Dos Santos G, Guillas G, et al. Recent recreational physical activity and breast cancer risk in postmenopausal women in the E3N cohort. *Cancer Epidemiology, Biomarkers & Prevention* 2014; 23(9):1893-1902.
- Liu L, Shi Y, Li T, et al. Leisure time physical activity and cancer risk: evaluation of the WHO's recommendation based on 126 high-quality epidemiological studies. *British Journal of Sports Medicine* 2016; 50(6):372-378.
- Schmid D, Behrens G, Keimling M, et al. A systematic review and meta-analysis of physical activity and endometrial cancer risk. *European Journal of Epidemiology* 2015; 30(5):397-412.
- Du M, Kraft P, Eliassen AH, et al. Physical activity and risk of endometrial adenocarcinoma in the Nurses' Health Study. *International Journal of Cancer* 2014; 134(11):2707-2716.
- Friedenreich C, Cust A, Lahmann PH, et al. Physical activity and risk of endometrial cancer: The European prospective investigation into cancer and nutrition. *International Journal of Cancer* 2007; 121(2):347-355.
- Borch KB, Weiderpass E, Braaten T, et al. Physical activity and risk of endometrial cancer in the Norwegian Women and Cancer (NOWAC) study. *International Journal of Cancer* 2017; 140(8):1809-1818.
- Behrens G, Jochem C, Keimling M, et al. The association between physical activity and gastroesophageal cancer: systematic review and meta-analysis. *European Journal of Epidemiology* 2014; 29(3):151-170.