Induction of Ferroptosis Protocol

Rev.1.1, October, 2023

By Mark Simon, Director, Nutritional Oncology Research Institute, CEO, NORI Nutraceuticals

What is Ferroptosis?

Ferroptosis is a form of programmed or regulated cell death different from apoptosis. Ferroptosis is an iron mediated mechanism that causes catastrophic damage to the mitochondrial membrane and other cellular lipid membranes. This is also known as lipid peroxidation chain reactions. Cancer cells are far more sensitive to ferroptosis than normal cells because of a multitude of factors. The primary factor is an elevated level of oxidative stress (ROS) compared to normal cells. Another is an accumulation of intracellular iron.

Ferroptosis is initiated by a Fenton reaction between hydrogen peroxide and ferrous iron which produces hydroxyl radical. The reaction proceeds further ultimately resulting a chain reaction which causes irreversible damage to phospholipid cell membranes. This is called lipid peroxidation.



Conditions Required to Trigger Ferroptosis Selectively in Cancer Cells

There is a set of conditions necessary to initiate and sustain ferroptosis in cancer cells. All of these conditions can be met with diet, lifestyle and nutraceuticals. Off-label drugs that support ferroptosis may be incorporated if available through an integrative oncologist.

- 1. Intracellular Free Iron
- 2. Lowered Antioxidant Defense
- 3. High Ratio of Polyunsaturated to Monosaturated Fatty Acids (PUFA/MUFA).
- 4. Minimized Hypoxia

Ferroptosis Induction Protocol

Dietary Methionine and Cysteine Restriction

The primary goal of methionine and cysteine restriction is to limit glutathione synthesis. Glutathione is a tripeptide synthesized from cysteine, glycine and glutamate. Cancer cells are highly dependent on glutathione for regulation of oxidation (ROS).

The diet offers many side benefits which help to lower inflammation, suppress angiogenesis, lower IGF-1, stabilize glucose and insulin and shift the tumor microenvironment towards an alkaline pH.

Dietary Polyunsaturated and Mono-saturated Fatty Acid Balance

Overall fat intake should be minimized. Use hempseed oil and borage oil to increase n-3 and n-6 polyunsaturated fatty acids (PUFAs). Minimize mono-saturated fatty acids(MUFAs) such as those found in olive oil, avocado and canola oil.

Iron Overload - Ferric Ammonium Citrate

A sufficient amount of intracellular free iron (ferrous) must be present for the initiation of ferroptosis. Cancer cells exhibit altered iron metabolism and excessive iron storage by ferritin.

Studies have demonstrated that exogenous iron alone can induce ferroptosis or augment other ferroptosis inducers. Ferric iron is converted to ferrous iron which is the form of iron that reacts with hydrogen peroxide.

Ferric Ammonium Citrate is a highly water soluble form of iron. The recommended dosing schedule is one 25 mg sublingual tablet 2 times a day for a total of 50 mg of elemental iron per day.

Thioredoxin Reductase Inhibition - Sodium Selenite, Piperlongumine

Sodium Selenite and Piperlongumine are both inhibitors of the thioredoxin antioxidant system. Both inhibit the thioredoxin reductase enzyme which blocks recycling of reduced thioredoxin.

Sodium Selenite dosage is 0.3 mg/kg using a combination of sublingual and slow release oral tablets.

Piperlongumine is administered as a whole herb preparation of Piper Longum fruit and root parts. Dosage is 4 tablets 4 times a day.

Exclusion of all Antioxidant Supplements

Antioxidants in food can not be controlled but concentrated vegetable juices should be avoided. Juicing may be counterproductive in this regard.

Do not take any supplements containing Vitamin C, Vitamin E tocopherol, NAC, Vitamin A, Polyphenols

Exclusion of all Iron Chelators and Iron Sequestors

Curcumin

Overcoming Hypoxia

Regular exercise is recommended to improve oxygenation of tissues and improvement is blood flow. Hypoxic areas of a tumor may be more resistant to ferroptosis compared to non-hypoxic areas of a tumor.

References

Lei, G., Zhuang, L. & Gan, B. Targeting ferroptosis as a vulnerability in cancer. *Nat Rev Cancer* **22**, 381–396 (2022). <u>https://doi.org/10.1038/s41568-022-00459-0</u>

Zhang, C., Liu, X., Jin, S. *et al.* Ferroptosis in cancer therapy: a novel approach to reversing drug resistance. *Mol Cancer* **21**, 47 (2022). <u>https://doi.org/10.1186/s12943-022-01530-y</u>

Karthikeyan Subburayan, Faisal Thayyullathil, Siraj Pallichankandy, Anees Rahman Cheratta, Sehamuddin Galadari, Superoxide-mediated ferroptosis in human cancer cells induced by sodium selenite, Translational Oncology,Volume 13, Issue 11, 2020

Chen Xin, Yu Chunhua, Kang Rui, Tang Daolin, Iron Metabolism in Ferroptosis, Frontiers in Cell and Developmental Biology, Vol. 8, 2020

Jiaojiao Li & Wei Zhang (2023) From iron chelation to overload as a therapeutic strategy to induce ferroptosis in hematologic malignancies, Hematology, 27:1, 1163-1170, DOI: <u>10.1080/16078454.2022.2132362</u>

Mortensen MS, Ruiz J, Watts JL. Polyunsaturated Fatty Acids Drive Lipid Peroxidation during Ferroptosis. Cells. 2023 Mar 4;12(5):804. doi: 10.3390/cells12050804. PMID: 36899940; PMCID: PMC10001165.

For Further Information visit <u>www.nutritionaloncology.net</u> and join the Ferroptosis Support Group on Facebook <u>https://</u> <u>www.facebook.com/groups/1771229133327428</u>