

EXAMPLE SCENARIO

E0651 MEDICARE (Traditional) TIMELINE & DOCUMENTS in this Order- Timeline with dates in this order & documentation requirements

MEASUREMENTS signed by doctor

1. **MEASUREMENT #1 – 3/1/25**
Baseline measurement
 - Key is 4-week trial of conservative therapy with measurements taken before and after trial period.
 - Rx can not be written before the trial period is done.
2. **MEASUREMENT #2 – 4/1/25**
Unresponsive with elevation, exercise, and compression garments, at least 4 weeks

MEDICAL DOCUMENTS (signed by doctor) dated on or after 4/1/25:

1. **RX- 4/1/25 - 4 Week Trial of Conservative Therapy must be documented before prescribing.**
2. **LMN - 4/1/25 - On or before RX Date.**
3. **QSD: "Office notes for Compression" - 4/1/25 (QSD= Qualifying Supporting Document)**
4. **Medical Assessment and Agreement Statement - 4/1/25 - On or before RX Date.**
5. **DOCTOR'S F2F (Face to Face) - On or before 4/1/25, 6 month time frame for Face to Face.**
6. **DOCTOR'S Medical Records – Must support the Medical Necessity and substantiate the information provided on the "FORMS"**

DELIVERY DOCUMENTS (signed by patient):

1. **DELIVERY TICKET - 4/1/25 (or later)**
2. **CONTRAINDICATIONS WITH PRE AND POST TRIAL MEASUREMENTS 4/1/25 (or later):**
 - a. **PRE-TRIAL MEASUREMENT #3 at delivery - 4/1/25 (or later)**
 - b. **POST-TRIAL MEASUREMENT #4 at delivery – 4/1/25 (or later)**

STATEMENT NEEDED IN DOCTOR'S MEDICAL RECORD: "SAMPLE"

The chronic lymphedema has been unresponsive with at least **4 weeks of elevation, exercise, and compression garments** used on both legs. Detailed measurements have been taken over time to confirm that the chronic lymphedema continues to persist.

Initial treatment with the PCD (E0651) was completed with bilateral leg sleeves @ 50 mmHg without any pain or discomfort. John Doe was able to operate the PCD, apply & remove the leg sleeves and connect all tubing without any assistance.

Ordering PCD (E0651) & Bilateral leg sleeves for home use. Pressure: 50 mmHg, Duration: 60 minutes, Frequency: 2x's daily, for treatment of persistent lymphedema.

Follow up in 4-6 weeks to review pressure, duration & frequency. Retake BLE measurements, adjust settings as needed to treat/control persistent lymphedema.