

CLINICAL WRITING

Here are some examples of clinical statements.

Mood & Affect

- Client presented with euthymic mood and congruent affect throughout the session.
- Client presented with anxious mood and constricted affect.
- Client demonstrated increased emotional distress when discussing identified stressors.
- Client's affect appeared blunted but appropriate to the content discussed.
- Client reported increased depressive symptoms since the previous session.
- Client demonstrated improved emotional regulation compared to previous sessions.
- Client appeared tearful when discussing interpersonal and/or trauma-related concerns.

Processing & Exploration

- Therapist facilitated exploration of the client's thoughts, emotions, and behavioral responses related to ____.
- Client processed the impact of ____ on current functioning.
- Therapist supported client in identifying connections between past experiences and present-day responses.
- Client explored underlying beliefs associated with ____.
- Therapist utilized open-ended questions to facilitate insight and self-reflection.
- Client identified recurring patterns related to ____.
- Client explored the meaning they have assigned to ____.
- Therapist supported client in examining alternative perspectives.
- Client demonstrated increased insight into the relationship between thoughts, emotions, and behaviors.
- Client processed unresolved emotions related to ____.
- Client explored the impact of ____ on interpersonal relationships.
- Therapist provided space for client to process and verbalize previously unprocessed emotions.

Engagement & Participation

- Client was engaged and participatory throughout the session.
- Client demonstrated appropriate engagement with therapeutic interventions.
- Client was receptive to clinician feedback and redirection.
- Client required minimal prompting to participate in therapeutic discussion.
- Client demonstrated increased willingness to explore underlying emotional concerns.
- Client was guarded initially but became increasingly engaged as rapport was established.
- Client demonstrated limited engagement when discussing emotionally distressing topics.

Coping & Emotional Regulation

- Client identified adaptive coping strategies currently utilized to manage emotional distress.
- Client reported increased difficulty utilizing coping skills during periods of heightened stress.
- Client practiced grounding techniques during session with clinician support.
- Client demonstrated improved ability to identify emotional triggers and corresponding responses.
- Client continues to develop healthy emotion-regulation strategies.
- Client reported utilizing avoidance as a primary coping mechanism.
- Client demonstrated increased capacity to tolerate and process uncomfortable emotions.

CBT Statements

- Therapist utilized CBT interventions to identify maladaptive thought patterns.
- Client identified automatic thoughts associated with ____.
- Therapist assisted client in examining evidence supporting and challenging the identified thought.
- Client explored cognitive distortions contributing to emotional distress.
- Therapist facilitated cognitive restructuring.
- Client practiced developing a more balanced alternative thought.
- Client demonstrated increased ability to recognize the connection between cognition and emotional response.
- Therapist challenged all-or-nothing thinking in a supportive manner.
- Client identified patterns of catastrophizing related to ____.
- Client explored how assumptions and interpretations contribute to emotional distress.
- Client practiced separating thoughts from factual evidence.

CLINICAL WRITING

Here are some examples of clinical statements.

DBT / Emotion Regulation

- Therapist provided psychoeducation regarding emotion regulation.
- Client identified emotional triggers associated with ____.
- Therapist assisted client in identifying early signs of emotional escalation.
- Client practiced identifying and labeling emotions.
- Therapist reinforced use of distress-tolerance strategies.
- Client identified adaptive coping strategies for managing emotional distress.
- Therapist reviewed grounding techniques.
- Client practiced mindfulness skills during session.
- Client demonstrated increased awareness of physiological cues associated with emotional activation.
- Therapist supported client in identifying factors that contribute to vulnerability to emotional dysregulation.
- Client explored the difference between emotional urges and behavioral choices.
- Client practiced utilizing coping skills while discussing a distressing topic.

Trauma-Informed Language

- Therapist utilized a trauma-informed approach throughout the session.
- Therapist maintained a focus on emotional safety and stabilization.
- Client explored the impact of previous traumatic experiences on current functioning.
- Therapist provided psychoeducation regarding trauma responses.
- Client identified trauma-related triggers.
- Client explored patterns of hyperarousal and/or avoidance.
- Therapist normalized the client's responses within the context of their reported experiences.
- Client explored the relationship between past experiences and current beliefs about safety.
- Therapist supported client in identifying present-day cues that activate trauma-related responses.
- Client demonstrated increased awareness of trauma-related patterns.
- Therapist reinforced the distinction between past and present experiences.
- Client practiced grounding while discussing trauma-related material.
- Therapist monitored client activation and adjusted interventions to support regulation.

Parts Work / Internal Systems

- Therapist utilized a parts-informed approach to facilitate exploration of internal experiences.
- Client identified conflicting internal experiences related to ____.
- Client explored the protective function of a previously identified part.
- Therapist supported client in approaching internal experiences with curiosity rather than judgment.
- Client identified how different parts respond to perceived threat or emotional distress.
- Client explored the needs underlying a protective response.
- Therapist facilitated increased self-compassion toward previously criticized aspects of self.
- Client demonstrated increased ability to differentiate between internal experiences and present circumstances.

Grief & Loss

- Client processed emotions associated with grief and loss.
- Therapist normalized the variability and non-linear nature of the grieving process.
- Client explored changes in identity and functioning following the loss.
- Client identified grief-related triggers.
- Therapist supported client in processing unresolved emotions related to the loss.
- Client explored the impact of grief on interpersonal relationships.
- Therapist reinforced use of healthy coping strategies during periods of increased grief.
- Client demonstrated increased ability to tolerate and verbalize grief-related emotions.

CLINICAL WRITING

Here are some examples of clinical statements.

Anxiety

- Client identified situations associated with increased anxiety.
- Therapist provided psychoeducation regarding the physiological response to anxiety.
- Client identified anticipatory thoughts contributing to anxiety.
- Therapist assisted client in differentiating perceived threat from immediate danger.
- Client practiced grounding and relaxation techniques.
- Client explored avoidance behaviors that may reinforce anxiety.
- Therapist supported gradual development of tolerance for uncertainty.
- Client demonstrated increased awareness of anxiety-related thought patterns.

Anger / Emotional Dysregulation

- Client identified triggers contributing to increased anger.
- Therapist assisted client in identifying underlying emotions associated with anger.
- Client explored the relationship between perceived invalidation and emotional escalation.
- Therapist provided psychoeducation regarding the function of anger.
- Client identified physiological warning signs associated with escalating anger.
- Client practiced strategies for reducing emotional intensity.
- Client explored alternative behavioral responses to anger-provoking situations.
- Client demonstrated increased ability to pause before responding.

Interpersonal Relationships

- Client explored interpersonal patterns contributing to current distress.
- Therapist supported client in identifying communication patterns.
- Client explored boundaries within interpersonal relationships.
- Therapist provided psychoeducation regarding healthy boundaries.
- Client identified difficulty expressing needs directly.
- Client practiced assertive communication.
- Client explored the relationship between attachment experiences and current interpersonal expectations.
- Client identified patterns of people-pleasing and/or difficulty with saying no.
- Therapist supported client in identifying controllable versus uncontrollable aspects of interpersonal situations.

Skill Building

- Therapist modeled the identified skill and provided corrective feedback.
- Client practiced the skill during session.
- Client demonstrated understanding of the intervention.
- Client required minimal/moderate prompting to utilize the skill.
- Client required additional support to generalize the skill to real-world situations.
- Therapist reviewed barriers to implementation.
- Client identified situations in which the skill could be applied.
- Client developed a plan for practicing the skill between sessions.
- Client demonstrated increased confidence in utilizing the skill.
- Therapist reinforced previously established coping strategies.

CLINICAL WRITING

Here are some examples of clinical statements.

Psychoeducation

- Therapist provided psychoeducation regarding ____.
- Therapist reviewed the relationship between ____ and ____.
- Client demonstrated understanding of the information presented.
- Client asked appropriate questions and actively participated in discussion.
- Therapist provided education regarding common trauma responses.
- Therapist provided education regarding cognitive distortions.
- Therapist provided education regarding emotional regulation and distress tolerance.
- Therapist reviewed the rationale for utilizing ____.
- Client was able to identify how the information relates to their current experiences.

Risk/Safety Documentation

- Client denied current suicidal or homicidal ideation.
- Client denied current intent or plan to harm self or others.
- Client reported passive thoughts of death without current intent or plan.
- Risk factors and protective factors were assessed during session.
- Client identified future-oriented goals and protective factors.
- No acute safety concerns were identified during the session.

A few more helpful clinical phrases

- Client demonstrated increased psychological insight into the relationship between past experiences and current functioning.
- Client continues to exhibit maladaptive behavioral patterns that contribute to functional impairment.
- Client demonstrated increased capacity for self-reflection and emotional identification.
- Client's responses suggested limited awareness of the impact of current behaviors on interpersonal functioning.
- Client demonstrated emerging ability to recognize and challenge maladaptive cognitive patterns.
- Client appeared to experience heightened physiological and emotional arousal when discussing trauma-related material.
- Client demonstrated increased tolerance for discussing previously avoided material.
- Client continues to benefit from structured therapeutic support, psychoeducation, and skills development.