

The client sample was analyzed for plant-based cannabinoids by Liquid Chromatography (LC). The collected data was compared to data collected for certified reference standards at known concentrations.

63505-CN

Weight %	Concentration (mg/mL)			
0.12	1.13	•		
ND	ND			
3.51	32.58			
0.02	0.18			
0.03	0.28			
ND	ND			
3.69	34.21	0%	Cannabinoids (wt%)	3.5%
0.12	1.13			
3.51	32.58			
	0.12 ND 3.51 0.02 0.03 ND ND ND ND ND ND ND ND ND ND ND 3.69 0.12	0.12 1.13 ND ND 3.51 32.58 0.02 0.18 0.03 0.28 ND ND 3.69 34.21 0.12 1.13	0.12 1.13 ND ND 3.51 32.58 0.02 0.18 0.03 0.28 ND ND 1.13 0%	0.12 1.13 ND ND 3.51 32.58 0.02 0.18 0.03 0.28 ND ND 3.69 34.21 0% 0.12 1.13

Ratio of Total CBD to THC 28.7:1

Limit of Quantitation (LOQ) = 0.01 wt%

Max THC (and Max CBD) are calculated values for total cannabinoids after heating, assuming complete decarboxylation of the acid to the neutral form. It is calculated based on the weight loss of the acid group during decarboxylation: $Max THC = (0.877 \times THCA) + THC$. This calculation does not include other cannabinoid isomers (eg. D8-THC and exo-THC). ND = None detected above the limits of detection (LOD), which is half of LOQ.

END OF REPORT

420 Fortune Blvd • Milford, MA 01757 • 617-221-3356 www.ProVerdeLabs.com