

**SCIP SAMPLE QUESTIONS**

**SCIP SCREENING MODULE**

<b>SS3. Agoraphobia:</b>	<b>Kappa</b>
<p><i>Have you been afraid of being alone (at home or outside of the home), traveling in a car, train or plane, being in an open space (e.g. park) or being in a closed space (e.g. store), or being in crowds?</i></p> <p>0      Absent or non-significant 1      Symptom present</p>	<b>0.52</b>

<b>SS7. Witnessing or experiencing traumatic events:</b>	<b>Kappa</b>
<p><i>Have you ever witnessed or experienced a traumatic event that involved actual or threatened death or serious injury to you or someone else (e.g. physical or sexual abuse, terrorist attack, natural disaster, war...)? Did you feel intense fear and helplessness?</i></p> <p>0            No traumatic events 1            One traumatic event 2            More than one traumatic event</p>	<b>0.75</b>

<b>SS9. Depressed mood:</b>	<b>Kappa</b>
<p><i>Have you been feeling sad, depressed or in low spirits?</i></p> <p>0      Absent or non-significant 1      Symptom present</p>	<b>0.86</b>

<b>SS13. Irritable mood:</b>	<b>Kappa</b>
<p><i>Have you felt that you were easily irritated without reason? Have you found yourself so irritable that you shout at people or start arguments or actually become aggressive?</i></p> <p>0 Absent or non-significant 1 Symptom present</p>	<b>0.75</b>

<b>SS22. I would like to ask you about alcohol use over the <u>past year</u>:</b>	<b>Kappa</b>
<p><i>Did alcohol cause problems for you at work or school, problems with family or friends, legal problems, other problems such as getting in physical fights?</i></p> <p>0 Absent or non-significant 1 Symptom present</p>	<b>0.89</b>

<b>SS26. Worry about weight gain:</b>	<b>Kappa</b>
<p><i>Have you worried about gaining weight to the point that you did things such as self-induced vomiting, using diet pills, laxatives, or heavy exercise?</i></p> <p>0 Absent or non-significant 1 Symptom present</p>	<b>0.73</b>

**SCIP MODULES**

<b>MA1. PANIC ATTACKS WITHOUT PHOBIA:</b>	<b>kappa</b>
<p><i>Have you gotten suddenly anxious and frightened for a short time (up to 60 minutes)?</i></p> <p><i>During that time, did you feel that your heart was racing or pounding, or did you start shaking or sweating, or did you feel you were choking?</i></p> <p>0 Patient had no panic attacks.</p> <p>1 Patient had panic attacks.</p>	<b>0.92</b>

<b>MA16. GENERALIZED ANXIETY:</b>	<b>Kappa</b>
<p><i>Have you had excessive worry and anxiety for long periods (hours each day lasting several months), not just during panic attacks?</i></p> <p><i>Is it difficult to control the anxiety?</i></p> <p>0 Patient has no generalized anxiety.</p> <p>1 Patient has generalized anxiety.</p>	<b>0.84</b>

<b>Flashbacks:</b>	<b>Kappa</b>
<p><b>PTSD3.</b> <i>Did you have a sense or feeling that the event was happening again: the sense of reliving the event (flashbacks), auditory/visual hallucinations related to the event, or body/somatosensory experiences of the event?</i></p> <p>0 Patient had no significant symptom.</p> <p>1 Patient has a sense or feeling that the event is happening again, the sense of reliving the event (flashbacks).</p>	<b>0.87</b>

<b><i>Avoiding people, places, activities:</i></b>	<b>Kappa</b>
<b><i>PTSD7. Did you try to avoid things that reminded you of the event (such as certain people, certain places, or some activities)?</i></b>	<b>0.94</b>
<p>0 Patient had no significant symptom.</p> <p>1 Patient avoids things that are reminders of the event (such as certain people, certain places or some activities).</p>	
<b><i>MB3. Anhedonia:</i></b>	<b>Kappa</b>
<b><i>Have you been unable to experience pleasure and enjoy things that you used to enjoy like exercising, enjoying your hobbies, or socializing with friends?</i></b>	<b>0.87</b>
<b><i>Have you lost interest in your work or pleasure in your social activities?</i></b>	
<p>0 Patient has no anhedonia.</p> <p>1 Patient has anhedonia less than half the time.</p> <p>2 Patient has anhedonia more than half the time.</p>	
<b><i>MB9. Psychomotor slowing/ retardation:</i></b>	<b>Kappa</b>
<b><i>Have you felt as though you were talking or moving more slowly than normal for you when depressed?</i></b>	<b>0.72</b>
<p>0 Patient has normal energy and activity.</p> <p>1 Patient has psychomotor retardation less than half the time.</p> <p>2 Patient has psychomotor retardation more than half the time.</p>	
<b><i>MB26. Expansive (elated) mood:</i></b>	<b>Kappa</b>
<b><i>Have you sometimes felt very happy, elated, on top of the world without much reason?</i></b>	<b>0.75</b>
<p>0 Patient has no elated mood.</p> <p>1 Patient has elated mood less than half the time.</p> <p>2 Patient has elated mood more than half the time.</p>	

<b><i>MC1. Auditory hallucinations:</i></b>	<b>kappa</b>
<p><b><i>Do you hear noises (like music, whispering sounds) or voices talking to you when there is no one around?</i></b></p> <p><b><i>Are the voices like a real voice or just thoughts in your mind?</i></b></p> <p>0 Patient has no auditory hallucinations.  1 Patient has auditory hallucinations.  2 Patient has auditory hallucinations with command.</p>	<b>0.90</b>

<b><i>MC10. Delusions of spying:</i></b>	<b>Kappa</b>
<p><b><i>Do you think someone is spying on you or recording your conversations?</i></b></p> <p>0 Patient has no delusions of being spied upon.  1 Patient has delusions of being spied upon some of the time.  2 Patient has delusions of being spied upon most of the time.</p>	<b>0.86</b>

<b><i>MC19. Delusion of thought reading:</i></b>	<b>kappa</b>
<p><b><i>Do you think that you can read people's minds?</i></b>  <b><i>Do you think that other people can read your thoughts?</i></b></p> <p>0 Patient has no delusions of thought reading.  1 Patient has delusions of thought reading.</p>	<b>0.83</b>

<b>MC27_1</b>	<b><i>Derailment (looseness of association):</i></b>	<b>Kappa</b>
	0 Normal speech. 1 Patient has derailment (looseness of association): speech shifts to different topics, related or unrelated, but eventually comes back to the main topic. 2 Patient has severe derailment (looseness of association): speech shifts to different topics, mostly unrelated and never comes back to main topic.	<b>0.65</b>

<b>MD2. Withdrawal:</b>		<b>Kappa</b>
	<b><i>When you stopped or cut down on alcohol use, did you have withdrawal symptoms? (Interviewer gives examples of alcohol withdrawal symptoms)</i></b>  0 Patient had no withdrawal symptoms from alcohol. 1 Patient had withdrawal symptoms from alcohol.	<b>0.93</b>

<b>MD4. Unable to Control:</b>		<b>Kappa</b>
	<b><i>Did you use more of the drug than you planned or intended to use? Did you use the drug for a longer period of time than you had planned or intended?</i></b> 0 Patient had control over the drug use. 1 Patient used the drug more than what was planned or intended.	<b>0.97</b>

### **Most Recent SCIP References**

1. Aboraya A, Nasrallah HA, Elswick D, Elshazly A, Estephan N, Aboraya D, Berzingi Se, Chumber J, Berzingi Sa, Justice J, Zafar J, Dohar S. Measurement-based Care in Psychiatry: Past, Present and Future. *Innov Clin Neurosci*. 2018;15:25-38.
2. Aboraya A. The Reliability and Validity of the Core Schizophrenia Symptoms Scale of the Standard for Clinicians' Interview in Psychiatry *Schizophrenia Bulletin*. 2018;44:S365-S366.
3. Aboraya A. Instruction Manual for the Standard for Clinicians' Interview in Psychiatry (SCIP). *Innovations in CLINICAL NEUROSCIENCE*. 2016;13:41-77.
4. Aboraya A, Nasrallah H, Muvvala S, El-Missiry A, Mansour H, Hill C, Elswick D, Price EC. The Standard for Clinicians' Interview in Psychiatry (SCIP): A Clinician-administered Tool with Categorical, Dimensional, and Numeric Output-Conceptual Development, Design, and Description of the SCIP. *Innov Clin Neurosci*. 2016;13:31-77.
5. Aboraya A. The Validity Results of the Standard for Clinicians' Interview in Psychiatry (SCIP). *Schizophrenia Bulletin*. 2015;41:S103-S104.
6. Aboraya A, El-Missiry A, Barlowe J, John C, Ebrahimian A, Muvvala S, Brandish J, Mansour H, Zheng W, Chumber P, Berry J, Elswick D, Hill C, Swager L, Abo Elez W, Ashour H, Haikal A, Eissa A, Rabie M, El-Missiry M, El Sheikh M, Hassan D, Ragab S, Sabry M, Hendawy H, Abdel Rahman R, Radwan D, Sherif M, Abou El Asaad M, Khalil S, Hashim R, Border K, Menguito R, France C, Hu W, Shuttleworth O, Price E. The reliability of the standard for clinicians' interview in psychiatry (SCIP): a clinician-administered tool with categorical, dimensional and numeric output. *Schizophrenia research*. 2014;156:174-183.