

Glossary of Adult Psychopathology (GAP)
of the Standard for Clinicians' Interview in Psychiatry (SCIP)

SCREENING ITEMS:

Glossary Code	Item Title/Subtitle	Item Definitions	Item Questions	# of cases	K	SE
S1_GEN_ANX	Generalized anxiety	Patient feels very anxious and afraid out of proportion to the situation (with or without physical symptoms) for a prolonged period almost every day.	<i>Have you felt very anxious and afraid out of proportion to the situation (with or without physical symptoms) for a prolonged period almost every day?</i> 0 Absent or non-significant. 1 Symptom present. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	61	0.76	0.05
S2_PAN	Panic attacks	Panic attack: Patient suddenly feels anxious and frightened for a short time (up to 60 minutes) and develops physical symptoms such as fast heart beats, shaking, sweating, difficulty breathing.	<i>Did you have a panic attack, during which you suddenly felt anxious and frightened for a short time (up to 60 minutes) and developed physical symptoms (e.g., fast heart beats, shaking, sweating)?</i> 0 Absent or non-significant. 1 Symptom present. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	54	0.81	0.05
S3_SOC_PHOB	Social Phobia	Patient is afraid and anxious when he/she does things in front of people, such as eating or speaking in public.	<i>Have you been afraid and anxious when you do things in front of people, such as eating or speaking in public?</i> 0 Absent or non-significant. 1 Symptom present. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	22	0.51	0.05

S4_AGORA_PHOB	Agoraphobia	Patient is afraid of being alone (at home or outside of home), traveling in a car, train or plane, being in an open space (e.g., park) or being in a closed space (e.g., store), or being in crowds.	<i>Have you been afraid of being alone (at home or outside of home), traveling in a car, train or plane, being in an open space (e.g., park) or being in a closed space (e.g., store), or being in crowds?</i> 0 Absent or non-significant. 1 Symptom present. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	26	0.52	0.05
S5_OBS	Screening for obsessions	The patient has an intrusive thought or image that does not make sense and keeps coming back to his/her mind even when the patient tries not to have the thought or the image.	<i>Do you ever have an intrusive thought or image that does not make sense and keeps coming back to your mind even when you try not to have the thought or the image?</i> 0 Absent or non-significant. 1 Symptom present. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	38	0.70	0.04
S6_COMP	Screening for compulsions	The patient has to do things over and over, that is, checking things he/she has done (such as washing hands even if they are clean, checking doors or repeating mental acts such as counting or praying).	<i>Do you find that you have to do things over and over, that is, checking things you have done (such as washing your hands even if they are clean, checking doors or repeating mental acts such as counting or praying)?</i> 0 Absent or non-significant. 1 Symptom present. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	31	0.58	0.05
S7_TRAUMA	Witness or experience traumatic events	The patient witnessed or experienced a traumatic event that involved actual or threatened death or serious injury to the patient or someone else (e.g., physical or sexual abuse, terrorist attack, natural disaster, war...). The patient felt intense fear and helplessness.	<i>Have you ever witnessed or experienced a traumatic event that involved actual or threatened death or serious injury to you or someone else (e.g., physical or sexual abuse, terrorist attack, natural disaster, war...)? Did you feel intense fear and helplessness?</i> 0 No traumatic events. 1 One traumatic event. 2 More than one traumatic event. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	69	0.75	0.05

S8_RE_EXP	Re-experience traumatic events	The patient has re-experienced the event in a distressing way (e.g., flashbacks, nightmares, bad dreams).	<i>Over the past month, have you re-experienced the event in a distressing way (e.g., flashbacks, nightmares, bad dreams)?</i> 0 Absent or non-significant. 1 Symptom present. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	34	0.89	0.05
S9_DEP	Depressed mood	The patient has been feeling sad, depressed or in low spirits.	<i>Have you been feeling sad, depressed or in low spirits?</i> 0 Absent or non-significant. 1 Symptom present. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	158	0.86	0.04
S10_ANHED	Anhedonia	The patient has been unable to experience pleasure and enjoy things that he/she used to enjoy like exercising, enjoying hobbies or socializing with friends.	<i>Have you been unable to experience pleasure and enjoy things that you used to enjoy like exercising, enjoying your hobbies or socializing with friends?</i> 0 Absent or non-significant. 1 Symptom present. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	125	0.87	0.04
S11_SUICIDAL	Suicidal ideation, intention and plan	The patient had thoughts about harming self. The patient had the intention to carry out the suicidal thoughts. The patient had specific suicidal plans. The patient had one or more suicide attempts during the past month.	<i>Have you had thoughts about harming yourself?</i> <i>Have you had the intention to carry out the suicidal thoughts?</i> <i>Have you had suicidal plans?</i> <i>Have you attempted to harm yourself during the past month?</i> <i>(multiple choices allowed).</i> 0 Absent. 1 Current suicidal ideations. 2 Current suicidal intention. 3 Current suicidal plans. 4 Recent suicide attempts (past month). . Not sure <i>or</i> Not applicable <i>or</i> Missing.	79	0.61	0.04

S12_SELF_MUTIL	Self-mutilation behaviors	The patient engages in self-mutilation behaviors without intending to die (e.g., burning, cutting, scratching).	<i>Have you attempted self-mutilation behaviors without intent to die (e.g., burning, cutting, scratching)?</i> 0 Absent. 1 Current self-mutilation (past month). . Not sure <i>or</i> Not applicable <i>or</i> Missing.	79	0.61	0.04
S13_ELATED	Elated mood	The patient feels very happy, elated, on top of the world for no apparent reason.	<i>Have you felt very happy, elated, on top of the world for no apparent reason?</i> 0 Absent or non-significant. 1 Symptom present. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	76	0.72	0.05
S14_IRRITAB	Irritable mood	The patient feels easily irritated without reason. The patient feels so irritable that he/she shouts at people or starts arguments or actually becomes aggressive.	<i>Have you felt easily irritated without reason? Have you found yourself so irritable that you shout at people or start arguments or actually become aggressive?</i> 0 Absent or non-significant. 1 Symptom present. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	65	0.75	0.05
S15_MIXED	Mixed mood (same day mood changes)	The patient has mixed mood swings: periods of <u>depression and elation or irritability on the same day.</u>	<i>Have you had mixed mood swings: periods of <u>depression and elation or irritability on the same day?</u></i> 0 Absent or non-significant. 1 Symptom present. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	44	0.50	0.05
S16_PARANOID	Paranoid delusions	The patient feels that someone is spying on him/her or trying to harm the patient or has a plot or conspiracy against the patient.	<i>Have you felt that someone is spying on you or trying to harm you or has a plot or conspiracy against you?</i> 0 Absent or non-significant. 1 Symptom present. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	97	0.83	0.04

S17_DEL	Other delusions	The patient has strange thoughts such as that he/she is a prominent person in society, or he/she is divine or God, or he/she receives special messages from TV or newspapers.	<i>Do you have strange thoughts such as that you are a prominent person in society, or you yourself are divine or you are God, or you receive special messages from TV or newspapers?</i> 0 Absent or non-significant. 1 Symptom present. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	39	0.77	0.04
S18_AUDH	Auditory hallucinations	The patient has experiences of hearing voices or noises that other people cannot hear.	<i>Have you had experiences of hearing voices or noises that other people cannot hear?</i> 0 Absent or non-significant. 1 Symptom present. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	92	0.76	0.04
S19_VISH	Visual hallucinations	The patient has experiences of seeing things (shadows, objects, people) that other people cannot see.	<i>Have you had experiences of seeing things (shadows, objects, people) that other people cannot see?</i> 0 Absent or non-significant. 1 Symptom present. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	51	0.68	0.05
S20_VIOL	Violence	The patient has been violent (destructive to objects or violent towards people) in the past (with or without the influence of alcohol or drugs).	<i>Have you been violent (destructive to objects or violent towards people) in the past (with or without the influence of alcohol or drugs)?</i> 0 No history of violence. 1 Yes, due to alcohol and/or drugs. 2 Yes, without use of alcohol or drugs. 3 Both, with and without use of alcohol or drugs. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	74	0.64	0.04

S21_DIS_BEH	Disorganized behavior	There is evidence of disorganized behavior by observation during the interview (agitation, odd appearance, inappropriate social behavior, inappropriate affect).	<i>There is evidence of disorganized behavior by observation during the interview (agitation, odd appearance, inappropriate social behavior, inappropriate affect).</i> 0 Absent or non-significant. 1 Symptom present. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	32	0.54	0.04
S22_DIS_TH	Disorganized thoughts	Evidence of disorganized thoughts by observation during the interview (loose associations, tangentiality, incoherent speech).	<i>Evidence of disorganized thoughts by observation during the interview (loose associations, tangentiality, incoherent speech).</i> 0 Absent or non-significant. 1 Symptom present. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	39	0.65	0.04
S23_ALC_PROB	Alcohol problems	Alcohol causes problems for the patient at work or school, problems with family or friends, legal problems, or other problems such as physical fights.	<i>During the past year, did alcohol cause problems for you at work or school, problems with family or friends, legal problems, or other problems such as getting in physical fights?</i> 0 Absent or non-significant. 1 Symptom present. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	53	0.89	0.06
S24_DRG_PROB	Drug problems	Drug use () causes problems for the patient at work or school, problems with family or friends, legal problems, or other problems such as getting in physical fights.	<i>During the past year, did drug use () cause problems for you at work or school, problems with family or friends, legal problems, or other problems such as getting in physical fights?</i> 0 Absent or non-significant. 1 Symptom present. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	17	0.78	0.06

S25_SOM_SX	Somatic symptoms	The patient had visited doctors due to physical illness and the doctors did the necessary work up and could not find a medical explanation for the symptoms.	<i>Have you visited doctors due to physical illness and the doctors did the necessary work up and could not find a medical explanation (have patient give examples)?</i> 0 Absent or non-significant. 1 Symptom present. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	33	0.81	0.05
S26_PAIN	Pain symptoms	The patient had pain and the doctor did the necessary work up and could not really explain why.	<i>Have you had pain and your doctor did the necessary work up and could not really explain why?</i> 0 Absent or non-significant. 1 Symptom present. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	24	0.93	0.05
S27_ANOREXIA	Worry about weight gain	The patient worried about gaining weight to the point that he/she engaged in self-induced vomiting, using diet pills, laxatives, or heavy exercise.	<i>Have you worried about gaining weight to the point that you did things such as self-induced vomiting, using diet pills, laxatives, or heavy exercise?</i> 0 Absent or non-significant. 1 Symptom present. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	12	0.73	0.05
S28_BINGE	Binge eating	The patient had episodes of binge eating (eating within one or two-hour period what most people would consider an unusually large amount of food).	<i>Do you have episodes of binge eating (eating within one or two-hour period what most people would consider an unusually large amount of food)?</i> 0 Absent or non-significant. 1 Symptom present. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	27	0.97	0.12

S29_SUS_ATTN	Sustained attention impairment	The patient has difficulty concentrating on one thing for a long time (e.g., reading a book, writing a letter).	<i>Do you have difficulty concentrating on one thing for a long time (e.g., reading a book, writing a letter)?</i> 0 Absent or non-significant. 1 Patient has difficulty concentrating for a long period of time. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	39	0.95	0.12
S30_FIDGETY	Fidgety	The patient has difficulty remaining seated (fidgets with hands and feet, squirms or wiggles in seat) when he/she is expected to remain seated (e.g., in a meeting or a church service).	<i>Do you have difficulty remaining seated (fidget with hands and feet, squirm or wiggle in seat) when expected to remain seated (e.g., in a meeting or a church service)?</i> 0 = Absent or non-significant. 1 = Patient fidgets with hands and feet, or wiggles in seat. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	41	0.81	0.12

PSYCHOPATHOLOGY ITEMS:

Glossary Code	Item Titles/subtitles	Item Definitions	Item Questions	# of Cases	K	SE
G1_POOR_CONC_1 (*)	Attention impairment (poor concentration)	The patient has difficulty paying attention and concentrating when reading an article, watching a TV show or a movie, or doing work or school assignments.	<i>Do you have difficulty paying attention and concentrating when reading an article, watching a TV show or a movie, or doing your work or school assignments?</i> 0 = Absent or non-significant. 1 = Patient has poor attention and concentration. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	41	0.92	0.12
G1_SUS_ATTN_IMPAIR_2	Sustained attention impairment	The patient has difficulty concentrating on one thing for a long time (e.g., reading a book, writing a letter).	<i>Do you have difficulty concentrating on one thing for a long time (e.g., reading a book, writing a letter)?</i> 0 = Absent or non-significant. 1 = Patient has difficulty concentrating for a long period of time. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	39	0.95	0.12
G1_AVOID_TASKS_3	Avoiding sustained attention tasks	The patient avoids tasks that require a lot of concentration at work, school, or home (e.g., reading a book, writing a letter).	<i>Do you avoid tasks that require a lot of concentration at work, school, or home (e.g., reading a book, writing a letter)?</i> 0 = Absent or non-significant. 1 = Patient avoids tasks that require sustained mental effort. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	34	0.97	0.12

An asterisk (*) below the glossary code indicates that the item is repeated elsewhere in the GAP.

G1_ATTN_SPOKEN_4	Attention when spoken to	Patient has difficulty concentrating on what people say to him/her, even when they are speaking to the patient directly.	<i>Do you have difficulty concentrating on what people say to you, even when they are speaking to you directly?</i> 0 = Absent or non-significant. 1 = Patient has difficulty concentrating on what people say. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	32	0.97	0.12
G1_MANAGE_TIME_5	Organization and meeting deadlines	The patient has difficulty with tasks that require organization and keeping track of many things all at once (e.g., planning and organizing work duties or household chores). The patient has difficulty managing his/her time (e.g., usually fails to meet deadlines).	<i>Do you have difficulty with tasks that require organization and keeping track of many things all at once (e.g., planning and organizing your work or household chores)?</i> <i>Do you have difficulty managing your time (e.g., usually fail to meet deadlines)?</i> 0 = Absent or non-significant. 1 = Patient has difficulty with tasks that require organization or meeting deadlines. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	30	0.82	0.12
G1_CHANGE_ACTIVITY_6	Changing activities	The patient changes from one activity to another without finishing the first.	<i>Do you change from one activity to another without finishing the first?</i> 0 = Absent or non-significant. 1 = Patient changes from one activity to another without finishing the first. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	40	0.92	0.12

G1_DISTRACTION_7 (*)	Distraction (Attention is distracted by environmental noises)	The patient is easily distracted from tasks by activity or noise in the surroundings.	<i>Are you easily distracted from tasks by</i> <i>activity or noise around you?</i> 0 = Absent or non-significant. 1 = Patient is easily distracted from tasks by activity or noise around. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	43	0.97	0.12
G1_MISPLACE_THINGS_8	Misplacing things	The patient loses or misplaces things more often than others (e.g., wallets, keys, cell phones).	<i>Do you lose or misplace things more often</i> <i>than others (e.g., wallets, keys, cell phones)?</i> 0 = Absent or non-significant. 1 = Patient loses or misplaces things more often than others. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	43	0.94	0.12
G1_FORGET_ACTIVITY_9	Forgetting daily activities	The patient forgets daily activities more often than others (e.g., appointments, paying bills, returning phone calls).	<i>Do you forget daily activities more often</i> <i>than others (e.g., appointments, paying bills,</i> <i>returning phone calls)?</i> 0 = Absent or non-significant. 1 = Patient forgets daily activities more often than others. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	24	0.94	0.12
G1_LOSE_TRACK_10	Losing track	The patient loses track of what he/she is doing (e.g., forgets why he/she went to get something).	<i>Do you lose track of what you are doing</i> <i>(e.g., forget why you went to get</i> <i>something)?</i> 0 = Absent or non-significant. 1 = Patient loses track of what he/she is doing. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	40	0.92	0.12

G1_FIDGETY_11	Fidgety	The patient has difficulty remaining seated (fidgets with hands and feet, squirms or wiggles in seat) when he/she is expected to remain seated (e.g., in a meeting or a church service).	<i>Do you have difficulty remaining seated (fidget with hands and feet, squirm or wiggle in seat) when expected to remain seated (e.g., in a meeting or a church service)?</i> 0 = Absent or non-significant. 1 = Patient fidgets with hands and feet, or wiggles in seat. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	41	0.81	0.12
G1_LEAVE_SEAT_12	Leaving seats	The patient leaves his/her seat in meetings or other situations (e.g., during an appointment or a church service) where the expectation is to remain seated.	<i>Do you leave your seat in meetings or other situations (e.g., during an appointment or a church service) where you are expected to remain seated?</i> 0 = Absent or non-significant. 1 = Patient leaves seat in meetings or other situations where expected to remain seated. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	30	0.88	0.12
G1_GETUP_MOVE_13	Restlessness/moving	The patient feels restless, fidgety and must get up and move around.	<i>Do you feel restless, fidgety and you must get up and move around?</i> 0 = Absent or non-significant. 1 = Patient feels restless, fidgety and must move around. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	49	0.61	0.12

G1_HYPERACTIVE_14	Hyperactivity	The patient feels overly active and compelled to do things, like he/she is driven by a motor.	<i>Do you feel overly active and compelled to do things, like you are driven by a motor?</i> 0 = Absent or non-significant. 1 = Patient feels overly active and compelled to do things. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	22	0.97	0.12
G1_WAIT_LINE_15	Waiting in line	The patient has difficulty waiting in line for his/her turn when the situation calls for it.	<i>Is it difficult for you to wait in line for your turn when the situation calls for it?</i> 0 = Absent or non-significant. 1 = Patient has difficulty waiting in line. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	23	1.00	0.12
G1_TALK_ALOT_16	Talking too much	The patient thinks he/she talks too much. Others say that the patient talks too much.	<i>Do you think you talk too much?</i> <i>Do others say that you talk too much?</i> 0 = Absent or non-significant. 1 = Patient talks too much. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	12	1.00	0.12
G1_LOUD_NOISY_17	Loud and noisy	The patient thinks that he/she is a loud and noisy person. Other people sometimes ask the patient to quiet down or lower his/her voice.	<i>Do you think that you are a loud and noisy person?</i> <i>Do other people sometimes ask you to quiet down or lower your voice?</i> 0 = Absent or non-significant. 1 = Patient or others feel the patient is loud and noisy. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	22	0.58	0.11

G1_IMPULSIVE_18	Impulsivity	The patient is impulsive (e.g., acts before he/she thinks adequately about consequences of actions).	<i>Are you impulsive (e.g., act before you think adequately about consequences of actions)?</i> 0 = Absent or non-significant. 1 = Patient is impulsive. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	41	0.92	0.12
G1_DISTURB_OTHERS_19	Disturbing others	The patient disturbs others or intrudes on others (e.g., when people are talking or when people are involved in activities).	<i>Do you disturb others or intrude on others (e.g., when people are talking or when people are involved in activities)?</i> 0 = Absent or non-significant. 1 = Patient disturbs others or intrudes on others. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	23	0.97	0.12
G1_BLURT_OUT_20	Blurt out answers	The patient has tendency to blurt out an answer before another person has finished asking the question.	<i>Do you have tendency to blurt out an answer before another person has finished asking the question?</i> 0 = Absent or non-significant. 1 = Patient blurts out the answers. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	32	0.89	0.12

G2_GEN_ANX_21	Generalized anxiety	The patient has excessive worry and anxiety for long periods (hours each day lasting several months), not just during panic attacks. The patient has difficulty controlling the anxiety.	<i>Have you had excessive worry and anxiety for long periods (hours each day lasting several months), not just during panic attacks?</i> <i>Is it difficult to control the anxiety?</i> 0 Patient has no generalized anxiety. 1 Patient has generalized anxiety. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	25	0.84	0.04
G2_RESTLESS_22	Restlessness with anxiety	The patient feels restless, keyed up or on edge.	<i>Did you feel restless, keyed up or on edge?</i> 0 No. 1 Yes . Not sure <i>or</i> Not applicable <i>or</i> Missing.	26	0.74	0.04
G2_TENSION_23	Tension with anxiety	The patient feels tense in muscles.	<i>Did you feel tense in your muscles?</i> 0 No. 1 Yes. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	22	0.77	0.04
G2_EXHAUSION_24	Exhaustion with anxiety	The patient feels tired, or easily exhausted even without work.	<i>Did you feel tired, or easily exhausted even without work?</i> 0 No. 1 Yes. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	22	0.79	0.05

G2_POOR_CONC_25 (*)	Poor concentration with anxiety	The patient has difficulty paying attention and concentrating when reading an article, watching a TV show or a movie, or doing work or school assignments due to anxiety.	<i>Did you have difficulty concentrating when anxious?</i> 0 No. 1 Yes. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	27	0.76	0.05
G2_IRRITAB_26	Irritability with anxiety	The patient feels irritable when anxious.	<i>Did you feel irritable when anxious?</i> 0 No. 1 Yes. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	28	0.83	0.04
G2_INSOMNIA_27	Insomnia with anxiety	The patient has difficulty falling asleep or staying asleep when anxious.	<i>Did you have difficulty falling asleep or staying asleep when anxious?</i> 0 No. 1 Yes. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	25	0.82	0.05

G3_PAN_28	Panic attacks	The patient suddenly becomes anxious and frightened for a short period of time (up to 60 minutes). During that time, the patient feels that the heart is racing or pounding, or he/she is shaking or sweating, or choking.	<i>Did you have a panic attack, when you suddenly become anxious and frightened for a short period of time (up to 60 minutes)? During that time, did you feel that your heart was racing or pounding, or did you start shaking or sweating, or did you feel you were choking?</i> 0 Patient had no panic attacks. 1 Patient had panic attacks. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	30	0.92	0.06
G3_PAN_WORRY_29	Worry about having another panic attack	After a panic attack, the patient worries about having another attack. The patient worries about its effects (e.g., losing control, having a heart attack or going crazy).	<i>After a panic attack, did you worry about having another attack? Did you worry about its effects (e.g., losing control, having a heart attack or going crazy)?</i> 0 After a panic attack, patient did not worry about having another one. 1 After a panic attack, patient worried about having another one or its effects. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	25	0.81	0.04
G3_PAN_END_30	Action to end or prevent panic attacks	The patient does something to end the attack, like leaving a store, calling someone, taking deep breaths. The patient does something to prevent attacks (like avoiding places that trigger the panic attacks).	<i>Did you have to do something to end the attack, like leaving a store, calling someone, taking deep breaths? Do you do anything to prevent attacks (like avoiding places that trigger the panic attacks)?</i> 0 After a panic attack, patient does nothing to end or prevent another panic attack. 1 After a panic attack, patient does something to end or prevent another panic attack. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	26	0.87	0.04

G4_SOC_PHOB_31	Social phobia	The patient becomes afraid and anxious when he/she does things in front of people, such as eating or speaking in public. The patient avoids social situations or endures them with intense fear.	<i>Have you been afraid and anxious when you do things in front of people, such as eating or speaking in public?</i> <i>Do you avoid social situations or endure them with intense fear?</i> 0 Patient had no social phobia. 1 Patient had social phobia. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	22	0.51	0.05
G4_AGORA_PHOB_32	Agoraphobia	The patient is afraid of being alone (at home or outside of home), traveling in a car, train or plane, being in an open space (e.g., park) or being in a closed space (e.g., store), or being in crowds. The patient avoids these situations, or requires a companion, or endures with intense fear.	<i>Have you been afraid of being alone (at home or outside of home), traveling in a car, train or plane, being in an open space (e.g., park) or being in a closed space (e.g., store), or being in crowds?</i> <i>Do you avoid these situations, or require a companion, or endure with intense fear?</i> 0 Patient had no agoraphobia. 1 Patient had agoraphobia. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	26	0.52	0.05

G4_SP_PHOB_32A	Specific phobia	The patient has strong fears of certain objects or situations (e.g., heights, animals, spiders, snakes, seeing blood, receiving injections). The patient avoids objects, situations, or endures with intense fear.	<i>Have you had strong fears of certain objects or situations (e.g., heights, animals, spiders, snakes, seeing blood, receiving injections)?</i> <i>Do you avoid objects or situations, or endure with intense fear?</i> 0 Patient had no specific phobias. 1 Patient had a specific phobia. . Not sure or Not applicable or Missing.			
G5_OBS_33	Obsessions	The patient has an intrusive thought or image that does not make sense and keeps coming back to the patient's mind even when he/she tries to avoid the thought or the image.	<i>Do you ever have an intrusive thought or image that does not make sense and keeps coming back to your mind even when you try to avoid the thought or the image?</i> 0 Patient has no obsessions. 1 Patient has obsessions less than 1 hour/day. 2 Patient has obsessions 1-4 hours/day. 3 Patient has obsessions more than 4 hours/day. . Not sure or Not applicable or Missing.	26	0.85	0.04
G5_OBS_AGGRESS_34	Aggressive Obsessions	<u>Potential examples:</u> fears of harming oneself, fears of unintentionally hurting someone, urges to stab someone, thoughts of “losing control” and hurting a partner or significant other.	<i>Do you have obsessive thoughts related to aggression?</i> 0 No. 1 Yes. . Not sure or Not applicable or Missing.			

G5_OBS_CONTAM_35	Contamination Obsessions	Potential examples: fear of contamination in public bathrooms, outdoors, or public places, fear of what might happen after touching one's own bodily secretions, fear of getting germs or viruses from others.	<i>Do you have obsessive thoughts related to contamination?</i> 0 No. 1 Yes. . Not sure <i>or</i> Not applicable <i>or</i> Missing.			
G5_OBS_SEX_36	Sexual Obsessions	Potential examples: disturbing thoughts about sex, unwanted images of sexual acts toward strangers, family members or children, fear of being found with pornography.	<i>Do you have obsessive thoughts related to sex/sexuality?</i> 0 No. 1 Yes. . Not sure <i>or</i> Not applicable <i>or</i> Missing.			
G5_OBS_RELIG_37	Religious Obsessions	Potential examples: fear of having blasphemous or "sinful" thoughts, concerns for engaging in sinful or forbidden behavior, feeling the need to complete a certain number of prayers or "good deeds".	<i>Do you have obsessive thoughts related to religion?</i> 0 No. 1 Yes. . Not sure <i>or</i> Not applicable <i>or</i> Missing.			
G5_OBS_SOMAT_38	Somatic Obsessions	Potential examples: worries about having an undiagnosed illness like cancer or heart disease, fear of contracting a deadly disease, images of your own death.	<i>Do you have obsessive thoughts related to your health?</i> 0 No. 1 Yes. . Not sure <i>or</i> Not applicable <i>or</i> Missing.			

G5_OBS_APPEAR_39	Appearance Obsessions	Potential examples: worries about the size of certain body parts (ears, nose, mouth), fears of certain body parts being disgusting to others.	<i>Do you have obsessive thoughts about your physical appearance?</i> 0 No. 1 Yes. . Not sure <i>or</i> Not applicable <i>or</i> Missing.			
G5_COMP_40	Compulsions	The patient has to do things over and over, that is, checking things he/she has done (such as washing hands even if they are clean, checking doors or repeating mental acts such as counting or praying). The patient gets very anxious or tense if he/she does not repeat the act over and over.	<i>Do you find that you have to do things over and over, that is, checking things you have done (such as washing your hands even if they are clean, checking doors or repeating mental acts such as counting or praying)?</i> <i>Do you get very anxious or tense if you do not repeat the act over and over?</i> 0 Patient has no compulsions. 1 Patient has compulsions less than 1 hour/day. 2 Patient has compulsions 1-4 hours/day. 3 Patient has compulsions more than 4 hours/day. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	18	0.77	0.04
G5_COMP_CHECK_41	Checking Compulsions	Potential examples: checking door locks, switches, or appliances many times before leaving the house, checking one's appearance for an excessive amount of time before leaving the house.	<i>Do you feel the need to check things over and over again?</i> 0 No. 1 Yes. . Not sure <i>or</i> Not applicable <i>or</i> Missing.			

G5_COMP_CLEAN_42	Cleaning/Washing Compulsions	<u>Potential examples:</u> excessively changing your clothes, excessive handwashing, tooth brushing, showering, scrubbing surfaces.	<i>Do you feel the need to clean/wash things over and over again?</i> 0 No. 1 Yes. . Not sure <i>or</i> Not applicable <i>or</i> Missing.			
G5_COMP_REPEAT_43	Repeating Compulsions	<u>Potential examples:</u> re-reading text out of concerns that one “missed something,” re-writing or re-tracing words, re-entering buildings or living spaces, re-tying one’s shoes.	<i>Do you feel the need to repeat things many times?</i> 0 No. 1 Yes. . Not sure <i>or</i> Not applicable <i>or</i> Missing.			
G5_COMP_ORDER_44	Ordering/Arranging Compulsions	<u>Potential examples:</u> re-arranging one’s pantry or refrigerator, spending excessive amounts of time organizing one’s desk or workspace.	<i>Do you feel the need to arrange and rearrange things, order and reorder items over and over again?</i> 0 No. 1 Yes. . Not sure <i>or</i> Not applicable <i>or</i> Missing.			
G5_COMP_HOARD_45	Hoarding/Collecting	<u>Potential examples:</u> picking up, collecting or buying useless things, owning an excessive amount of items and being unable to donate or get rid of them.	<i>Do you feel the need to collect and/or hoard things?</i> 0 No. 1 Yes. . Not sure <i>or</i> Not applicable <i>or</i> Missing.			

G5_COMP_MENTAL_46	Mental Compulsions	<u>Potential examples:</u> repetitively counting certain numbers or avoiding certain “bad” numbers, repeating certain words, counting senseless things (e.g., ceiling tiles), reciting prayers or statements.	<i>Do you engage in any repetitive mental behaviors?</i> 0 No. 1 Yes. . Not sure <i>or</i> Not applicable <i>or</i> Missing.			
G6_DEP_47	Depressed mood	The patient feels sad, depressed or in low spirits.	<i>Have you been feeling sad, depressed or in low spirits?</i> 0 Patient has no depressed mood. 1 Patient has depressed mood less than half the time. 2 Patient has depressed mood more Than half the time. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	128	0.91	0.04
G6_ANHEDONIA_48 (*)	Anhedonia (Loss of pleasure and interest)	The patient has been unable to experience pleasure and enjoy things that he/she used to enjoy like exercising, enjoying hobbies, or socializing with friends.	<i>Have you been unable to experience pleasure and enjoy things that you used to enjoy like exercising, enjoying your hobbies, or socializing with friends?</i> 0 Patient has no anhedonia. 1 Patient has anhedonia less than half the time. 2 Patient has anhedonia more than half the time. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	121	0.87	0.04

G6_HOPELESSNESS_49	Hopelessness	The patient feels hopeless about his/her future.	<i>Have you felt hopeless about your future?</i> 0 Patient is not hopeless. 1 Patient feels hopeless less than half the time. 2 Patient feels hopeless more than half the time. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	11	0.82	0.04
G6_POOR_CONC_50 (*)	Attention impairment (poor concentration)	The patient has decreased concentration and he/she is unable to complete a task (e.g., at work, reading an article, reading a book, or watching a movie), even though the patient was able to do that before.	<i>Have you found that your concentration has decreased and you are unable to complete a task (e.g., at work, reading an article, reading a book, or watching a movie), even though you were able to do that before?</i> 0 Patient has no concentration problems. 1 Patient has difficulty concentrating less than half the time. 2 Patient has difficulty concentrating more than half the time. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	116	0.80	0.04
G6_PSYCHOM_SLOW_51 (*)	Psychomotor retardation (slowing)	The patient is talking or moving more slowly than normal when depressed.	<i>Have you felt as though you were talking or moving more slowly than normal for you when depressed?</i> 0 Patient has normal activity. 1 Patient has psychomotor retardation less than half the time. 2 Patient has psychomotor retardation more than half the time. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	97	0.72	0.04

G6_WORTHLESS_52	Worthlessness (low self-esteem)	The patient has low self-esteem and feels that he/she is a worthless person in the society or a failure.	<i>Have you felt that you are a worthless person in the society or a failure?</i> 0 Patient has no feeling of worthlessness. 1 Patient feels worthless less than half the time. 2 Patient feels worthless more than half the time. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	97	0.78	0.04
G6_GUILT_53	Guilt	The patient feels guilty or ashamed of himself/herself for something the patient has done or thought.	<i>Have you felt guilty or ashamed of yourself for something you have done or thought?</i> 0 Patient has no feeling of guilt. 1 Patient feels guilty less than half the time. 2 Patient feels guilty more than half the time. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	86	0.80	0.04
G6_SUICDALITY_54	Suicidality	The patient had thoughts, intentions or plans to harm or kill himself/herself during the past month. The patient may have made one or more suicide attempt during the past month.	<i>Have you had thoughts, intentions or plans to harm or kill yourself during the past month?</i> <i>Have you made a suicide attempt during the past month?</i> 0 Patient had no suicidal thoughts or made a suicide attempt during the past month. 1 Patient had thoughts, intentions or plans to harm self. 2 Patient made one recent suicide attempt during the past month. 3 Patient made two or more recent suicide attempts during the past month. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	68	0.64	0.04

G6_CRY_DEP_55	Crying when depressed	Patient has crying spells when depressed.	<i>Have you cried when depressed?</i> 0 Patient has no crying spells. 1 Patient has crying spells due to sadness less than half the time. 2 Patient has crying spells due to sadness more than half the time. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	11	0.76	0.04
G6_FATIGUE_56	Fatigue and loss of energy	Patient feels tired and exhausted during the day, even when he/she sleeps well and does not work very hard. Even though patient may feel tired and exhausted, he/she may have no decrease in the three main goal-directed activities in life: school, work, or social activities with other people (item # 130).	<i>Have you felt tired and exhausted during the day, even when you slept well and did not work very hard?</i> 0 Patient has no fatigue or loss of energy. 1 Patient feels tired and exhausted less than half the time. 2 Patient feels tired and exhausted more than half the time. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	97	0.72	0.04
G6_APPT_LOSS_57	Loss of appetite when depressed	Patient has marked loss of appetite when depressed.	<i>Have you lost your appetite when depressed?</i> 0 Patient had no loss of appetite. 1 Patient had marked loss of appetite for 2 weeks or less. 2 Patient had marked loss of appetite for more than 2 weeks. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	93	0.79	0.04
G6_APPT_INCREASE_58	Increased appetite when depressed	Patient has marked increase of appetite when depressed.	<i>Has your appetite increased when depressed?</i> 0 Patient had no increase of appetite. 1 Patient had marked increase of appetite for 2 weeks or less. 2 Patient had marked increase of appetite for more than 2 weeks. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	93	0.79	0.04

G6_WT_LOSS_59	Weight loss	Patient had marked loss of weight when depressed.	<i>Did you lose weight when depressed?</i> 0 Patient had no weight loss or minimal weight loss. 1 Patient lost more than 5% of body weight in a month. 2 Patient lost more than 15% of body weight in a year. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	62	0.71	0.04
G6_WT_GAIN_60	Weight gain	Patient had marked weight gain when depressed.	<i>Did you gain weight when depressed?</i> 0 Patient had no weight gain or minimal weight gain. 1 Patient gained more than 5% of body weight in a month. 2 Patient gained more than 15% of body weight in a year. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	15	0.76	0.05
G6_INIT_INSOMNIA_61	Initial insomnia	Patient has difficulty falling asleep (one hour or more) more than half the time when depressed.	<i>Have you had difficulty falling asleep when depressed?</i> 0 Patient has no sleeping problems. 1 Patient has difficulty falling asleep (one hour or more) more than half the time when depressed. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	103	0.79	0.04
G6_MID_INSOMNIA_62	Middle insomnia	Patient has difficulty staying asleep (awakens and stays awake one hour or more) more than half the time when depressed.	<i>Have you had difficulty staying asleep when depressed?</i> 0 Patient has no sleeping problems. 1 Patient has difficulty staying asleep (awakens and stays awake one hour or more) more than half the time when depressed. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	79	0.65	0.04

G6_LATE_INSOMNIA_63	Late insomnia	Patient has early awakenings (at least 2 hours earlier than intended) more than half the time when depressed.	<i>Have you been waking much earlier than your usual?</i> 0 Patient has no sleeping problems. 1 Patient has early waking (at least 2 hours) more than half the time when depressed. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	46	0.62	0.04
G6_HYPER_SOMNIA_64	Hypersomnia	Patient has excessive sleep (sleeps longer than 12 hours in a 24-hour period including naps) more than half the time.	<i>Have you been sleeping a lot more than usual when depressed?</i> 0 Patient has no hypersomnia. 1 Patient has excessive sleep (sleeps longer than 12 hours in a 24-hour period including naps) more than half the time. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	26	0.68	0.05
G6_DECREASED_LIBIDO_65	Decreased libido	Patient has much lower or no interest in sex or sexual activities.	<i>Has your interest in sex or your sexual activity been less than usual when depressed?</i> 0 Patient has no change in sexual activities or interest in sex. 1 Patient has much lower or no interest in sex or sexual activities. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	74	0.80	0.04

G7_ELATED_MOOD_66	Elated (euphoric) mood (Expansive mood)	Elated mood: the patient feels very happy and on top of the world. Expansive mood: the patient feels very happy and excessively friendly towards others.	<i>Have you sometimes felt very happy, elated, on top of the world without much reason?</i> 0 Patient has no elated mood. 1 Patient has elated mood less than half the time. 2 Patient has elated mood more than half the time. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	71	0.75	0.04
G7_IRRIT_MOOD_67	Irritable mood	The patient feels very easily irritated without reason to the point that he/she may shout at people or start arguments or actually become aggressive.	<i>Have you sometimes felt that you were easily irritated without reason? Have you found yourself so irritable that you shout at people or start arguments or actually become aggressive?</i> 0 Patient has no irritable mood. 1 Patient has irritable mood less than half the time. 2 Patient has irritable mood more than half the time. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	70	0.76	0.04
G7_MIXED_MOOD_68	Mixed mood (mood lability) [Same day mood changes]	Patient has periods of <u>depression and elation or irritability on the same day.</u>	<i>Have you had mixed mood swings: periods of <u>depression and elation or irritability on the same day?</u></i> 0 Patient has no mixed mood swings. 1 Patient has mixed mood less than half the time. 2 Patient has mixed mood more than half the time. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	41	0.58	0.05

G7_RACING_THOT_69	Racing thoughts	The patient has too many different thoughts racing through his/her mind compared with normal.	<i>Have you felt that you had too many different thoughts racing through your mind compared with normal?</i> 0 Patient has no racing thoughts. 1 Patient has racing thoughts less than half the time. 2 Patient has racing thoughts more than half the time. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	71	0.85	0.04
G7_PRESS_SPEECH_70	Pressured speech	Patient has been talking faster than usual during this time (for example, people said that they were unable to understand the patient because he/she was speaking too fast or the patient felt a pressure to continue talking).	<i>Have you been talking faster than usual during this time (for example, people said that they were unable to understand you because you were speaking too fast or you felt a pressure to continue talking)?</i> 0 Patient has normal speech. 1 Patient has pressured speech less than half the time. 2 Patient has pressured speech more than half the time. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	53	0.72	0.04
G7_FLIGHT_IDEAS_71	Flight of ideas	Flight of ideas (a combination of pressured speech and derailment) is observed during the clinical interview.	<i>Flight of Ideas (a combination of pressured speech and derailment):</i> 0 Patient has no flight of ideas. 1 Patient has flight of ideas. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	15	0.62	0.06

G7_MORE_ACTIVE_72	Increase in activities	The patient has been more active and more productive than usual at work or school.	<i>Have you been more active and had more energy than usual?</i> <i>Did you do more things than usual at work or school?</i> 0 Patient has no increased energy. 1 Patient has too much energy less than half the time. 2 Patient has too much energy more than half the time. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	68	0.83	0.04
G7_LESS_SLEEP_73	Decreased sleep	The patient needed to sleep four hours or less (in a 24-hour period including naps) and felt rested.	<i>Have you needed less sleep than usual and without getting tired?</i> 0 Patient has normal sleep. 1 Patient sleeps four hours or less (in a 24-hour period including naps) and feels rested. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	56	0.78	0.04
G7_DISTRACTION_74 (*)	Distraction (Attention is distracted by environmental noises)	The patient is easily distracted by unimportant activities or external noises happening in the surroundings.	<i>Do you find yourself easily distracted by unimportant activities or external noises happening around you?</i> 0 Patient has no distraction. 1 Patient has been easily distracted by external stimuli less than half the time. 2 Patient has been easily distracted by external stimuli more than half the time. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	63	0.79	0.04

G7_GRANDIOSE_75	Grandiosity	The patient feels more self-confident than usual, or he/she has special powers or special abilities.	<i>Have you felt more self-confident than usual?</i> <i>Have you felt that you have special powers or special abilities?</i> 0 Patient has no grandiosity. 1 Patient has grandiose thoughts, but not of a delusional quality. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	40	0.81	0.04
G7_OVERSPENDING_76	Overspending (poor judgment in new activities)	The patient has poor judgement with regard to engaging in new activities, usually involving money and sexuality. Money: spending more money than he/she could afford, writing bad checks, or investing money foolishly. Sexual indiscretions: e.g., having sex in a public setting.	<i>Have you done something that you regretted later (e.g., spending a lot of money that you could not afford, writing bad checks, or investing money foolishly, sexual indiscretions)?</i> 0 Patient did not go on a spending spree. 1 Patient went on a spending spree during manic phase. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	49	0.74	0.04
G7_HYPERSEXUAL_77	Hypersexuality	The patient's interest in sex or sexual activity has notably increased.	<i>Have you found that your interest in sex or your sexual activity has increased much compared with normal?</i> 0 Patient has no change in sexual activities or interest in sex. 1 Patient has been feeling hypersexual more than usual during manic phase. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	24	0.69	0.04

G7_CLANG_78	Clanging	Clanging: Words and phrases are associated together inappropriately by similarity in sound rather than meaning. Punning: association by double meaning. Assonance: association by similar syllables.	0 Normal speech. 1 Clanging: association by sound. 2 Punning: association by double meaning. 3 Assonance: association by similar syllables. 4 Mixed. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	12	0.49	0.04
--------------------	-----------------	--	---	-----------	-------------	-------------

G8_UNDER_WT_79	Being underweight	The patient has been very thin and could not maintain a minimal normal weight (low weight is defined as a weight that is less than minimally normal).	<i>Have you ever been very thin and could not maintain a minimal normal weight?</i> <i>Have people ever said you weighed much less than normal?</i> 0 = Absent or non-significant. 1 = Symptoms present. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	32	0.83	0.11
G8_WT_FEELING_80	Weight affects feelings	The patient's weight and shape are very important and affect how he/she feels about himself/herself to the point that the patient does not worry about the health risks of being so little.	<i>Do you feel that your weight and shape are very important and affect how you feel about yourself to the point that you do not worry about the health risks of being so little?</i> 0 = Absent or non-significant. 1 = Symptoms present. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	50	0.75	0.12
G8_FEAR_WT_GAIN_81	Fear of weight gain	The patient has an intense fear of gaining weight or becoming fat, even though he/she is underweight.	<i>Do you have an intense fear of gaining weight or becoming fat, even though you are underweight?</i> 0 = Absent or non-significant. 1 = Symptoms present. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	20	1.00	0.12
G8_FASTING_82	Losing weight by fasting	The patient tries to lose weight by fasting (not eating anything at all for at least 24 hours).	<i>Do you try to lose weight by fasting (not eating anything at all for at least 24 hours)?</i> 0 = Absent or non-significant. 1 = Symptoms present. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	32	0.95	0.12

G8_EXERCISE_83	Losing weight by exercise	The patient tries to lose weight by exercising too much (more than one-hour a day for at least one week).	<i>Do you try to lose weight by exercising too much (more than one hour a day for at least one week)?</i> 0 = Absent or non-significant. 1 = Symptoms present. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	22	0.86	0.12
G8_DIET_PILLS_84	Losing weight by diet pills	The patient tries to lose weight by using diet pills.	<i>Do you try to lose weight by using diet pills?</i> 0 = Absent or non-significant. 1 = Symptoms present. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	22	0.97	0.12
G8_VOMIT_85	Losing weight by vomiting	The patient tries to lose weight by inducing vomiting.	<i>Do you try to lose weight by inducing vomiting?</i> 0 = Absent or non-significant. 1 = Symptoms present. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	27	0.94	0.12
G8_LAXATIVE_86	Losing weight by laxatives	The patient tries to lose weight by taking laxatives or using enemas.	<i>Do you try to lose weight by taking laxatives or using enemas?</i> 0 = Absent or non-significant. 1 = Symptoms present. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	14	1.00	0.12
G8_OTHER_METHOD_87	Losing weight by other methods	The patient tries to lose weight by taking diuretics or by other methods.	<i>Do you try to lose weight by taking diuretics?</i> <i>Do you try to lose weight by other methods?</i> 0 = Absent or non-significant. 1 = Symptoms present. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	8 (**)	1.00	0.12

**Kappa is unstable if the number of positive cases for a given item is < 10.

G8_BINGE_88	Binge eating	The patient has episodes of binge eating (eating within one or two-hour period what most people would consider an unusually large amount of food).	<i>Do you have episodes of binge eating (eating within one or two-hour period what most people would consider an unusually large amount of food)?</i> 0 = Absent or non-significant. 1 = Symptoms present. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	27	0.97	0.12
G8_BINGE_FREQ_89	Binge eating frequency	Binge eating frequency.	0 None. 1 1-3 times per month. 2 At least once a week for 3 months. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	27	0.85	0.09
G8_LOSE_CONTRL_90	Losing control with binge eating	During the episodes of binge eating, the patient feels that he/she had lost control and could not stop eating.	<i>During the episodes of binge eating, did you feel that you had lost control and could not stop eating?</i> 0 = Absent or non-significant. 1 = Symptoms present. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	17	0.96	0.12
G8_EAT_FAST_91	Eating fast during binge eating	During the episodes of binge eating, the patient eats much more rapidly than usual.	<i>During the episodes of binge eating, did you eat much more rapidly than usual?</i> 0 = Absent or non-significant. 1 = Symptoms present. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	16	1.00	0.12
G8_EAT_FULL_92	Eating until uncomfortably full during binge eating	During the episodes of binge eating, the patient eats until he/she felt uncomfortably full.	<i>During the episodes of binge eating, did you eat until you felt uncomfortably full?</i> 0 = Absent or non-significant. 1 = Symptoms present. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	25	0.94	0.12

G8_EAT_NOT_HUNGRY_93	Eating when not hungry during binge eating	During the episodes of binge eating, the patient eats a large amount of food when he/she did not feel physically hungry.	<i>During the episodes of binge eating, did you eat a large amount of food when you did not feel physically hungry?</i> 0 = Absent or non-significant. 1 = Symptoms present. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	22	0.97	0.12
G8_EAT_ALONE_94	Eating alone during binge eating	During the episodes of binge eating, the patient eats alone because he/she was embarrassed by how much he/she was eating.	<i>During the episodes of binge eating, did you eat alone because you were embarrassed by how much you were eating?</i> 0 = Absent or non-significant. 1 = Symptoms present. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	16	0.96	0.12
G8_EAT_DISGUSTED_95	Feeling disgusted and guilty during binge eating	During the episodes of binge eating, the patient felt disgusted with himself/herself, depressed or guilty by overeating.	<i>During the episodes of binge eating, did you feel disgusted with yourself, depressed or guilty by your overeating?</i> 0 = Absent or non-significant. 1 = Symptoms present. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	22	0.86	0.12
G8_EAT_DISTRESSED_96	Distressed by overeating during binge eating	During the episodes of binge eating, the patient felt quite upset or very distressed by overeating.	<i>During the episodes of binge eating, did you feel quite upset or very distressed by your overeating?</i> 0 = Absent or non-significant. 1 = Symptoms present. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	24	0.77	0.11

G8_FASTING_B_97	Fasting after binge eating	After binge eating episodes, the patient tries to lose weight by fasting (not eating anything at all for at least 24 hours).	<i>After binge eating episodes, did you try to lose weight by fasting (not eating anything at all for at least 24 hours)?</i> 0 = Absent or non-significant. 1 = Symptoms present. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	19	0.93	0.12
G8_EXERCISE_B_98	Exercise after binge eating	After binge eating episodes, the patient tries to lose weight by exercising too much (more than one hour a day for at least one week).	<i>After binge eating episodes, did you try to lose weight by exercising too much (more than one hour a day for at least one week)?</i> 0 = Absent or non-significant. 1 = Symptoms present. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	12	0.95	0.12
G8_DIET_PILLS_B_99	Using diet pills after binge eating	After binge eating episodes, the patient tries to lose weight by using diet pills.	<i>After binge eating episodes, did you try to lose weight by using diet pills?</i> 0 = Absent or non-significant. 1 = Symptoms present. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	12	0.95	0.12
G8_VOMIT_B_100	Vomiting after binge eating	After binge eating episodes, the patient tries to lose weight by inducing vomiting.	<i>After binge eating episodes, did you try to lose weight by inducing vomiting?</i> 0 = Absent or non-significant. 1 = Symptoms present. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	17	1.00	0.12

G8_LAXATIVE_B_101	Taking laxatives after binge eating	After binge eating episodes, the patient tries to lose weight by taking laxatives or enemas.	<i>After binge eating episodes, did you try to lose weight by taking laxatives or enemas?</i> 0 = Absent or non-significant. 1 = Symptoms present. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	14	1.00	0.12
G8_OTHER_METHOD_B_102	Other losing weight methods after binge eating	After binge eating episodes, the patient tries to lose weight by taking diuretics or by other methods.	<i>After binge eating episodes, did you try to lose weight by taking diuretics? Do you try to lose weight by other methods?</i> 0 = Absent or non-significant. 1 = Symptoms present. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	9	1.00	0.12
G8_BINGE_BHV_FREQ_103	Binge eating compensatory behavior frequency	Binge eating compensatory behavior frequency.	0 None. 1 1-3 times per month. 2 At least once a week for 3 months. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	25	0.87	0.09
G8_OTHER_EATING_104	Other eating behaviors	Any other eating behaviors.	<i>Do you have any other eating behaviors?</i> 0 = Absent or non-significant. 1 = Symptoms present. . = Not sure <i>or</i> Not applicable <i>or</i> Missing.	4	0.39	0.09

G9_TRAUMA_105	Witness or experience traumatic events	The patient witnessed or experienced a traumatic event that involved actual or threatened death or serious injury to the patient or someone else (e.g., physical or sexual abuse, rape, terrorist attack, natural disaster, war...).	<i>Have you ever witnessed or experienced a traumatic event that involved actual or threatened death or serious injury to you or someone else (e.g., physical or sexual abuse, rape, terrorist attack, natural disaster, war...)?</i> <i>Did you feel intense fear and helplessness?</i> 0 Patient had no traumatic events. 1 Patient has experienced one traumatic event. 2 Patient has experienced several traumatic events. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	69	0.75	0.05
G9_TRAUMA_DISTRESS_106	Distressing recollection of events	The patient had recurrent upsetting memories (distressing recollection) of the event.	<i>Did you have recurrent upsetting memories (distressing recollection) of the event?</i> 0 Patient had no significant symptom. 1 Patient has recurrent upsetting memories (distressing recollection) of the event. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	30	0.88	0.05
G9_TRAUMA_NGTMARE_107	Bad dreams or nightmares	The patient had recurrent upsetting dreams or nightmares of the event.	<i>Did you have recurrent upsetting dreams or nightmares of the event?</i> 0 Patient had no significant symptom. 1 Patient has recurrent upsetting dreams and nightmares of the event. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	26	0.94	0.05

G9_TRAUMA_FLASH_108	Flashbacks	The patient had a sense or feeling that the event was happening again: the sense of reliving the event (flashbacks), auditory/visual hallucinations related to the event, or body/somatosensory experiences of the event.	<i>Did you have a sense or feeling that the event was happening again: the sense of reliving the event (flashbacks), auditory/visual hallucinations related to the event, or body/somatosensory experiences of the event?</i> 0 Patient had no significant symptom. 1 Patient has a sense or feeling that the event is happening again, the sense of reliving the event (flashbacks). . Not sure <i>or</i> Not applicable <i>or</i> Missing.	23	0.87	0.05
G9_TRAUMA_AVOIDTH_109	Avoidance of thoughts and feelings	The patient tries to avoid thoughts and feelings associated with the event.	<i>Did you try to avoid thoughts and feelings associated with the event?</i> 0 Patient had no significant symptom. 1 Patient tries not to think about the event. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	27	0.94	0.05
G9_TAUMA_AVOIDPLE_110	Avoidance of people, places, and activities	The patient tries to avoid things that reminded him/her of the event (such as certain people, certain places, or some activities).	<i>Did you try to avoid things that reminded you of the event (such as certain people, certain places, or some activities)?</i> 0 Patient had no significant symptom. 1 Patient avoids things that are reminders of the event (such as certain people, certain places, or some activities). . Not sure <i>or</i> Not applicable <i>or</i> Missing.	27	0.94	0.05

G9_TRAUMA_AMNESIA_111	Amnesia	The patient has difficulty remembering some or all important aspects of the event.	<i>Did you have difficulty remembering some or all important aspects of the event?</i> 0 Patient had no significant symptom. 1 Patient has difficulty remembering some or all important aspects of the event. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	15	0.70	0.06
G9_TRAUMA_ASOCIAL_112 (*)	Diminished social interest (asociality)	The patient spends less time or shows less interest in activities with friends, family or hobbies that he/she used to enjoy due to the event.	<i>Did you spend less time or show less interest in activities with friends, family or hobbies that you used to enjoy due to the event?</i> 0 Patient had no significant symptom. 1 Patient spends less time or shows less interest in activities with friends/family or hobbies due to the event. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	17	0.83	0.05
G9_TRAUMA_DETACH_113	Detachment and isolation	The patient feels distant, cut off, or isolated from other people due to the event.	<i>Did you feel distant, cut off, or isolated from other people due to the event?</i> 0 Patient had no significant symptom. 1 Patient feels distant, cut off, or isolated from other people due to the event. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	22	0.87	0.05

G9_TRAUMA_NUMB_114	Diminished emotional feelings (Diminished experience of emotions)	The patient feels emotionally numb. The patient has trouble experiencing feelings (happiness, love feelings) due to the event.	<i>Did you feel emotionally numb? Did you have trouble experiencing feelings (happiness, love feelings) due to the event?</i> 0 Patient had no significant symptom. 1 Patient feels emotionally numb. Patient has trouble experiencing feelings (such as happiness or love feelings) due to the event. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	24	0.88	0.05
G9_TRAUMA_INSOMNIA_115	Insomnia	The patient has difficulty falling or staying asleep due to the event.	<i>Did you have difficulty falling or staying asleep due to the event?</i> 0 Patient had no significant symptom. 1 Patient has difficulty falling or staying asleep due to the event. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	16	0.78	0.05
G9_TRAUMA_ANGER_116	Anger	The patient has periods of irritability or sudden outbursts of anger due to the event.	<i>Did you have periods of irritability or sudden outbursts of anger due to the event?</i> 0 Patient had no significant symptom. 1 Patient has periods of irritability or sudden outbursts of anger due to the event. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	19	0.80	0.05

G9_TRAUMA_POOR_CONC_117 (*)	Attention impairment (poor concentration)	The patient has difficulty concentrating due to the event. The patient has decreased concentration and he/she is unable to complete a task (e.g., at work, reading an article, reading a book, or watching a movie), even though the patient was able to do that before.	<i>Did you have difficulty concentrating due to the event?</i> 0 Patient had no significant symptom. 1 Patient has difficulty concentrating due to the event. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	14	0.78	0.05
G9_TRAUMA_VIGIL_118	Hypervigilance	The patient feels very alert or watchful of things going on around even when there is no need to be.	<i>Did you feel very alert or watchful of things going on around you even when there was no need to be?</i> 0 Patient had no significant symptom. 1 Patient feels very alert or watchful of things going on around even when there is no need to be. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	17	0.87	0.05
G9_TRAUMA_STARTLE_119	Startle response	The patient feels jumpy and easily startled. The patient is easily scared or makes a sudden movement or jump when he/she hears noises or caught by surprise.	<i>Did you feel jumpy and easily startled? Were you easily scared or did you make a sudden movement or jump when you heard noises or if you were caught by surprise?</i> 0 Patient had no significant symptom. 1 Patient feels jumpy and has a startle response. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	20	0.86	0.05

G9_TRAUMA_EMOTION_120	Psychological distress due to events	The patient gets emotionally upset (e.g., anxiety, agitation, shame, guilt) when something reminded him/her of the event.	<i>Did you get emotionally upset (e.g., anxiety, agitation, shame, guilt) when something reminded you of the event?</i> 0 Patient had no significant symptom. 1 Patient gets emotionally upset (e.g., anxiety, agitation, shame, guilt) when reminded of the event. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	26	0.91	0.05
G9_TRAUMA_PHYSICAL_121	Physical reactions due to events	The patient has physical reactions (e.g., fast heart beats, fast breathing, sweating) when something reminded him/her of the event.	<i>Did you have physical reactions (e.g., fast heart beats, fast breathing, sweating) when something reminded you of the event?</i> 0 Patient had no significant symptom. 1 Patient has physical reactions (e.g., fast heart beats, fast breathing, sweating) when reminded of the event. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	24	0.93	0.05
G9_TRAUMA_DAZE_122	Daze (feeling out of touch with surroundings)	The patient feels out of touch with things going on around (e.g., being in a daze or not aware of surroundings).	<i>Did you feel out of touch with things going on around you (e.g., being in a daze or not aware of surroundings)?</i> 0 Patient had no significant symptom. 1 Patient feels out of touch with things going on around. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	16	0.82	0.05

G10_AGITATION_123	Agitation	Patient may be verbally agitated causing people to be annoyed, verbally aggressive causing people to feel insulted or scared, physically agitated toward self-such as pacing, or destructive to objects.	0 Patient has no verbal agitation. 1 Patient is verbally agitated causing people to feel annoyed (e.g., makes loud noises, shouts angrily, constant whining or constant attention seeking). 2 Patient is verbally aggressive, causing people to feel insulted or scared (e.g., cursing or using foul language, makes threats to others or self). 3 Patient is physically agitated towards self (e.g., pacing up and down or disrobing). 4 Patient is destructive to objects (e.g., slams doors, throws clothes or objects, kicks wall or furniture, breaks objects, smashes windows). . Not sure <i>or</i> Not applicable <i>or</i> Missing.	33	0.48	0.04
G10_VIOLENCE_124	Violence toward others	Patient is physically violent and threatening towards people without touching victims or the patient touches victims with or without resulting injury.	0 Patient is not violent towards people. 1 Patient is physically violent and threatening towards people without touching (e.g., makes threatening gesture, swings at people). 2 Patient is physically violent and touches victims with or without resulting injury (e.g., grabs at clothes, strikes, kicks, pulls hair, attacks). . Not sure <i>or</i> Not applicable <i>or</i> Missing.	25	0.64	0.04

G10_ODD_APPEAR_125	Odd appearance and behavior	Patient wears odd clothes. Social behavior towards others is inappropriate. Sexual behavior towards others is inappropriate.	0 Patient has appropriate appearance and behaviors. 1 Patient has inappropriate appearance and behaviors. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	19	0.67	0.06
G10_INAPPRO_AFFECT_126	Inappropriate affect	<u>Inappropriate affect:</u> The patient's expression of affect is non-congruent with social or emotional context, e.g., laughing when talking about death of a loved one.	0 Patient has appropriate affect. 1 Patient has inappropriate affect. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	14	0.77	0.06
G11_ALOGIA_127	Alogia	Alogia: poverty of speech, or poverty of content of speech.	0 Patient has normal speech. 1 Patient has alogia less than half time. 2 Patient has alogia more than half time. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	29	0.62	0.05
G11_ANHEDONIA_128 (*)	Anhedonia (Loss of pleasure and interest)	The patient has been unable to experience pleasure and enjoy things that he/she used to enjoy like exercising, enjoying hobbies, or socializing with friends.	<i>Have you been unable to experience pleasure and enjoy things that you used to enjoy like exercising, enjoying your hobbies, or socializing with friends?</i> 0 Patient has no anhedonia. 1 Patient has anhedonia less than half the time. 2 Patient has anhedonia more than half the time. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	121	0.87	0.04

G11_FLAT_AFFECT_129	Blunted/flat affect (Decrease in affective expression) <u>Other descriptors in literature:</u> apathy, emotional apathy, emotional blunting, emotional indifference, blunting of emotional tone.	Decreased facial expression, e.g., not smiling or laughing at a joke, poor eye contact, indifference to things around, loss of emotional reaction and indifference.	0 Patient has broad affect. 1 Patient has blunted affect. 2 Patient has flat affect. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	42	0.68	0.05
G11_AVOLITION_130	Avolition (Decrease in goal-directed activities) <u>(Other descriptors in Literature:</u> apathy, motor apathy, general apathy, lack of drive, lack of energy, lack of interest.	Decrease in the three main goal-directed activities in life: school, work, or social activities with other people.	0 Patient has normal activities. 1 Patient has avolition less than half time. 2 Patient has avolition more than half time. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	35	0.74	0.04
G11_ASOCIAL_131 (*)	Diminished social interest (asociality)	The patient spends less time or shows less interest in activities with friends, family or hobbies that he/she used to enjoy.	<i>Did you spend less time or show less interest in activities with friends, family or hobbies that you used to enjoy?</i> 0 Patient had no significant symptom. 1 Patient spends less time or shows less interest in activities with friends, family or hobbies. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	35	0.74	0.04

G11_POOR_CONC_132 (*)	Attention impairment (poor concentration)	The patient has decreased concentration and he/she is unable to complete a task (e.g., at work, reading an article, reading a book, or watching a movie), even though the patient was able to do that before	<i>Have you found that your concentration has decreased and you are unable to complete a task (e.g., at work, reading an article, reading a book, or watching a movie), even though you were able to do that before?</i> 0 Patient has no concentration problems. 1 Patient has difficulty concentrating less than half the time. 2 Patient has difficulty concentrating more than half the time. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	41	0.92	0.12
G11_PSYCHOM_SLOW_133 (*)	Psychomotor retardation/slowing	Patient is moving slow or talking slow compared to his/her normal. Patient is observed usually sitting or lying down and talking very little.	<i>Have you felt as though you were talking or moving more slowly than normal for you?</i> 0 Patient has normal activity. 1 Patient has psychomotor retardation less than half the time. 2 Patient has psychomotor retardation more than half the time. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	97	0.72	0.04

G11_POOR_CARE_134	Poor self-care	<u>Poor self-care:</u> Patient does not wear proper clothes for the weather, does not shower regularly, does not eat properly, very dirty and disheveled.	0 Patient has proper self-care. 1 Patient has poor self-care less than half time. 2 Patient has poor self-care more than half time. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	27	0.79	0.06
G12_AUD_HAL_135	Auditory hallucinations (Hallucination quality)	The patient hears noises (like music, whispering sounds) or voices talking to him/her when there is no one around.	Do you hear noises (like music, whispering sounds) or voices talking to you when there is no one around? Are the voices like a real voice or just thoughts in your mind? 0 Patient has no auditory hallucinations. 1 Patient has auditory hallucinations. 2 Patient has auditory hallucinations with command. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	54	0.90	0.04
G12_FREQ_AUDH_136	Frequency of auditory hallucinations	The frequency is how many days per month on average the patient hears noises or voices.	How often do you hear noises (like music, whispering sounds) or voices talking to you when there is no one around? 0 Patient has no auditory hallucinations. 1 Patient has auditory hallucinations (1-4 days/ month). 2 Patient has auditory hallucinations (5-15 days/month). 3 Patient has auditory hallucinations (>15 days/month). . Not sure <i>or</i> Not applicable <i>or</i> Missing.	54	0.93	0.05

G12_DURATION_AUDH_137	Hallucination duration	The duration is how many hours per day on average the patient hears noises or voices.	On days when you hear noises or voices, how often do you hear them? 0 Patient has no auditory hallucinations. 1 Patient has auditory hallucinations (less than 1 hour / day). 2 Patient has auditory hallucinations (1-4 hours/day). 3 Patient has auditory hallucinations (more than 4 hours / day). . Not sure <i>or</i> Not applicable <i>or</i> Missing.	46	0.92	0.05
G12_AUD_THOT_138	Audible thoughts	Patient experiences auditory hallucinations, with voices speaking his/her thoughts aloud.	<i>Do you think that your thoughts are so loud that someone close to you can hear what you are thinking?</i> 0 Patient has no audible thoughts. 1 Patient has audible thoughts. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	7	1.00	0.05
G12_VOICES_ARGUE_139	Voices arguing	Two or more hallucinatory voices argue about what the patient is doing or thinking. The subject is usually the patient, who is referred to in a third person.	<i>Do you hear two or more voices that argue about what you are doing or thinking?</i> 0 Voices do not argue with the patient. 1 Voices argue about what the patient is doing or thinking. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	40	0.77	0.04

G12_VOICES_COMMENT_140	Voices commenting	One or more hallucinatory voices comment on what the patient is doing or thinking as they occur.	<i>Do you hear a voice or voices commenting on what you are doing or thinking?</i> 0 Voices do not comment about the patient. 1 Voices comment on what the patient is doing or thinking. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	40	0.77	0.04
G12_INT_HAL_141	Internal hallucinations	The patient hears the voices or noises as coming from inside his/her own head.	<i>Do you hear the voices or noises inside your head, or as though coming from outside?</i> 0 No AH 1 Mostly external hallucinations 2 Almost equal 3 Mostly internal hallucinations . Not sure <i>or</i> Not applicable <i>or</i> Missing.	50	0.84	0.04
G12_2X3HAL_142	Second/third hallucinations	The patient hears voices talking directly to him/her (second person) or talking to each other about him/her (third person).	<i>Do you hear voices talking directly to you (second person) or talking to each other about you (third person)?</i> 0 Voices do not talk to the patient or about the patient. 1 Voices talk to the patient directly most of the time. 2 Voices talk to the patient directly and talk to each other about the patient. 3 Voices talk to each other about the patient most of the time. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	45	0.78	0.04

G12_OBS_HAL_143	Observed hallucinations	Patient has been observed talking to self, talking to a mirror, or running a conversation with unseen person.	0 Patient has not been observed talking to self. 1 Patient has been observed talking to self, talking to a mirror, or running a conversation with unseen person. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	12	0.55	0.04
G12_VIS_HAL_144	Visual hallucinations	The patient sees things other people cannot see (e.g., shadows, objects or people).	<i>Do you see things other people cannot see (e.g., shadows, objects or people)?</i> 0 Patient has no visual hallucinations. 1 Patient has visual hallucinations. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	27	0.81	0.04
G12_OLF_HAL_145	Olfactory hallucinations	The patient notices unusual smells that other people do not notice. For example: patient smells flowers, while no flowers around.	<i>Do you sometimes notice unusual smells that other people do not notice?</i> 0 Patient has no olfactory hallucinations. 1 Patient has olfactory hallucinations. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	8	0.78	0.05
G12_TACT_HAL_146	Tactile hallucinations	The patient feels strange sensations on his/her body and no clear explanation for them. For example: crawling sensation under the skin, the feeling of being touched and there is no one else around.	<i>Do you sometimes feel strange sensations on your body and you have no explanation for them?</i> 0 Patient has no tactile hallucinations. 1 Patient has tactile hallucinations. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	10	0.95	0.05

G12_GUST_HAL_146A	Gustatory hallucinations	The patient experiences strange tastes in the mouth that he/she cannot explain.	<i>Do you sometimes experience strange tastes in your mouth that you cannot explain?</i> 0 Patient has no gustatory hallucinations. 1 Patient has gustatory hallucinations. . Not sure <i>or</i> Not applicable <i>or</i> Missing.			
G13_SOM_PASSIVITY_147	Somatic passivity	_Bizarre delusions associated with somatic sensations. _The patient describes bodily sensations (e.g., crawling sensation under the skin, being tortured by electricity, receiving electric shocks, sensation of burning) attributed to an outside person or force.	<i>Do you feel or have bodily sensations (e.g., something is crawling under your skin) and you think it is caused by an outside person or force?</i> 0 Patient has no somatic passivity. 1 Patient has somatic passivity. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	7	0.58	0.04
G13_THOT_INSERT_148	Thought insertion	An outside person or force is inserting a stream of thoughts to the patient's mind.	<i>Do you think that thoughts in your mind are not your own thoughts and that they were inserted into your mind by an outside person or force?</i> 0 Patient has no delusions of thought insertion. 1 Patient has delusions of thought insertion. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	16	0.76	0.04

G13_THOT_WITHDR_149	Thought withdrawal	An outside person or force is removing or taking thoughts from the patient's mind.	<i>Do you think that your thoughts were taken out of your mind by an outside person or force?</i> 0 Patient has no delusions of thought withdrawal. 1 Patient has delusions of thought withdrawal. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	6	0.80	0.04
G13_THOT_BROADCAST_150	Thought broadcasting	The patient experiences that his/her thoughts are diffused and broadcast to the outside world.	<i>Do you think that your thoughts are broadcast so that people are able to know what you are thinking even if they are in different places?</i> 0 Patient has no delusions of thought broadcast. 1 Patient has delusions of thought broadcast. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	16	0.71	0.04

G13_PARAND_ DEL_151	Paranoid/ persecutory delusions	<u>Delusion is a false belief that is held by the patient with conviction (firm belief) and despite evidence to the contrary (fixed belief).</u> The patient feels that people are against him/her or talking about or laughing at the patient. Patient may feel someone is trying to poison or kill the patient.	<i>Have you felt that people are against you, or talking about you or laughing at you?</i> <i>Do you think someone is trying to harm you (e.g., trying to poison your food or trying to kill you)?</i> 0 Patient has no paranoid/persecutory delusions. 1 Patient has paranoid/persecutory delusions some of the time. 2 Patient has paranoid/persecutory delusions most of the time. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	50	0.86	0.04
G13_CONSP_ DEL_152	Conspiracy delusions	The patient thinks there is a plot or a conspiracy against him/her by anyone (e.g. a person, FBI, CIA).	<i>Do you think there is a plot or a conspiracy against you by anyone (e.g., a person, FBI, CIA)?</i> 0 Patient has no delusions of conspiracy. 1 Patient has delusions of conspiracy some of the time. 2 Patient has delusions of conspiracy most of the time. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	49	0.84	0.04
G13_REF_DEL_ 153	Delusions of reference	When the patient watches TV, listens to the radio, or reads the newspaper, he/she thinks that special messages are intended specifically for him/her.	<i>When you are watching TV, listening to the radio, or reading the newspaper, do you think that special messages are intended specifically for you?</i> 0 Patient has no delusions of reference. 1 Patient has delusions of reference some of the time. 2 Patient has delusions of reference most of the time. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	31	0.81	0.05

G13_READ_DEL_154	Delusions of reading thoughts	Patient thinks that he/she can read people's minds, or other people can read his/her thoughts.	<i>Do you think that you can read people's minds?</i> <i>Do you think that other people can read your thoughts?</i> 0 Patient has no delusions of thought reading. 1 Patient has delusions of thought reading. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	17	0.83	0.04
G13_RELIG_DEL_155	Religious delusions	The patient has unusual religious thoughts, experiences, or practices that most friends or relatives consider very strange. The patient may feel that he/she is divine or God.	<i>Do you have unusual religious thoughts, experiences, or practices that your friends or relatives consider very strange?</i> <i>Do you feel that you yourself are divine or that you are God?</i> 0 Patient has no religious delusions. 1 Patient has religious delusions some of the time. 2 Patient has religious delusions most of the time. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	17	0.80	0.04
G13_GRAND_DEL_156	Grandiose delusions	The patient feels that he/she has special powers or great abilities that most people do not have. The patient may think that he/she is a prominent person in society.	<i>Do you feel that you have special powers or great abilities that most people do not have?</i> <i>Do you think that you are a prominent person in society?</i> 0 Patient has no grandiose delusions. 1 Patient has grandiose delusions some of the time. 2 Patient has grandiose delusions most of the time. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	16	0.77	0.05

G13_CTRL_DEL_157	Delusions of control	The patient experiences that his/her actions, impulses and emotions are controlled by an outside person or force.	<i>Do you think that some outside person or force is controlling your actions, impulses or emotions against your will?</i> 0 Patient has no delusions of being controlled. 1 Patient has delusions of being controlled. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	9	0.68	0.05
G13_OTHER_DEL_158	Other delusions	The patient has other strange thoughts or beliefs that other people do not have.	<i>Do you have any other strange thoughts or beliefs that other people do not have?</i> 0 Patient has no other delusions. 1 Patient has other delusions. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	12	0.40	0.05
G13_BIZAR_DEL_159	Bizarreness of delusions	Delusions are bizarre if they are completely impossible (e.g., patient believes he/she was born on Mars and brought to earth on a spaceship).	0 Patient has no bizarre delusions. 1 Patient has bizarre delusions. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	14	0.43	0.05

G14_DERAIL_160	Derailment (loose associations)	Derailment (loose associations): Patient's speech shifts to different topics, related or unrelated, but eventually returns to the main topic or may never return to the main topic.	0 Normal speech. 1 Patient has derailment (loose associations): speech shifts to different topics, related or unrelated, but eventually comes back to the main topic. 2 Patient has severe derailment (loose associations): speech shifts to different topics, mostly unrelated and never comes back to main topic. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	37	0.65	0.06
G14_TANGENT_161	Tangentiality	Patient's reply to a question is related in some distant way, or totally unrelated.	0 Normal speech. 1 Patient has some tangentiality: replying to a question is related in some distant way. 2 Patient has severe tangentiality: replying to a question is totally unrelated. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	28	0.57	0.06
G14_INCOHER_SP_162	Incoherent speech	Patient has incoherent speech: each sentence by itself makes sense. However, the first sentence is unrelated to the next sentence.	0 Normal speech. 1 Patient has incoherent speech: each sentence by itself makes sense. However, the first sentence is unrelated to the next sentence. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	18	0.41	0.06
G14_ILLOG_SP_163	Illogical speech	Patient's speech taken together is not logical.	0 Normal speech. 1 Patient has illogical speech: taken together, speech is not logical. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	13	0.25	0.05

G14_WORD_SALAD_164	Other disorganized thoughts (e.g., word salad, clanging)	Patient has other disorganized thoughts such as word salad (incoherence of speech at the level of the sentence, that is, the first word has nothing to do with the second word) or clanging (words and phrases are associated together inappropriately by similarity in sound rather than meaning).	0 Normal speech. 1 Other disorganized thoughts. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	39	0.65	0.04
---------------------------	---	--	--	-----------	-------------	-------------

G15_ALC_TOL_165	Alcohol tolerance	The patient needed to use a lot more alcohol than he/she previously used to get the same effect (compared when the patient first started to drink). The patient noticed that the same amount of alcohol he/she takes now has less effect than before (compared when the patient first started to drink).	<i>Did you use a lot more alcohol than you previously used to get the same effect (compared when you first started to drink)?</i> <i>Did you notice that the same amount of alcohol you take now has less effect than before (compared when you first started to drink)?</i> 0 Patient had no tolerance to alcohol. 1 Patient had tolerance to alcohol. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	39	0.99	0.06
G15_ALC_WITHDR_166	Alcohol withdrawal	When the patient stopped or cut down on alcohol use, he/she had alcohol withdrawal symptoms. Interviewer gives examples of alcohol withdrawal symptoms.	<i>When you stopped or cut down on alcohol use, did you have withdrawal symptoms? (Patient gives examples of alcohol withdrawal symptoms).</i> 0 Patient had no withdrawal symptoms from alcohol. 1 Patient had withdrawal symptoms from alcohol. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	33	0.93	0.06
G15_ALC_AVOIDW_167	Drinking alcohol to avoid withdrawal	The patient drinks alcohol to avoid withdrawal symptoms. The patient uses benzodiazepines (e.g., Ativan, Klonopin, Xanax) to avoid alcohol withdrawal symptoms.	<i>Did you drink alcohol to avoid withdrawal symptoms?</i> <i>Did you use benzodiazepines (e.g., Ativan, Klonopin, Xanax) to avoid withdrawal symptoms?</i> 0 Patient did not have to drink alcohol or use benzodiazepines to avoid withdrawal symptoms. 1 Patient drank alcohol or used benzodiazepines to avoid withdrawal symptoms. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	29	0.96	0.06

G15_UNABLE_CTRL_A_168	Unable to control alcohol	The patient drank more alcohol than he/she planned or intended. The patient drank alcohol for a longer period of time than he/she planned or intended.	<i>Did you drink more alcohol than you planned or intended?</i> <i>Did you drink alcohol for a longer period of time than you planned or intended?</i> 0 Patient had control over alcohol drinking. 1 Patient was unable to control alcohol drinking. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	51	0.96	0.06
G15_UNABLE_REDUCE_A_169	Unable to reduce or stop alcohol	The patient tried to reduce or stop alcohol use and did not succeed.	<i>Did you try to reduce or stop alcohol use and not succeed?</i> 0 Patient was able to stop or reduce alcohol drinking. 1 Patient was unable to stop or reduce alcohol drinking. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	47	0.85	0.06
G15_TIME_ALC_170	Time spent to drink alcohol	On days of alcohol use, the patient spends substantial time obtaining, using or recovering from the effect of alcohol. Alcohol drinking consumes much of the patient's time.	<i>On days of alcohol use, did you spend substantial time obtaining, using or recovering from the effect of alcohol?</i> <i>Did alcohol drinking consume much of your time?</i> 0 Patient did not lose substantial time due to alcohol. 1 Patient lost substantial time due to alcohol. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	37	0.94	0.06

G15_FAIL_OBLIG_171	Failure to fulfil major obligations	The patient's alcohol use resulted in failure to fulfill major role obligations (work, school or home).	<i>Did alcohol use result in failure to fulfill major role obligations (work, school or home)?</i> 0 Alcohol had no effect on work, school or social obligations. 1 Alcohol had negative effect on work, school or social obligations. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	36	0.92	0.06
G15_REDUCE_SOCIAL_172	Giving up social or recreational activities	The patient's alcohol use resulted in giving up or reducing important social or recreational activities.	<i>Did alcohol use result in giving up or reducing important social or recreational activities?</i> 0 Alcohol use had no effect on social or recreational activities. 1 Alcohol use had negative effect on social or recreational activities. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	36	0.92	0.06
G15_LESS_WORK_173	Less time working due to alcohol use	Alcohol use resulted in less time working (missing one or more days of work).	<i>Did alcohol use result in less time working?</i> 0 Alcohol did not result in less time working. 1 Alcohol resulted in missing one or more days of work. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	36	0.92	0.06
G15_WORK_PROB_174	Alcohol-related work problems	Alcohol use resulted in work-related problems.	<i>Did alcohol cause any work-related problems?</i> 0 Alcohol caused no work problems. 1 Alcohol caused work problems. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	29	0.83	0.06

G15_FIGHT_ALC_175	Fighting when intoxicated	The patient got in physical fights when intoxicated.	<i>Did you get in physical fights when intoxicated?</i> 0 Patient did not get in physical fights when intoxicated. 1 Patient got in physical fights when intoxicated. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	31	0.90	0.06
G15_FAMILY_PROBLM_176	Alcohol-related family problems	Alcohol use resulted in family-related problems.	<i>Did alcohol cause any family problems?</i> 0 Alcohol caused no family problems. 1 Alcohol caused family problems. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	51	0.82	0.06
G15_LEGAL_PROBLM_177	Alcohol-related legal problems	Alcohol use resulted in legal problems.	<i>Did alcohol cause any legal problems?</i> 0 Alcohol caused no legal problems. 1 Alcohol caused legal problems. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	29	0.92	0.06
G15_MED_PROBLM_178	Alcohol-induced medical problems	Alcohol use resulted in medical problems (e.g., peptic ulcer disease, liver cirrhosis).	<i>Did alcohol cause any medical problems (e.g., peptic ulcer disease, liver cirrhosis)?</i> 0 Alcohol caused no medical problems. 1 Alcohol caused medical problems. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	11	0.70	0.06
G15_EMOTION_PROBLM_179	Alcohol-induced emotional problems	Alcohol use resulted in emotional problems (e.g., depression, hallucinations).	<i>Did alcohol cause any emotional problems (e.g., depression, hallucinations)?</i> 0 Alcohol caused no emotional problems. 1 Alcohol caused emotional problems. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	24	0.90	0.06

G15_CONTIN_ALC_180	Alcohol use in spite of problems	The patient continued to use alcohol even though alcohol caused problems.	<i>Did you continue to use alcohol even though you had problems?</i> 0 Patient had no alcohol problems. 1 Patient continued to use alcohol even though alcohol caused problems. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	57	0.87	0.06
G15_ALC_HAZARD_181	Alcohol use in hazardous situations	Patient used alcohol in a situation, in which it was physically hazardous (e.g., driving a car or operating machinery).	<i>Did you use alcohol in a situation, in which it was physically hazardous (e.g., driving a car or operating machinery)?</i> 0 Patient did not use alcohol in hazardous situations. 1 Patient used alcohol in hazardous situations. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	42	0.77	0.06
G15_ALC_BINGE_182	Alcohol binge	The patient goes on binges when he/she kept drinking alcohol for a couple of days or more without sobering up.	<i>Did you go on binges when you kept drinking alcohol for a couple of days or more without sobering up?</i> 0 Patient had no binge problem. 1 Patient had binge problem. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	37	0.88	0.06
G15_ALC_BLACKOUT_183	Alcohol blackout	The patient had a blackout after drinking so much alcohol that the next day the patient could not remember what he/she said or did.	<i>Did you have a blackout after drinking so much alcohol that the next day you could not remember what you said or did?</i> 0 Patient had no blackout. 1 Patient had blackout. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	53	0.98	0.06

G16_DRG_TOL_184	Drug tolerance	The patient used a lot more of the drug than he/she previously used to get the same effect (compared when the patient first started to use the drug). The patient noticed that the same amount of the drug he/she takes now has less effect than before (compared when the patient first started to use the drug).	<i>Did you use a lot more of the drug than you previously used to get the same effect (compared when you first started to use the drug)?</i> <i>Did you notice that the same amount of the drug you take now has less effect than before (compared when you first started to use the drug)?</i> 0 Patient had no tolerance to the drug. 1 Patient had tolerance to the drug. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	49	0.95	0.06
G16_DRG_WITHDR_185	Drug withdrawal	When the patient stopped or cut down on the drug use, the patient had withdrawal symptoms of the drug. (The interviewer gives examples of the drug withdrawal symptoms).	<i>When you stopped or cut down on the drug use, did you have withdrawal symptoms?</i> <i>(Patient gives examples of the drug withdrawal symptoms).</i> 0 Patient had no withdrawal symptoms from the drug. 1 Patient had withdrawal symptoms from the drug. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	46	0.97	0.06
G16_DRG_AVOIDW_186	Using drug to avoid withdrawal	The patient had to use the drug to avoid withdrawal symptoms.	<i>Did you have to use the drug to avoid withdrawal symptoms?</i> 0 Patient did not have to use the drug to avoid withdrawal symptoms. 1 Patient had to use the drug to avoid withdrawal symptoms. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	40	0.94	0.06

G16_UNABLE_CTRL_D_187	Unable to control drug use	The patient used more of the drug than he/she planned or intended to use. The patient used the drug for a longer period of time than he/she had planned or intended.	<i>Did you use more of the drug than you planned or intended to use?</i> <i>Did you use the drug for a longer period of time than you had planned or intended?</i> 0 Patient had control over the drug use. 1 Patient used the drug more than what was planned or intended. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	55	0.97	0.06
G16_UNABLE_REDUCE_D_188	Unable to reduce or stop drug use	The patient tried to reduce or stop the drug use and he/she did not succeed.	<i>Did you try to reduce or stop the drug use and not succeed?</i> 0 Patient was able to stop or reduce the drug use. 1 Patient was unable to stop or reduce the drug use. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	54	0.97	0.06
G16_TIME_DRG_189	Time spent to use drug	On days of drug use, the patient spent substantial time obtaining, using or recovering from the effect of the drug. The drug use consumed much of the patient's time.	<i>On days of drug use, did you spend substantial time obtaining, using or recovering from the effect of the drug?</i> <i>Did drug use consume much of your time?</i> 0 Patient did not lose substantial time due to the drug use. 1 Patient lost substantial time due to the drug use. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	56	0.88	0.06

G16_FAIL_OBLIG_190	Failure to fulfil major obligations	The drug use resulted in failure to fulfill major role obligations at work, school or home.	<i>Did the drug use result in failure to fulfill major role obligations at work, school or home?</i> 0 Drug use had no effect on work, school or social obligations. 1 Drug use had a negative effect on work, school or social obligations. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	50	0.95	0.06
G16_REDUCE_SOCIAL_191	Giving up social or recreational activities	The patient's drug use resulted in giving up or reducing important social or recreational activities.	<i>Did the drug use result in giving up or reducing important social or recreational activities?</i> 0 The drug had no effect on social or recreational activities. 1 The drug had negative effect on social or recreational activities. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	50	0.95	0.06
G16_LESS_WORK_192	Less time working due to drug use	Drug use resulted in less time working (missing one or more days of work).	<i>Did drug use result in less time working?</i> 0 Drug use did not result in less time working. 1 Drug use resulted in missing one or more days of work. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	50	0.95	0.06
G16_WORK_PROBLM_193	Drug-related work problems	Drug use resulted in work-related problems.	<i>Did drug use cause any work problems?</i> 0 Drug use caused no work problems. 1 Drug use caused work problems. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	34	0.83	0.06
G16_FIGHT_DRG_194	Fighting when using drug	The patient got in physical fights when using the drug.	<i>Did you get in physical fights when using the drug?</i> 0 Patient did not get in physical fights when using the drug. 1 Patient got in physical fights when using the drug. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	22	0.80	0.06

G16_FAMILY_PROBLM_195	Drug-related family problems	Drug use resulted in family-related problems.	<i>Did drug use cause any family problems?</i> 0 Drug use caused no family problems. 1 Drug use caused family problems. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	58	0.80	0.06
G16_LEGAL_PROBLM_196	Drug-related legal problems	Drug use resulted in legal problems.	<i>Did drug use cause any legal problems?</i> 0 Drug use caused no legal problems. 1 Drug use caused legal problems. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	22	0.80	0.06
G16_MOOD_PROBLM_197	Drug-induced mood problems	Drug use resulted in mood problems (e.g., depression, mood swings).	<i>Did drug use cause any mood problems (e.g., depression, mood swings)?</i> 0 Drug use caused no mood problems. 1 Drug use caused mood problems. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	19	0.76	0.06
G16_PSYCHOSIS_PROBM_198	Drug-induced psychosis	Drug use resulted in psychotic symptoms (e.g., delusions, hallucinations).	<i>Did drug use cause any psychotic symptoms (e.g., delusions, hallucinations)?</i> 0 Drug use caused no psychotic symptoms. 1 Drug use caused psychotic symptoms. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	16	0.59	0.06

G16_CONTIN_DRUG_199	Drug use in spite of problems	The patient continued to use the drug even though he/she had problems.	<i>Did you continue to use the drug even though you had problems?</i> 0 Patient had no problems from the drug use. 1 Patient continued to use the drug even though the drug caused problems. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	64	0.91	0.06
G16_DRG_HAZARD_200	Drug use in hazardous situations	The patient used the drug in a situation, in which it was physically hazardous (e.g., driving a car or operating machinery).	<i>Did you use the drug in a situation, in which it was physically hazardous (e.g., driving a car or operating machinery)?</i> 0 Patient did not use the drug in hazardous situations. 1 Patient used the drug in hazardous situations. . Not sure <i>or</i> Not applicable <i>or</i> Missing.	57	0.90	0.06

Kappa values were calculated based upon inter-rater interviews of 322 patients at William R. Sharpe Jr. Hospital, Chestnut Ridge Center (inpatient and outpatient), Ain Shams University Hospital and Mansoura University Hospital.