

VERELST PHARMA WEIGHT LOSS - scientific references

Through our experience in genetic sequencing we found that almost half of the world population has genetic variants related with a diminished activity of the methylation cycle, which leads to increased levels of homocysteine in plasma¹. This is associated with a higher susceptibility to obesity, metabolic syndrome and diabetes²⁻⁴. Supplementation with methylated B vitamins (such as L-5-

MTHF) has been proven beneficial to reduce homocysteine levels and improve symptoms of methylation related variants^{5,6}. Pregnenolone as a precursor is a rate limiting step in the formation of all known steroid hormones. The plasma levels of pregnenolone and DHEA have been shown to decline progressively with age and thus prevention of age-related morbidities diminishes⁷. Deficiency of pregnenolone, leading to lower levels of sex steroid hormones are a known cause of increased abdominal obesity and metabolic risk profile.⁸ The combination of the methylated vitamins and the steroid hormone precursor results in long lasting weight loss.

Dietary supplement 30 capsules Net weight 10 g.

VERELST
Pharma ■■■
Designed by VERELST Research in Genetics

WEIGHT LOSS
Through a new approach we have created a smart product that combats overweight giving fast and lasting results

Dr. Sonia Van Kerckhoven
PhD in Molecular Biology and Expert in Medical Genetics

REFERENCES

1. Thawnashom K, Tungtrongchitr R, Petmitr S, et al. Methylenetetrahydrofolate reductase (mthfr) polymorphism (c677t) in relation to homocysteine. *Southeast Asian J Trop Med Public Heal.* 2005;36(2):459-466.
2. Piyathilake CJ, Badiga S, Alvarez RD, Partridge EE, Johanning GL. A Lower Degree of PBMC L1 Methylation Is Associated with Excess Body Weight and Higher HOMA-IR in the Presence of Lower Concentrations of Plasma Folate. *PLoS One.* 2013;8(1). doi:10.1371/journal.pone.0054544.
3. Kheradmand M, Maghbooli Z, Salemi S, Sanjari M. Associations of MTHFR C677T polymorphism with insulin resistance, results of NURSE Study (Nursing Unacquainted Related Stress Etiologies). *J Diabetes Metab Disord.* 2017;16(1):1-8. doi:10.1186/s40200-017-0303-9.
4. Finer S, Saravanan P, Hitman G, Yajnik C. Review Article The role of the one-carbon cycle in the developmental origins of Type 2 diabetes and obesity. *Diabet Med.* 2014;31:263-

272. doi:10.1111/dme.12390.

5. Anderson S, Panka J, Rakobitsch R, Tyre K, Pulliam K. Anxiety and Methylenetetrahydrofolate Reductase Mutation Treated With S-Adenosyl Methionine and Methylated B Vitamins. *Integr Med A Clin J*. 2016;15(2):48. <http://www.ncbi.nlm.nih.gov/pubmed/27330489><http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC4898281><https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4898281/>.
6. Bernard J, Venn Rudolf Moser, Jim I Mann TJG. Comparison of the effect of low-dose supplementation with L-5-methyltetrahydrofolate or folic acid on plasma homocysteine: a randomized placebo-controlled study. *Am Soc Clin Nutr*. 2003;77(April):658-662.
7. Sih R, Kamel H, Horani M, Morley JE. Dehydroepiandrosterone and Pregnenolone. In: *Hormone Replacement Therapy*. ; 1999:241-259.
8. Tchernof A, Després J. Sex Steroid Hormones , Sex Hormone-Binding Globulin , and Obesity in Men and Women Key words. *Horm Metab Res*. 2000;32(11/12):526-536. doi:10.1055/s-2007-978681.
9. Ma Y, Liu D. Activation of Pregnane X Receptor by Pregnenolone 16 α -carbonitrile Prevents High-Fat Diet-Induced Obesity in AKR / J Mice. *PLoS One*. 2012;7(6). doi:10.1371/journal.pone.0038734.