Post-Vaccination Discharge Instructions

Thank you for bringing your pet in for their vaccinations today! Vaccinations play a crucial role in protecting your pet from serious diseases. Below are important post-vaccine care instructions to ensure your pet's comfort and well-being.

What to Expect After Vaccination

Mild side effects are normal after vaccinations and typically resolve within 24-48 hours. Common reactions include:

- Mild lethargy or decreased activity
- Slight swelling or tenderness at the injection site
- Temporary loss of appetite
- Low-grade fever

These symptoms are **normal** and should improve on their own. Encourage your pet to rest and provide fresh water and food as usual.

When to Contact the Veterinarian

While most pets tolerate vaccines well, some may experience more serious reactions. Please **contact us immediately** if your pet shows any of the following signs:

- Persistent vomiting or diarrhea
- Swelling around the face, eyes, or muzzle
- Difficulty breathing or excessive panting
- Hives or widespread skin redness
- Weakness, collapse, or extreme lethargy

If your pet has a history of vaccine reactions, let us know before their next vaccination appointment.

Additional Care Instructions

- Avoid excessive exercise or rough play for the next 24 hours.
- Monitor the injection site for swelling, redness, or pain beyond **48 hours**.
- Keep an eye on your pet's appetite and behavior.
- If your pet received an intranasal vaccine, mild sneezing or nasal discharge may occur.

Next Vaccination Due Dates

Your pet may require booster vaccinations or additional vaccines at their next visit. Please check your pet's vaccination schedule and plan accordingly.

If you have any concerns or questions, don't hesitate to call or text us at 606-392-2030.

Thank you for prioritizing your pet's health—we look forward to seeing you at their next visit!