


Ex 6.1

Q1

To obtain the graph of $y = 3 \sin x$ we first draw the graph of $y = \sin x$ in the interval $[0, 2\pi]$. The maximum and minimum values are 3 and -3 respectively.



We have,

$$\begin{aligned}y &= 2 \sin \left(x - \frac{\pi}{4} \right) \\ \Rightarrow (y - 0) &= 2 \sin \left(x - \frac{\pi}{4} \right)\end{aligned}$$


Shifting the origin at $\left(\frac{\pi}{4}, 0\right)$, we have:

$$x = X + \frac{\pi}{4} \text{ and } y = Y + L$$

Substituting these values in (i), we get

$$Y = 2 \sin X$$

Thus we draw the graph of $Y = 2 \sin X$ and shift it by $\frac{\pi}{4}$ to the right to get the required graph.





We have,

$$\begin{aligned}y &= 2 \sin(2x - 1) \\&\Rightarrow (y - 0)' = 2 \sin 2 \left(x - \frac{1}{2}\right)\end{aligned}$$


Shifting the origin at $\left(\frac{1}{2}, 0\right)$, we have

$$x = X + \frac{1}{2} \text{ and } y = Y - 0$$

Substituting these values in (i), we get

$$Y = 2 \sin 2X$$

Thus we draw the graph of $Y = 2 \sin 2X$ and shift it by $1/2$ to the right to get the required graph.



We have,

$$\begin{aligned}y &= 3 \sin(3x + 1) \\&\Rightarrow (y - 0)' = 3 \sin 3 \left(x + \frac{1}{3}\right)\end{aligned}$$


Shifting the origin at $\left(-\frac{1}{3}, 0\right)$, we have

$$x = X - \frac{1}{3} \text{ and } y = Y + 0$$

Substituting these values in (i), we get

$$Y = 3 \sin 3X$$

Thus we draw the graph of $Y = 3 \sin 3X$ and shift it by $1/3$ to the left to get the required graph.





We have,

$$y = 3 \sin \left(2x - \frac{\pi}{4} \right)$$
$$\Rightarrow (y - C) = 3 \sin 2 \left(x - \frac{\pi}{8} \right)$$


Shifting the origin at $\left(\frac{\pi}{8}, 0 \right)$, we have

$$x = X + \frac{\pi}{8} \text{ and } y = Y + C$$

Substituting these values in (i), we get


$$Y = 3 \sin 2X$$

Thus we draw the graph of $Y = 3 \sin 2X$ and shift it by $\frac{\pi}{8}$ to the right to get the required graph.





Q2



We have,

$$y = \sin\left(x + \frac{\pi}{4}\right)$$
$$\Rightarrow y - 0 = \sin\left(x - \frac{\pi}{4}\right) \quad (i)$$

Shifting the origin at $\left(-\frac{\pi}{4}, 0\right)$, we obtain


$$x = X - \frac{\pi}{4}, \quad y = Y + 0$$

Substituting these values in (i), we get

$$Y = \sin X.$$

Thus we draw the graph of $Y = \sin X$ and shift it by $\frac{\pi}{4}$ to the left to get the required graph.

To obtain the graph of $y = \sin 3x$ we first draw the graph of $y = \sin x$ in the interval $[0, 2\pi]$ and then divide the x-coordinates of the points where it crosses x-axis by 3.





Ex 6.2

Q1

We have,

$$y = \cos\left(x - \frac{\pi}{4}\right)$$

$$\Rightarrow y - 1 = \cos\left(x + \frac{\pi}{4}\right) \quad \text{--- i)}$$


Shifting the origin at $\left(-\frac{\pi}{4}, 0\right)$, we obtain

$$x = X - \frac{\pi}{4}, \quad y = Y + 0$$

Substituting these values in i), we get

$$Y = \cos X.$$

Thus we draw the graph of $Y = \cos X$ and shift it by $\frac{\pi}{4}$ to the left to get the required graph.



We have,

$$y = \cos\left(x + \frac{\pi}{4}\right)$$

$$\Rightarrow y - 0 = \cos\left(x + \frac{\pi}{4}\right) \quad \text{--- ii)}$$


Shifting the origin at $\left(\frac{\pi}{4}, 0\right)$, we obtain

$$x = X + \frac{\pi}{4}, \quad y = Y + 0$$

Substituting these values in ii), we get

$$Y = \cos X.$$

Thus we draw the graph of $Y = \cos X$ and shift it by $\frac{\pi}{4}$ to the right to get the required graph.





We have,

$$\begin{aligned}y &= 3 \cos(2x - 1) \\ \Rightarrow (y - C) &= 3 \cos 2\left(x - \frac{1}{2}\right)\end{aligned}$$


Shifting the origin at $\left(\frac{1}{2}, 0\right)$, we have

$$x = X + \frac{1}{2} \text{ and } y = Y + C$$

Substituting these values in (i), we get

$$Y = 3 \cos 2X$$

Thus we draw the graph of $Y = 3 \cos 2X$ and shift it by $\frac{1}{2}$ to the right to get the required graph.



We have,

$$\begin{aligned}y &= 2 \cos\left(x - \frac{\pi}{2}\right) \\ \Rightarrow y - 0 &= 2 \cos\left(x - \frac{\pi}{2}\right) \quad \text{---(i)}\end{aligned}$$


Shifting the origin at $\left(\frac{\pi}{2}, 0\right)$ we obtain

$$x = X + \frac{\pi}{2}, \quad y = Y + C$$

Substituting these values in (i), we get


$$Y = 2 \cos X.$$

Thus we draw the graph of $Y = 2 \cos X$ and shift it by $\frac{\pi}{2}$ to the right to get the required graph.





Q2



We have,

$$\begin{aligned}y &= \cos 2\left(x - \frac{\pi}{4}\right) \\ \Rightarrow y - 0 &= \cos 2\left(x - \frac{\pi}{4}\right)\end{aligned}$$


shifting the origin at $\left(\frac{\pi}{4}, 0\right)$, we obtain

$$x = X + \frac{\pi}{4}, \quad y = Y + 0$$

Substituting these values in (i), we get


$$Y = \cos 2X.$$

Thus we draw the graph of $Y = \cos 2X$ and shift it by $\frac{\pi}{4}$ to the right to get the required graph.





To obtain the graph of $y = \cos \frac{x}{2}$ we first draw the graph of $y = \cos x$ in the interval $[0, 2\pi]$ and then divide the x-coordinates of the points where it crosses x-axis by 1/2.





Ex 6.3

Q1

We know that

$$y = \sin^2 x = \frac{1 - \cos 2x}{2} = \frac{1}{2} - \frac{1}{2} \cos 2x$$

We have,

$$\begin{aligned} y &= \frac{1}{2} - \frac{1}{2} \cos 2x \\ \Rightarrow y - \frac{1}{2} &= -\frac{1}{2} \cos 2x \end{aligned} \quad (i)$$


Shifting the origin at $\left(0, -\frac{1}{2}\right)$, we obtain

$$x = X, y = Y + \frac{1}{2}$$

Substituting these values in (i), we get

$$Y = -\frac{1}{2} \cos 2X.$$

Thus we draw the graph of $Y = \cos 2X$, adjust the maximum and minimum values to $1/2$ and $-1/2$ and shift it by $\frac{1}{2}$ up to get the required graph.





Q2

We know that

$$y = \cos^2 x = \frac{1 + \cos 2x}{2} = \frac{1}{2} + \frac{1}{2} \cos 2x$$

We have,

$$\begin{aligned} y &= \frac{1}{2} + \frac{1}{2} \cos 2x \\ \Rightarrow y - \frac{1}{2} &= \frac{1}{2} \cos 2x \end{aligned} \quad \text{---(i)}$$


shifting the origin at $\left(0, -\frac{1}{2}\right)$, we obtain

$$x = X, y = Y + \frac{1}{2}$$

Substituting these values in (i), we get

$$Y = -\frac{1}{2} \cos 2X$$

Thus we draw the graph of $Y = \cos 2X$, adjust the maximum and minimum values to 1/2 and -1/2 and shift it by $\frac{1}{2}$ down to get the required graph.





Q3

We have,

$$y = \sin^2\left(x - \frac{\pi}{4}\right)$$
$$\rightarrow y - 0 = \sin^2\left(x - \frac{\pi}{4}\right) \quad \text{--- (i)}$$


Shifting the origin at $\left(\frac{\pi}{4}, 0\right)$, we obtain

$$x = X + \frac{\pi}{4}, \quad y = Y + 0$$

Substituting these values in (i), we get

$$Y = \sin^2 X,$$


Thus we draw the graph of $Y = \sin^2 X$ and shift it by $\frac{\pi}{4}$ to the right to get the required graph.






Q4

To obtain the graph of $y = \tan 2x$ we first draw the graph of $y = \tan x$ in the interval $\left(-\frac{\pi}{2}, -\frac{\pi}{2}\right)$ and then divide the x-coordinates of the points where it crosses x-axis by 2.



Q5


To obtain the graph of $y = 2 \tan 3x$ we first draw the graph of $y = \tan x$ in the interval $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ and then divide the x-coordinates of the points where it crosses x-axis by 3. We then stretch the graph vertically by a factor of 2.





Q6

To obtain the graph of $y = 2 \cot 2x$ we first draw the graph of $y = \cot x$ in the interval $(0, \pi)$ and then divide the x-coordinates of the points where it crosses x-axis by 2. We then stretch the graph vertically by a factor of 2.





Q7

We have,

$$y = \cos 2\left(x - \frac{\pi}{6}\right)$$
$$\Rightarrow y - 0 = \cos 2\left(x - \frac{\pi}{6}\right) \quad \text{--- (i)}$$


Shifting the origin at $\left(\frac{\pi}{6}, 0\right)$, we obtain

$$x = X + \frac{\pi}{6}, \quad y = Y + 0$$

Substituting these values in (i), we get

$$Y = \cos 2X.$$

Thus we draw the graph of $Y = \cos 2X$ and shift it by $\frac{\pi}{6}$ to the right to get the required graph.





Q8

We know that

$$y = \sin^2 x = \frac{1 - \cos 2x}{2} = \frac{1}{2} - \frac{1}{2} \cos 2x$$

We have,

$$\begin{aligned} y &= \frac{1}{2} - \frac{1}{2} \cos 2x \\ \Rightarrow y - \frac{1}{2} &= -\frac{1}{2} \cos 2x \quad \cdots (i) \end{aligned}$$

Shifting the origin at $\left(0, -\frac{1}{2}\right)$, we obtain


$$x = X, \quad y = Y + \frac{1}{2}$$

Substituting these values in (i), we get

$$Y = -\frac{1}{2} \cos 2X.$$


Thus we draw the graph of $Y = \cos 2X$, adjust the maximum and minimum values to $1/2$ and $-1/2$

and shift it by $\frac{1}{2}$ up to get the required graph.





Q9



Q10

