

SYSTEM OF LINEAR EQUATIONS (XII, R. S. AGGARWAL)

EXERCISE 8A (Pg. No.: 310)

Show that each one of the following systems of equations is inconsistent.

1. $x + 2y = 9; 2x + 4y = 7$

Sol. $A = \begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix}, X = \begin{bmatrix} x \\ y \end{bmatrix}, B = \begin{bmatrix} 9 \\ 7 \end{bmatrix}$ $|A| = \begin{vmatrix} 1 & 2 \\ 2 & 4 \end{vmatrix} = 4 - 4 = 0 \therefore \text{adj } A = \begin{bmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{bmatrix}'$

So the cofactor of the element of matrix A are $C_{11} = 4, C_{12} = -2, C_{21} = -2, C_{22} = 1$

$$\text{adj } A = \begin{bmatrix} 4 & -2 \\ -2 & 1 \end{bmatrix}' \Rightarrow \text{adj } A = \begin{bmatrix} 4 & -2 \\ -2 & 1 \end{bmatrix}$$

Now, $(\text{adj } A) \cdot B = \begin{bmatrix} 4 & -2 \\ -2 & 1 \end{bmatrix} \begin{bmatrix} 9 \\ 7 \end{bmatrix} = \begin{bmatrix} 36 - 14 \\ -18 + 7 \end{bmatrix} = \begin{bmatrix} 22 \\ -11 \end{bmatrix} \neq 0$

Hence, the system has no solution, it is inconsistent.

2. $2x + 3y = 5; 6x + 9y = 10$

Sol. $A = \begin{bmatrix} 2 & 3 \\ 6 & 9 \end{bmatrix}, X = \begin{bmatrix} x \\ y \end{bmatrix}, B = \begin{bmatrix} 5 \\ 10 \end{bmatrix} \therefore \text{adj } A = \begin{bmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{bmatrix}'$

So the cofactor of the element of matrix A are $C_{11} = 9, C_{12} = -6, C_{21} = -3, C_{22} = 2$

$$\text{adj } A = \begin{bmatrix} 9 & -6 \\ -3 & 2 \end{bmatrix}' \Rightarrow \text{adj } A = \begin{bmatrix} 9 & -3 \\ -6 & 2 \end{bmatrix}$$

Now, $\text{adj}(A) \times B = \begin{bmatrix} 9 & -3 \\ -6 & 2 \end{bmatrix} \begin{bmatrix} 5 \\ 10 \end{bmatrix} = \begin{bmatrix} 45 - 30 \\ -30 + 20 \end{bmatrix} = \begin{bmatrix} 15 \\ -10 \end{bmatrix} \neq 0$

Hence, the system has no solution, it is inconsistent.

3. $4x - 2y = 3; 6x - 3y = 5$

Sol. $A = \begin{bmatrix} 4 & -2 \\ 6 & -3 \end{bmatrix}, X = \begin{bmatrix} x \\ y \end{bmatrix}, B = \begin{bmatrix} 3 \\ 5 \end{bmatrix}, |A| = \begin{vmatrix} 4 & -2 \\ 6 & -3 \end{vmatrix} = -12 + 12 = 0 \therefore \text{adj } A = \begin{bmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{bmatrix}'$

So the cofactor of the element of matrix A are $C_{11} = -3, C_{12} = -6, C_{21} = 2, C_{22} = 4$

$$\text{adj } A = \begin{bmatrix} -3 & -6 \\ 2 & 4 \end{bmatrix}' \Rightarrow \text{adj } A = \begin{bmatrix} -3 & 2 \\ -6 & 4 \end{bmatrix}$$

Now, $(\text{adj } A) \cdot B = \begin{bmatrix} -3 & 2 \\ -6 & 4 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \end{bmatrix} = \begin{bmatrix} -9 + 10 \\ -18 + 20 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix} \neq 0$

Hence, the system has no solution, it is inconsistent.

4. $6x + 4y = 5; 9x + 6y = 8$

Sol. $A = \begin{bmatrix} 6 & 4 \\ 9 & 6 \end{bmatrix}, X = \begin{bmatrix} x \\ y \end{bmatrix}, B = \begin{bmatrix} 5 \\ 8 \end{bmatrix}, |A| = \begin{vmatrix} 6 & 4 \\ 9 & 6 \end{vmatrix} = 36 - 36 = 0 \therefore \text{adj } A = \begin{bmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{bmatrix}'$

So the cofactor of the element of matrix A are $C_{11} = 6, C_{12} = -9, C_{21} = -4, C_{22} = 6$

$$\therefore \text{adj } A = \begin{bmatrix} 6 & -9 \\ -4 & 6 \end{bmatrix}' = \begin{bmatrix} 6 & -4 \\ -9 & 6 \end{bmatrix}$$

$$\text{Now, } (\text{adj } A) \cdot B = \begin{bmatrix} 6 & -4 \\ -9 & 6 \end{bmatrix} \begin{bmatrix} 5 \\ 8 \end{bmatrix} = \begin{bmatrix} 30-32 \\ -45+48 \end{bmatrix} = \begin{bmatrix} -2 \\ 3 \end{bmatrix} \neq 0.$$

Hence, the system has no solution, it is inconsistent.

5. $x + y - 2z = 5; x - 2y + z = -2, -2x + y + z = 4$

Sol. $A = \begin{bmatrix} 1 & 1 & -2 \\ 1 & -2 & 1 \\ -2 & 1 & 1 \end{bmatrix}, X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, B = \begin{bmatrix} 5 \\ -2 \\ 4 \end{bmatrix},$

$$|A| = \begin{vmatrix} 1 & 1 & -2 \\ 1 & -2 & 1 \\ -2 & 1 & 1 \end{vmatrix} = 1 \begin{vmatrix} -2 & 1 \\ 1 & 1 \end{vmatrix} - 1 \begin{vmatrix} 1 & 1 \\ -2 & 1 \end{vmatrix} - 2 \begin{vmatrix} 1 & -2 \\ -2 & 1 \end{vmatrix} = 0$$

$$\therefore \text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}'$$

So the cofactor of the element of matrix A are, $\begin{bmatrix} C_{11} = -3 & C_{12} = -3 & C_{13} = -3 \\ C_{21} = -3 & C_{22} = -3 & C_{23} = -3 \\ C_{31} = -3 & C_{32} = -3 & C_{33} = -3 \end{bmatrix}$

$$\Rightarrow \text{adj } A = \begin{bmatrix} -3 & -3 & -3 \\ -3 & -3 & -3 \\ -3 & -3 & -3 \end{bmatrix}' \Rightarrow \text{adj } A = \begin{bmatrix} -3 & -3 & -3 \\ -3 & -3 & -3 \\ -3 & -3 & -3 \end{bmatrix}$$

$$\text{Now, } (\text{adj } A) \cdot B = \begin{bmatrix} -3 & -3 & -3 \\ -3 & -3 & -3 \\ -3 & -3 & -3 \end{bmatrix} \begin{bmatrix} 5 \\ -2 \\ 4 \end{bmatrix} = \begin{bmatrix} -15+6-12 \\ -15+6-12 \\ -15+6-12 \end{bmatrix} = \begin{bmatrix} -21 \\ -21 \\ -21 \end{bmatrix} \neq 0.$$

Hence, the system has no solution, it is inconsistent.

6. $2x - y + 3z = 1; 3x - 2y + 5z = -4; 5x - 4y + 9z = 14$

Sol. $A = \begin{bmatrix} 2 & -1 & 3 \\ 3 & -2 & 5 \\ 5 & -4 & 9 \end{bmatrix}, X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, B = \begin{bmatrix} 1 \\ -4 \\ 14 \end{bmatrix},$

$$|A| = \begin{vmatrix} 2 & -1 & 3 \\ 3 & -2 & 5 \\ 5 & -4 & 9 \end{vmatrix} = 2 \begin{vmatrix} -2 & 5 \\ -4 & 9 \end{vmatrix} - (-1) \begin{vmatrix} 3 & 5 \\ 5 & 9 \end{vmatrix} + 3 \begin{vmatrix} 3 & -2 \\ 5 & -4 \end{vmatrix} = 2(2) + (2) + 3(-2) = 4 + 2 - 6 = 0$$

$$\therefore \text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}'$$

<https://millionstar.godaddy.com/>

So the cofactor of the element of matrix A are,
$$\begin{bmatrix} C_{11} = 2 & C_{12} = -2 & C_{13} = -2 \\ C_{21} = -3 & C_{22} = 3 & C_{23} = 3 \\ C_{31} = 1 & C_{32} = -1 & C_{33} = -1 \end{bmatrix}$$

$$\text{adj } A = \begin{bmatrix} 2 & -2 & -2 \\ -3 & 3 & 3 \\ 1 & -1 & -1 \end{bmatrix}' \Rightarrow \text{adj } A = \begin{bmatrix} 2 & -3 & 1 \\ -2 & 3 & -1 \\ -2 & 3 & -1 \end{bmatrix}$$

$$\text{Now, } (\text{adj } A) \cdot B = \begin{bmatrix} 2 & -3 & 1 \\ -2 & 3 & -1 \\ -2 & 3 & -1 \end{bmatrix} \begin{bmatrix} 1 \\ -4 \\ 14 \end{bmatrix} = \begin{bmatrix} 2+12+14 \\ -2-12-14 \\ -2-12-14 \end{bmatrix} = \begin{bmatrix} 28 \\ -28 \\ -28 \end{bmatrix} \neq 0$$

Hence, the solution has no solution, it is inconsistent.

7. $x + 2y + 4z = 12$; $y + 2z = -1$; $3x + 2y + 4z = 4$

Sol. $A = \begin{bmatrix} 1 & 2 & 4 \\ 0 & 1 & 2 \\ 3 & 2 & 4 \end{bmatrix}$, $X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$, $B = \begin{bmatrix} 12 \\ -1 \\ 4 \end{bmatrix}$,

$$|A| = \begin{vmatrix} 1 & 2 & 4 \\ 0 & 1 & 2 \\ 3 & 2 & 4 \end{vmatrix} = 1 \begin{vmatrix} 1 & 2 \\ 2 & 4 \end{vmatrix} - 2 \begin{vmatrix} 0 & 2 \\ 3 & 4 \end{vmatrix} + 4 \begin{vmatrix} 0 & 1 \\ 3 & 2 \end{vmatrix} = (4-4) - 2(0-6) + 4(0-3) = 0 + 12 - 12 = 0$$

$$\therefore \text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}'$$

So the cofactor of the element of matrix A are,
$$\begin{bmatrix} C_{11} = 0 & C_{12} = 6 & C_{13} = -3 \\ C_{21} = 0 & C_{22} = -8 & C_{23} = 4 \\ C_{31} = 0 & C_{32} = -2 & C_{33} = 1 \end{bmatrix}$$

$$\therefore \text{adj } A = \begin{bmatrix} 0 & 6 & -3 \\ 0 & -8 & 4 \\ 0 & -2 & 1 \end{bmatrix}' \Rightarrow \text{adj } A = \begin{bmatrix} 0 & 0 & 0 \\ 6 & -8 & -2 \\ -3 & 4 & 1 \end{bmatrix}$$

$$\text{Now, } (\text{adj } A) \cdot B = \begin{bmatrix} 0 & 0 & 0 \\ 6 & -8 & -2 \\ -3 & 4 & 1 \end{bmatrix} \begin{bmatrix} 12 \\ -1 \\ 4 \end{bmatrix} = \begin{bmatrix} 0-0+0 \\ 72+8-8 \\ -36-4+4 \end{bmatrix} = \begin{bmatrix} 0 \\ 72 \\ -36 \end{bmatrix} \neq 0$$

Hence, the system has no solution, it is inconsistent.

8. $3x - y - 2z = 2$; $2y - z = -1$; $3x - 5y = 3$

Sol. $A = \begin{bmatrix} 3 & -1 & -2 \\ 0 & 2 & -1 \\ 3 & -5 & 0 \end{bmatrix}$, $X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$, $B = \begin{bmatrix} 2 \\ -1 \\ 3 \end{bmatrix}$,

$$|A| = \begin{vmatrix} 3 & -1 & -2 \\ 0 & 2 & -1 \\ 3 & -5 & 0 \end{vmatrix} = 3 \begin{vmatrix} 2 & -1 \\ -5 & 0 \end{vmatrix} + 1 \begin{vmatrix} 0 & -1 \\ 3 & 0 \end{vmatrix} - 2 \begin{vmatrix} 0 & 2 \\ 3 & -5 \end{vmatrix} \\ = 3(0-5) + 3 - 2(0-6) = -15 + 3 + 12 = 0$$

<https://millionstar.godaddysites.com/>

$$\therefore \text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}$$

So the cofactor of the element of matrix A are, $\begin{bmatrix} C_{11} = -5 & C_{12} = -3 & C_{13} = -6 \\ C_{21} = 10 & C_{22} = 6 & C_{23} = 12 \\ C_{31} = 5 & C_{32} = 3 & C_{33} = 6 \end{bmatrix}$

$$\text{adj } A = \begin{bmatrix} -5 & -3 & -6 \\ 10 & 6 & 12 \\ 5 & 3 & 6 \end{bmatrix} \Rightarrow \text{adj } A = \begin{bmatrix} -5 & 10 & 5 \\ -3 & 6 & 3 \\ -6 & 12 & 6 \end{bmatrix}$$

$$\text{Now, } (\text{adj } A) \cdot B = \begin{bmatrix} -5 & 10 & 5 \\ -3 & 6 & -3 \\ -6 & 12 & 6 \end{bmatrix} \begin{bmatrix} 2 \\ -1 \\ 3 \end{bmatrix} = \begin{bmatrix} -10 - 10 + 15 \\ -6 - 6 + 9 \\ -12 - 12 + 18 \end{bmatrix} = \begin{bmatrix} -5 \\ -3 \\ -6 \end{bmatrix} \neq 0$$

Hence, the system has no solution, it is inconsistent.

Solve each of the following systems of equations using matrix method.

9. $5x + 2y = 4$; $7x + 3y = 5$

Sol. $A = \begin{bmatrix} 5 & 2 \\ 7 & 3 \end{bmatrix}$, $X = \begin{bmatrix} x \\ y \end{bmatrix}$, $B = \begin{bmatrix} 4 \\ 5 \end{bmatrix}$, $|A| = \begin{vmatrix} 5 & 2 \\ 7 & 3 \end{vmatrix} = 15 - 14 = 1 \Rightarrow |A| \neq 0$. Hence, A^{-1} exists.

$$\therefore \text{adj } A = \begin{bmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{bmatrix}$$

So the cofactor of the element of matrix A are, $C_{11} = 3$, $C_{21} = -2$, $C_{12} = -7$, $C_{22} = 5$

$$\text{adj } A = \begin{bmatrix} 3 & -7 \\ -2 & 5 \end{bmatrix} \Rightarrow \text{adj } A = \begin{bmatrix} 3 & -2 \\ -7 & 5 \end{bmatrix} \text{ and } |A| = \begin{vmatrix} 5 & 2 \\ 7 & 3 \end{vmatrix} = (15 - 14) = 1$$

$$\therefore A^{-1} = \frac{1}{|A|} (\text{adj } A) = \frac{1}{1} \begin{bmatrix} 3 & -2 \\ -7 & 5 \end{bmatrix} = \begin{bmatrix} 3 & -2 \\ -7 & 5 \end{bmatrix}$$

$$\text{Now, } AX = B \Rightarrow X = A^{-1}B \Rightarrow X = \begin{bmatrix} 3 & -2 \\ -7 & 5 \end{bmatrix} \begin{bmatrix} 4 \\ 5 \end{bmatrix} \Rightarrow X = \begin{bmatrix} 12 - 10 \\ -28 + 25 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2 \\ -3 \end{bmatrix}$$

Hence, $x = 2$ and $y = -3$, it is inconsistent.

10. $3x + 4y - 5 = 0$; $x - y + 3 = 0$

Sol. $A = \begin{bmatrix} 3 & 4 \\ 1 & -1 \end{bmatrix}$, $X = \begin{bmatrix} x \\ y \end{bmatrix}$, $B = \begin{bmatrix} 5 \\ -3 \end{bmatrix}$ $\therefore \text{adj } A = \begin{bmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{bmatrix}$

So the cofactor of the element of matrix A are, $\begin{bmatrix} C_{11} = -1 & C_{21} = -4 \\ C_{12} = -1 & C_{22} = 3 \end{bmatrix}$

$$\text{adj } A = \begin{bmatrix} -1 & -1 \\ -4 & 3 \end{bmatrix} \Rightarrow \text{adj } A = \begin{bmatrix} -1 & -4 \\ -1 & 3 \end{bmatrix} \text{ and } |A| = \begin{vmatrix} 3 & 4 \\ 1 & -1 \end{vmatrix} = (-3 - 4) = -7 \neq 0$$

$$\text{and } A^{-1} = \frac{1}{|A|} (\text{adj } A) = \frac{1}{(-7)} \begin{bmatrix} -1 & -4 \\ -1 & 3 \end{bmatrix} = \frac{1}{7} \begin{bmatrix} 1 & 4 \\ 1 & -3 \end{bmatrix}$$

<https://millionstar.godaddy.com/>

$$\text{Now, } AX = B \Rightarrow X = A^{-1}B \Rightarrow X = \frac{1}{7} \begin{bmatrix} 1 & 5 \\ 1 & -3 \end{bmatrix} \begin{bmatrix} 1 \\ -3 \end{bmatrix}$$

$$\Rightarrow X = \frac{1}{7} \begin{bmatrix} 5-12 \\ 5+9 \end{bmatrix} \Rightarrow X = \frac{1}{7} \begin{bmatrix} -7 \\ 14 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -1 \\ 2 \end{bmatrix}. \quad \text{Hence, } x = -1 \text{ and } y = 2.$$

11. $x + 2y = 1; 3x + y = 4$

Sol. $A = \begin{bmatrix} 1 & 2 \\ 3 & 1 \end{bmatrix}, X = \begin{bmatrix} x \\ y \end{bmatrix}, B = \begin{bmatrix} 1 \\ 4 \end{bmatrix} \therefore \text{adj } A = \begin{bmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{bmatrix}'$

So the cofactor of the element of matrix A are, $\begin{bmatrix} C_{11} = 1 & C_{21} = -2 \\ C_{12} = -3 & C_{22} = 1 \end{bmatrix}$

$$\text{adj } A = \begin{bmatrix} 1 & -3 \\ -2 & 1 \end{bmatrix}' \Rightarrow \text{adj } A = \begin{bmatrix} 1 & -2 \\ -3 & 1 \end{bmatrix} \text{ and } |A| = \begin{vmatrix} 1 & 2 \\ 3 & 1 \end{vmatrix} \Rightarrow |A| = (1-6) = (-5)$$

$$\text{and } A^{-1} = \frac{1}{|A|} (\text{adj } A) = \frac{1}{(-5)} \begin{bmatrix} 1 & -2 \\ -3 & 1 \end{bmatrix} = \frac{1}{5} \begin{bmatrix} -1 & 2 \\ 3 & -1 \end{bmatrix}$$

$$\text{Now } AX = B \Rightarrow X = A^{-1}B \Rightarrow X = \frac{1}{5} \begin{bmatrix} -1 & 2 \\ 3 & -1 \end{bmatrix} \begin{bmatrix} 1 \\ 4 \end{bmatrix} \Rightarrow X = \frac{1}{5} \begin{bmatrix} -1+8 \\ 3-4 \end{bmatrix}$$

$$\Rightarrow X = \frac{1}{5} \begin{bmatrix} 7 \\ -1 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 7/5 \\ -1/5 \end{bmatrix}. \quad \text{Hence, } x = \frac{7}{5} \text{ and } y = -\frac{1}{5}.$$

12. $5x + 7y + 2 = 0; 4x + 6y + 3 = 0$

Sol. $A = \begin{bmatrix} 5 & 7 \\ 4 & 6 \end{bmatrix}, B = \begin{bmatrix} -2 \\ -3 \end{bmatrix}, X = \begin{bmatrix} x \\ y \end{bmatrix}, |A| = \begin{vmatrix} 5 & 7 \\ 4 & 6 \end{vmatrix} = (30-28) = 2. \text{ Hence, } A^{-1} \text{ exist.}$

$$\therefore \text{adj } A = \begin{bmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{bmatrix}'$$

So the cofactor of the element of matrix A are, $\begin{bmatrix} C_{11} = 6 & C_{21} = -7 \\ C_{12} = -4 & C_{22} = 5 \end{bmatrix}$

$$\text{adj } A = \begin{bmatrix} 6 & -4 \\ -7 & 5 \end{bmatrix}' = \begin{bmatrix} 6 & -7 \\ -4 & 5 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{|A|} (\text{adj } A) = \frac{1}{2} \begin{bmatrix} 6 & -7 \\ -4 & 5 \end{bmatrix}$$

$$\text{Now, } AX = B \Rightarrow X = A^{-1}B \Rightarrow X = \frac{1}{2} \begin{bmatrix} 6 & -7 \\ -4 & 5 \end{bmatrix} \begin{bmatrix} -2 \\ -3 \end{bmatrix} \Rightarrow X = \frac{1}{2} \begin{bmatrix} -12+21 \\ 8-15 \end{bmatrix}$$

$$\Rightarrow X = \frac{1}{2} \begin{bmatrix} 9 \\ -7 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 9/2 \\ -7/2 \end{bmatrix}.$$

$$\text{Hence, } x = \frac{9}{2} \text{ \& } y = -\frac{7}{2}$$

13. $2x - 3y + 1 = 0; x + 4y + 3 = 0$

Sol. $A = \begin{bmatrix} 2 & -3 \\ 1 & 4 \end{bmatrix}, X = \begin{bmatrix} x \\ y \end{bmatrix}, B = \begin{bmatrix} -1 \\ -3 \end{bmatrix}, |A| = \begin{vmatrix} 2 & -3 \\ 1 & 4 \end{vmatrix} = (8+3) = 11. \text{ Hence, } A^{-1} \text{ exist.}$

<https://millionstar.godaddysites.com/>

$$\therefore \text{adj } A = \begin{bmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{bmatrix}'$$

So the cofactor of the element of matrix A are, $\begin{bmatrix} C_{11} = 4 & C_{21} = 3 \\ C_{12} = -1 & C_{22} = 2 \end{bmatrix}$

$$\text{adj } A = \begin{bmatrix} 4 & -1 \\ 3 & 2 \end{bmatrix}' \Rightarrow \text{adj } A = \begin{bmatrix} 4 & 3 \\ -1 & 2 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{|A|}(\text{adj } A) = \frac{1}{11} \begin{bmatrix} 4 & 3 \\ -1 & 2 \end{bmatrix}$$

$$\text{Now, } AX = B \Rightarrow X = A^{-1}B \Rightarrow X = \frac{1}{11} \begin{bmatrix} 4 & 3 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} -1 \\ -3 \end{bmatrix}$$

$$\Rightarrow X = \frac{1}{11} \begin{bmatrix} -4-9 \\ 1-6 \end{bmatrix} \Rightarrow X = \frac{1}{11} \begin{bmatrix} -13 \\ -5 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -13/11 \\ -5/11 \end{bmatrix}. \text{ Hence, } x = \frac{-13}{11} \text{ and } y = \frac{-5}{11}$$

14. $4x - 3y = 3; 3x - 5y = 7$

Sol. $A = \begin{bmatrix} 4 & -3 \\ 3 & -5 \end{bmatrix}, X = \begin{bmatrix} x \\ y \end{bmatrix}, B = \begin{bmatrix} 3 \\ 7 \end{bmatrix}, |A| = \begin{vmatrix} 4 & -3 \\ 3 & -5 \end{vmatrix} = (-20 + 9) = (-11)$. Hence, A^{-1} exist.

$$\therefore \text{adj } A = \begin{bmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{bmatrix}'$$

So the cofactor of the element of matrix A are, $\begin{bmatrix} C_{11} = -5 & C_{21} = 3 \\ C_{12} = -3 & C_{22} = 4 \end{bmatrix}$

$$\text{adj } A = \begin{bmatrix} -5 & -3 \\ 3 & 4 \end{bmatrix}' \Rightarrow \text{adj } A = \begin{bmatrix} -5 & 3 \\ -3 & 4 \end{bmatrix}$$

$$A^{-1} = \frac{1}{|A|}(\text{adj } A) = \frac{1}{(-11)} \begin{bmatrix} -5 & 3 \\ -3 & 4 \end{bmatrix} = \frac{1}{11} \begin{bmatrix} 5 & -3 \\ 3 & -4 \end{bmatrix}$$

$$\text{Now, } AX = B \Rightarrow X = A^{-1}B \Rightarrow X = \frac{1}{11} \begin{bmatrix} 5 & -3 \\ 3 & -4 \end{bmatrix} \begin{bmatrix} 3 \\ 7 \end{bmatrix}$$

$$\Rightarrow X = \frac{1}{11} \begin{bmatrix} 15-21 \\ 9-28 \end{bmatrix} \Rightarrow X = \frac{1}{11} \begin{bmatrix} -6 \\ -19 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -6/11 \\ -19/11 \end{bmatrix}. \text{ Hence, } x = \frac{-6}{11} \text{ and } y = \frac{-19}{11}.$$

15. $2x + 8y + 5z = 5; x + y + z = -2; x + 2y - z = 2$

Sol. Let $A = \begin{bmatrix} 2 & 8 & 5 \\ 1 & 1 & 1 \\ 1 & 2 & -1 \end{bmatrix}, X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, B = \begin{bmatrix} 5 \\ -2 \\ 2 \end{bmatrix}$

$$|A| = 2 \begin{vmatrix} 1 & 1 \\ 2 & -1 \end{vmatrix} - 8 \begin{vmatrix} 1 & 1 \\ 1 & -1 \end{vmatrix} + 5 \begin{vmatrix} 1 & 1 \\ 1 & 2 \end{vmatrix} = 2(-3) - 8(-2) + 5(1) = -6 + 16 + 5 = 15 \Rightarrow |A| = 15$$

Hence, A^{-1} exist.

$$\therefore \text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}'$$

<https://millionstar.godaddy.com/>

So the cofactor of the element of matrix A are,

$$\begin{bmatrix} C_{11} = 9 & C_{12} = 2 & C_{13} = 1 \\ C_{21} = 18 & C_{22} = -7 & C_{23} = 4 \\ C_{31} = 3 & C_{32} = 3 & C_{33} = -6 \end{bmatrix}$$

$$\text{adj } A = \begin{bmatrix} -3 & 2 & 1 \\ 18 & -7 & 4 \\ 3 & 3 & -6 \end{bmatrix} \Rightarrow \text{adj } A = \begin{bmatrix} -3 & 18 & 3 \\ 2 & -7 & 3 \\ 1 & 4 & -6 \end{bmatrix} \therefore A^{-1} = \frac{1}{15} \begin{bmatrix} -3 & 18 & 3 \\ 2 & -7 & 3 \\ 1 & 4 & -6 \end{bmatrix}$$

$$\therefore AX = B \Rightarrow X = A^{-1}B \Rightarrow X = \frac{1}{15} \begin{bmatrix} -3 & 18 & 3 \\ 2 & -7 & 3 \\ 1 & 4 & -6 \end{bmatrix} \begin{bmatrix} 5 \\ -2 \\ 2 \end{bmatrix} \Rightarrow X = \frac{1}{15} \begin{bmatrix} -15 - 36 + 6 \\ 10 + 14 + 6 \\ 5 - 8 - 12 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{15} \begin{bmatrix} -45 \\ 30 \\ -15 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} -3 \\ 2 \\ -1 \end{bmatrix} \therefore x = -3, y = 2, z = -1$$

16. $x - y + z = 1; 2x + y - z = 2; x - 2y - z = 4$

Sol. $A = \begin{bmatrix} 1 & -1 & 1 \\ 2 & 1 & -1 \\ 1 & -2 & -1 \end{bmatrix}, X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, B = \begin{bmatrix} 1 \\ 2 \\ 4 \end{bmatrix}$

$$|A| = \begin{vmatrix} 1 & -1 & 1 \\ 2 & 1 & -1 \\ 1 & -2 & -1 \end{vmatrix} \Rightarrow |A| = 1(-1-2) + 1(-2+1) + 1(-4-1)$$

$$\Rightarrow |A| = -3 - 1 - 5 \Rightarrow |A| = (-9). \text{ Hence, } A^{-1} \text{ exist}$$

$$\therefore \text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}$$

So the cofactor of the element of matrix A are,

$$\begin{bmatrix} C_{11} = -3 & C_{12} = 1 & C_{13} = -5 \\ C_{21} = -3 & C_{22} = -2 & C_{23} = 1 \\ C_{31} = 0 & C_{32} = 3 & C_{33} = 3 \end{bmatrix}$$

$$\therefore \text{adj } A = \begin{bmatrix} -3 & 1 & -5 \\ -3 & -2 & 1 \\ 0 & 3 & 3 \end{bmatrix} \Rightarrow \text{adj } A = \begin{bmatrix} -3 & -3 & 0 \\ 1 & -2 & 3 \\ -5 & 1 & 3 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{|A|} (\text{adj } A) = \frac{1}{(-9)} \begin{bmatrix} -3 & -3 & 0 \\ 1 & -2 & 3 \\ -5 & 1 & 3 \end{bmatrix} = \frac{1}{9} \begin{bmatrix} 3 & 3 & 0 \\ -1 & 2 & -3 \\ 5 & -1 & -3 \end{bmatrix}$$

$$\text{Now, } AX = B \Rightarrow X = A^{-1}B \Rightarrow X = \frac{1}{9} \begin{bmatrix} 3 & 3 & 0 \\ -1 & 2 & -3 \\ 5 & -1 & -3 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 4 \end{bmatrix}$$

$$\Rightarrow X = \frac{1}{9} \begin{bmatrix} 3+6+0 \\ -1+4-12 \\ 5-2-12 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{9} \begin{bmatrix} 9 \\ -9 \\ -9 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \\ -1 \end{bmatrix}. \text{ Hence, } x=1, y=-1 \text{ and } z=-1.$$

17. $3x+4y+7z=4; 2x-y+3z=-3; x+2y-3z=8$

Sol. The given system of linear equations may be written in matrix form as $AX = B$ where

$$A = \begin{bmatrix} 3 & 4 & 7 \\ 2 & -1 & 3 \\ 1 & 2 & -3 \end{bmatrix}, X = \begin{bmatrix} x \\ y \\ z \end{bmatrix} \text{ and } B = \begin{bmatrix} 4 \\ -3 \\ 8 \end{bmatrix}$$

$$\text{Now, } |A| = \begin{vmatrix} 3 & 4 & 7 \\ 2 & -1 & 3 \\ 1 & 2 & -3 \end{vmatrix}$$

$$= 3(3-6) - 4(-6-3) + 7(4+1) = -9+36+35 = 62 \neq 0$$

$$C_{11} = (-1)^{1+1} \begin{vmatrix} -1 & 3 \\ 2 & -3 \end{vmatrix} = 3-6 = -3$$

$$C_{12} = (-1)^{1+2} \begin{vmatrix} 2 & 3 \\ 1 & -3 \end{vmatrix} = -\{-6-3\} = 9$$

$$C_{13} = (-1)^{1+3} \begin{vmatrix} 2 & -1 \\ 1 & 2 \end{vmatrix} = 4+1 = 5$$

$$C_{21} = (-1)^{2+1} \begin{vmatrix} 4 & 7 \\ 2 & -3 \end{vmatrix} = -(-12-14) = 26$$

$$C_{22} = (-1)^{2+2} \begin{vmatrix} 3 & 7 \\ 1 & -3 \end{vmatrix} = -8-7 = -16$$

$$C_{23} = (-1)^{2+3} \begin{vmatrix} 3 & 4 \\ 1 & 2 \end{vmatrix} = -(6-4) = -2$$

$$C_{31} = (-1)^{3+1} \begin{vmatrix} 4 & 7 \\ -1 & 3 \end{vmatrix} = 12+7 = 19$$

$$C_{32} = (-1)^{3+2} \begin{vmatrix} 3 & 7 \\ 2 & 3 \end{vmatrix} = -(9-14) = 5$$

$$C_{33} = (-1)^{3+3} \begin{vmatrix} 3 & 4 \\ 2 & -1 \end{vmatrix} = (-3-8) = -11$$

$$\therefore \text{Adj} \cdot A = \begin{bmatrix} -3 & 9 & 5 \\ 26 & -19 & -2 \\ 19 & 5 & -11 \end{bmatrix} = \begin{bmatrix} -3 & 26 & 19 \\ 9 & -16 & 5 \\ 5 & -2 & -11 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{|A|} \text{Adj} A$$

$$= \frac{1}{62} \begin{bmatrix} -3 & 26 & 19 \\ 9 & -16 & 5 \\ 5 & -2 & -11 \end{bmatrix}$$

$$\therefore AX = B$$

$$\Rightarrow X = A^{-1}B$$

<https://millionstar.godaddysites.com/>

$$\Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{62} \begin{bmatrix} -3 & 26 & 19 \\ 9 & -16 & 5 \\ 5 & -2 & -11 \end{bmatrix} \begin{bmatrix} 4 \\ -3 \\ 8 \end{bmatrix} = \frac{1}{62} \begin{bmatrix} -12 - 78 + 152 \\ 36 + 48 + 40 \\ 20 + 6 - 88 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{62} \begin{bmatrix} 62 \\ 124 \\ -62 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix}$$

Equation the corresponding element we get $x=1, y=2, z=-1$

18. $x+2y+z=7; x+3z=11; 2x-3y=1$

Sol. $A = \begin{bmatrix} 1 & 2 & 1 \\ 1 & 0 & 3 \\ 2 & -3 & 0 \end{bmatrix}, X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, B = \begin{bmatrix} 7 \\ 11 \\ 1 \end{bmatrix}$

$$|A| = \begin{vmatrix} 1 & 2 & 1 \\ 1 & 0 & 3 \\ 2 & -3 & 1 \end{vmatrix} \Rightarrow |A| = 1(0+9) - 2(0-6) + 1(-3-0) \Rightarrow |A| = 9+12-3 \Rightarrow |A| = 18$$

Hence, A^{-1} exist.

$$\therefore \text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}'$$

So the cofactor of the element of matrix A are, $\begin{bmatrix} C_{11}=9 & C_{12}=6 & C_{13}=-3 \\ C_{21}=-3 & C_{22}=-2 & C_{23}=7 \\ C_{31}=6 & C_{32}=-2 & C_{33}=-2 \end{bmatrix}$

$$\text{adj } A = \begin{bmatrix} 9 & 6 & -3 \\ -3 & -2 & 7 \\ 6 & -2 & -2 \end{bmatrix}' \Rightarrow \text{adj } A = \begin{bmatrix} 9 & -3 & 6 \\ 6 & -2 & -2 \\ -3 & 7 & -2 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{|A|} (\text{adj } A) = \frac{1}{18} \begin{bmatrix} 9 & -3 & 6 \\ 6 & -2 & -2 \\ -3 & 7 & -2 \end{bmatrix}$$

Now, $AX = B, X = A^{-1}B \Rightarrow X = \frac{1}{18} \begin{bmatrix} 9 & -3 & 6 \\ 6 & -2 & -2 \\ -3 & 7 & -2 \end{bmatrix} \begin{bmatrix} 7 \\ 11 \\ 1 \end{bmatrix} \Rightarrow X = \frac{1}{18} \begin{bmatrix} 63 - 33 + 6 \\ 42 - 22 - 2 \\ -21 + 77 - 2 \end{bmatrix}$

$$\Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{18} \begin{bmatrix} 36 \\ 18 \\ 54 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix}. \text{ Hence, } x=2, y=1 \text{ and } z=3.$$

19. $2x-3y+5z=16; 3x+2y-4z=-4; x+y-2z=-3$

Sol. Let $A = \begin{bmatrix} 2 & -3 & 5 \\ 3 & 2 & -4 \\ 1 & 1 & -2 \end{bmatrix}, X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, B = \begin{bmatrix} 16 \\ -4 \\ -3 \end{bmatrix}$

<https://millionstar.godaddysites.com/>

$$|A| = \begin{vmatrix} 2 & -3 & 5 \\ 3 & 2 & -4 \\ 1 & 1 & -2 \end{vmatrix} = 2(-4+4) + 3(-6+4) + 5(3-2) = 2(0) + 3(-2) + 5(1) = -6 + 5 = -1$$

$\Rightarrow |A| = -1$. Hence, A^{-1} exist.

$$\therefore \text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}'$$

So the cofactor of the element of matrix A are, $\begin{bmatrix} C_{11}=0 & C_{12}=2 & C_{13}=1 \\ C_{21}=-1 & C_{22}=-9 & C_{23}=-5 \\ C_{31}=2 & C_{32}=23 & C_{33}=13 \end{bmatrix}$

$$\text{adj } A = \begin{bmatrix} 0 & 2 & 1 \\ -1 & -9 & -5 \\ 2 & 23 & 13 \end{bmatrix}' \Rightarrow \text{adj } A = \begin{bmatrix} 0 & -1 & 2 \\ 2 & -9 & 23 \\ 1 & -5 & 13 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{|A|} (\text{adj } A) = -1 \begin{bmatrix} 0 & -1 & 2 \\ 2 & -9 & 23 \\ 1 & -5 & 13 \end{bmatrix} \Rightarrow A^{-1} = \begin{bmatrix} 0 & 1 & -2 \\ -2 & 9 & -23 \\ -1 & 5 & -13 \end{bmatrix}$$

$$\therefore AX = B, X = A^{-1}B \Rightarrow X = \begin{bmatrix} 0 & 1 & -2 \\ -2 & 9 & -23 \\ -1 & 5 & -13 \end{bmatrix} \begin{bmatrix} 16 \\ -4 \\ -3 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0-4+6 \\ -32-36+69 \\ -16-20+39 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix} \text{ Hence } x=2, y=1, z=3$$

20. $x+y+z=4; 2x-y+z=-1; 2x+y-3z=-9$

Sol. Let $A = \begin{bmatrix} 1 & 1 & 1 \\ 2 & -1 & 1 \\ 2 & 1 & -3 \end{bmatrix}$, $X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$, $B = \begin{bmatrix} 4 \\ -1 \\ -9 \end{bmatrix}$

$$|A| = \begin{vmatrix} 1 & 1 & 1 \\ 2 & -1 & 1 \\ 2 & 1 & -3 \end{vmatrix} \Rightarrow |A| = (3-1) - (-6-2) + (2+2) \Rightarrow |A| = 2+8+4 \Rightarrow |A| = 14$$

Hence, A^{-1} exist.

$$\therefore \text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}'$$

So the cofactor of the element of matrix A are, $\begin{bmatrix} C_{11}=2 & C_{12}=8 & C_{13}=4 \\ C_{21}=4 & C_{22}=-5 & C_{23}=1 \\ C_{31}=2 & C_{32}=1 & C_{33}=-3 \end{bmatrix}$

<https://millionstar.godaddysites.com/>

$$\therefore \text{adj } A = \begin{bmatrix} 2 & 8 & 4 \\ 4 & -5 & 1 \\ 2 & 1 & -3 \end{bmatrix} = \begin{bmatrix} 2 & 4 & 2 \\ 8 & -5 & 1 \\ 4 & 1 & -3 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{|A|}(\text{adj } A) \Rightarrow A^{-1} = \frac{1}{14} \begin{bmatrix} 2 & 4 & 2 \\ 8 & -5 & 1 \\ 4 & 1 & -3 \end{bmatrix}$$

$$AX = B \Rightarrow X = A^{-1}B$$

$$\Rightarrow X = \frac{1}{14} \begin{bmatrix} 2 & 4 & 2 \\ 8 & -5 & 1 \\ 4 & 1 & -3 \end{bmatrix} \begin{bmatrix} 4 \\ -1 \\ -9 \end{bmatrix} \Rightarrow X = \frac{1}{14} \begin{bmatrix} 8-4-18 \\ 32+5-9 \\ 16-1+27 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{14} \begin{bmatrix} -14 \\ 28 \\ 42 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} -1 \\ 2 \\ 3 \end{bmatrix} \text{ Hence } x = -1, y = 2, z = 3$$

21. $2x - 3y + 5z = 11$; $3x + 2y - 4z = -5$; $x + y - 2z = -3$

Sol. $A = \begin{bmatrix} 2 & -3 & 5 \\ 3 & 2 & -4 \\ 1 & 1 & -2 \end{bmatrix}$, $X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$, $B = \begin{bmatrix} 11 \\ -5 \\ -3 \end{bmatrix}$

$$|A| = \begin{vmatrix} 2 & -3 & 5 \\ 3 & 2 & -4 \\ 1 & 1 & -2 \end{vmatrix} \Rightarrow |A| = 2(-4+4) + 3(-6+4) + 5(3-2) \Rightarrow |A| = 0 - 6 + 5 \Rightarrow |A| = (-1)$$

Hence, A^{-1} exist.

$$\therefore \text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}'$$

So the cofactor of the element of matrix A are, $\begin{bmatrix} C_{11} = 0 & C_{12} = 2 & C_{13} = 1 \\ C_{21} = -1 & C_{22} = -9 & C_{23} = -5 \\ C_{31} = 2 & C_{32} = 23 & C_{33} = 13 \end{bmatrix}$

$$\text{adj } A = \begin{bmatrix} 0 & 2 & 1 \\ -1 & -9 & -5 \\ 2 & 23 & 13 \end{bmatrix}' \Rightarrow \text{adj } A = \begin{bmatrix} 0 & -1 & 2 \\ 2 & -9 & 23 \\ 1 & -5 & 13 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{|A|}(\text{adj } A) = \frac{1}{(-1)} \begin{bmatrix} 0 & -1 & 2 \\ 2 & -9 & 23 \\ 1 & -5 & 13 \end{bmatrix} = \begin{bmatrix} 0 & 1 & -2 \\ -2 & 9 & -23 \\ -1 & 5 & -13 \end{bmatrix}$$

Now, $AX = B$, $X = A^{-1}B$

$$\Rightarrow X = \begin{bmatrix} 0 & 1 & -2 \\ -2 & 9 & -23 \\ -1 & 5 & -13 \end{bmatrix} \begin{bmatrix} 11 \\ -5 \\ -3 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0-5+6 \\ -22-45+69 \\ -11-25+39 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$$

Hence, $x=1, y=2$ and <https://millionstar.godaddysites.com/>

$$x+y+z=1; x-2y+3z=2; 5x-3y+z=3$$

Sol. $A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & -2 & 3 \\ 5 & -3 & 1 \end{bmatrix}, X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, B = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$

$$|A| = \begin{vmatrix} 1 & 1 & 1 \\ 1 & -2 & 3 \\ 5 & -3 & 1 \end{vmatrix} \Rightarrow |A| = 1(-2+9) - 1(1-15) + 1(-3+10) \Rightarrow |A| = 7+14+7 \Rightarrow |A| = 28$$

Hence, A^{-1} exist.

$$\therefore \text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}$$

So the cofactor of the element of matrix A are, $\begin{bmatrix} C_{11}=7 & C_{12}=14 & C_{13}=7 \\ C_{21}=-4 & C_{22}=-4 & C_{23}=8 \\ C_{31}=5 & C_{32}=-2 & C_{33}=-3 \end{bmatrix}$

$$\therefore \text{adj } A = \begin{bmatrix} 7 & 14 & 7 \\ -4 & -4 & 8 \\ 5 & -2 & -3 \end{bmatrix} \Rightarrow \text{adj } A = \begin{bmatrix} 7 & -4 & 5 \\ 14 & -4 & -2 \\ 7 & 8 & -3 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{|A|} (\text{adj } A) = \frac{1}{28} \begin{bmatrix} 7 & -4 & 5 \\ 14 & -4 & -2 \\ 7 & 8 & -3 \end{bmatrix}$$

Now, $AX = B, X = A^{-1}B$

$$\Rightarrow X = \frac{1}{28} \begin{bmatrix} 7 & -4 & 5 \\ 14 & -4 & -2 \\ 7 & 8 & -3 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \Rightarrow X = \frac{1}{28} \begin{bmatrix} 7-8+15 \\ 14-8-6 \\ 7+16-9 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{28} \begin{bmatrix} 14 \\ 0 \\ 14 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1/2 \\ 0 \\ 1/2 \end{bmatrix}$$

Hence, $x = \frac{1}{2}, y = 0$ and $z = \frac{1}{2}$

23. $x+y+z=6; x+2z=7; 3x+y+z=12$

Sol. $A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 0 & 2 \\ 3 & 1 & 1 \end{bmatrix}, X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, B = \begin{bmatrix} 6 \\ 7 \\ 12 \end{bmatrix}$

$$|A| = \begin{vmatrix} 1 & 1 & 1 \\ 1 & 0 & 2 \\ 3 & 1 & 1 \end{vmatrix} \Rightarrow |A| = 1(0-2) - 1(1-6) + 1(1-0) \Rightarrow |A| = -2+5+1 \Rightarrow |A| = 4$$

Hence, A^{-1} exist.

<https://millionstar.godaddysites.com/>

$$\therefore \text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}$$

So the cofactor of the element of matrix A are, $\begin{bmatrix} C_{11} = -2 & C_{12} = 5 & C_{13} = 1 \\ C_{21} = 0 & C_{22} = -2 & C_{23} = 2 \\ C_{31} = 2 & C_{32} = -1 & C_{33} = -1 \end{bmatrix}$

$$\text{adj } A = \begin{bmatrix} -2 & 5 & 1 \\ 0 & -2 & 2 \\ 2 & -1 & -1 \end{bmatrix}' \Rightarrow \text{adj } A = \begin{bmatrix} -2 & 0 & 2 \\ 5 & -2 & -1 \\ 1 & 2 & -1 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{|A|} (\text{adj } A) = \frac{1}{4} \begin{bmatrix} -2 & 0 & 2 \\ 5 & -2 & -1 \\ 1 & 2 & -1 \end{bmatrix}$$

Now, $AX = B$, $X = A^{-1}B$

$$\Rightarrow X = \frac{1}{4} \begin{bmatrix} -2 & 0 & 2 \\ 5 & -2 & -1 \\ 1 & 2 & -1 \end{bmatrix} \begin{bmatrix} 6 \\ 7 \\ 12 \end{bmatrix} \Rightarrow X = \frac{1}{4} \begin{bmatrix} -12+0+24 \\ 30-14-12 \\ 6+14-12 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{4} \begin{bmatrix} 12 \\ 4 \\ 8 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 3 \\ 1 \\ 2 \end{bmatrix}$$

Hence, $x = 3$, $y = 1$ and $z = 2$

24. $2x + 3y + 3z = 5$; $x - 2y + z = -4$; $3x - y - 2z = 3$

Sol. $A = \begin{bmatrix} 2 & 3 & 3 \\ 1 & -2 & 1 \\ 3 & -1 & -2 \end{bmatrix}$, $X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$, $B = \begin{bmatrix} 5 \\ -4 \\ 3 \end{bmatrix}$

$$|A| = \begin{vmatrix} 2 & 3 & 3 \\ 1 & -2 & 1 \\ 3 & -1 & -2 \end{vmatrix} \Rightarrow |A| = 2(4+1) - 3(-2-3) + 3(-1+6) \Rightarrow |A| = 10+15+15 \Rightarrow |A| = 40$$

Hence, A^{-1} exist

$$\therefore \text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}'$$

So the cofactor of the element of matrix A are, $\begin{bmatrix} C_{11} = 5 & C_{12} = 5 & C_{13} = 5 \\ C_{21} = 3 & C_{22} = -13 & C_{23} = 11 \\ C_{31} = 9 & C_{32} = 1 & C_{33} = -7 \end{bmatrix}$

$$\text{adj } A = \begin{bmatrix} 5 & 5 & 5 \\ 3 & -13 & 11 \\ 9 & 1 & -7 \end{bmatrix}' \Rightarrow \text{adj } A = \begin{bmatrix} 5 & 3 & 9 \\ 5 & -13 & 1 \\ 5 & 11 & -7 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{|A|} (\text{adj } A) = \frac{1}{40} \begin{bmatrix} 5 & 3 & 9 \\ 5 & -13 & 1 \\ 5 & 11 & -7 \end{bmatrix}$$

<https://millionstarsgodaaddysites.com/>

Now, $AX = B, X = A^{-1}B \Rightarrow X = \frac{1}{40} \begin{bmatrix} 5 & -13 & 1 \\ 5 & 11 & -7 \end{bmatrix} \begin{bmatrix} -4 \\ 3 \end{bmatrix}$

$$\Rightarrow X = \frac{1}{40} \begin{bmatrix} 25-12+27 \\ 25+52+3 \\ 25-44-21 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{40} \begin{bmatrix} 40 \\ 80 \\ -40 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix}$$

Hence, $x=1, y=2$ and $z=-1$

25. $4x-5y-11z=12; x-3y+z=1; 2x+3y-7z=2$

Sol. $A = \begin{bmatrix} 4 & -5 & -11 \\ 1 & -3 & 1 \\ 2 & 3 & -7 \end{bmatrix}, X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, B = \begin{bmatrix} 12 \\ 1 \\ 2 \end{bmatrix}$

$$|A| = \begin{vmatrix} 4 & -5 & -11 \\ 1 & -3 & 1 \\ 2 & 3 & -7 \end{vmatrix} \Rightarrow |A| = 4(21-3) + 5(-7-2) - 11(3+6) \Rightarrow |A| = 72 - 45 - 99$$

$\Rightarrow |A| = (-72)$. Hence, A^{-1} exist.

$$\therefore \text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}'$$

So the cofactor of the element of matrix A are, $\begin{bmatrix} C_{11}=18 & C_{12}=9 & C_{13}=9 \\ C_{21}=-68 & C_{22}=-6 & C_{23}=-22 \\ C_{31}=-38 & C_{32}=-15 & C_{33}=-7 \end{bmatrix}$

$$\text{adj } A = \begin{bmatrix} 18 & 9 & 9 \\ -68 & -6 & -22 \\ -38 & -15 & -7 \end{bmatrix}' \Rightarrow \text{adj } A = \begin{bmatrix} 18 & -68 & -38 \\ 9 & -6 & -15 \\ 9 & -22 & -7 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{|A|} (\text{adj } A) = \frac{1}{(-72)} \begin{bmatrix} 18 & -68 & -38 \\ 9 & -6 & -15 \\ 9 & -22 & -7 \end{bmatrix} = \frac{1}{72} \begin{bmatrix} -18 & 68 & 38 \\ -9 & 6 & 15 \\ -9 & 22 & 7 \end{bmatrix}$$

Now, $AX = B, X = A^{-1}B \Rightarrow X = \frac{1}{72} \begin{bmatrix} -18 & 68 & 38 \\ -9 & 6 & 15 \\ -9 & 22 & 7 \end{bmatrix} \begin{bmatrix} 12 \\ 1 \\ 2 \end{bmatrix}$

$$\Rightarrow X = \frac{1}{72} \begin{bmatrix} -216+68+76 \\ -108+6+30 \\ -108+22+14 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{72} \begin{bmatrix} -72 \\ -72 \\ -72 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} -1 \\ -1 \\ -1 \end{bmatrix}$$

Hence, $x=-1, y=-1$ and $z=-1$

26. $x-y+2z=7; 3z+4y-5z=-5; 2x-y+3z=12$

Sol. $A = \begin{bmatrix} 1 & -1 & 2 \\ 3 & 4 & -5 \\ 2 & -1 & 3 \end{bmatrix}, X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, B = \begin{bmatrix} 7 \\ -5 \\ 12 \end{bmatrix}$

Now, $|A| = \begin{vmatrix} 1 & -1 & 2 \\ 3 & 4 & -5 \\ 2 & -1 & 3 \end{vmatrix} = 1(12-5) + 1(9+10) + 2(-3-8) = 7 + 19 - 22 = 4 \neq 0$

Hence A^{-1} exist and system have unique solution

$$C_{11} = (-1)^{1+1} \begin{vmatrix} 4 & -5 \\ -1 & 3 \end{vmatrix} = 12 - 5 = 7$$

$$C_{12} = (-1)^{1+2} \begin{vmatrix} 3 & -5 \\ 2 & 3 \end{vmatrix} = -(9+10) = -19$$

$$C_{13} = (-1)^{1+3} \begin{vmatrix} 3 & 4 \\ 2 & -1 \end{vmatrix} = +(-3-8) = -11$$

$$C_{21} = (-1)^{2+1} \begin{vmatrix} -1 & 2 \\ -1 & 3 \end{vmatrix} = -(3+2) = -5$$

$$C_{22} = (-1)^{2+2} \begin{vmatrix} 1 & 2 \\ 2 & 3 \end{vmatrix} = +(3-4) = -1$$

$$C_{23} = (-1)^{2+3} \begin{vmatrix} 1 & -1 \\ 2 & -1 \end{vmatrix} = -(-1+2) = -1$$

$$C_{31} = (-1)^{3+1} \begin{vmatrix} -1 & 2 \\ 4 & -5 \end{vmatrix} = +(5-8) = -3$$

$$C_{32} = (-1)^{3+2} \begin{vmatrix} 1 & 2 \\ 3 & -5 \end{vmatrix} = -(-5-6) = 11$$

$$C_{33} = (-1)^{3+3} \begin{vmatrix} 1 & -1 \\ 3 & 4 \end{vmatrix} = +(4+3) = 7$$

$$\therefore \text{adj}A = \begin{bmatrix} 7 & -19 & -11 \\ 1 & -1 & -1 \\ -3 & 11 & 7 \end{bmatrix}^T = \begin{bmatrix} 7 & 1 & -3 \\ -19 & -1 & 11 \\ -11 & -1 & 7 \end{bmatrix}$$

$$\Rightarrow A^{-1} = \frac{1}{|A|} \text{adj}A = \frac{1}{4} \begin{bmatrix} 7 & 1 & -3 \\ -19 & -1 & 11 \\ -11 & -1 & 7 \end{bmatrix}$$

$$\therefore AX = B$$

$$\Rightarrow X = A^{-1}B$$

$$\Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{4} \begin{bmatrix} 7 & 1 & -3 \\ -19 & -1 & 11 \\ -11 & -1 & 7 \end{bmatrix} \begin{bmatrix} 7 \\ -5 \\ 12 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{4} \begin{bmatrix} 49-5-36 \\ -133+5+132 \\ -77+5+84 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{4} \begin{bmatrix} 8 \\ 4 \\ 12 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix}$$

Equation the corresponding elements, we get $x = 3$

$$6x - 9y - 20z = -4; 4x - 15y + 10z = -1; 2x - 3y - 5z = -1$$

Sol. Let $A = \begin{bmatrix} 6 & -9 & -20 \\ 4 & -15 & 10 \\ 2 & -3 & -5 \end{bmatrix}$, $X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$, $B = \begin{bmatrix} -4 \\ -1 \\ -1 \end{bmatrix}$

$$|A| = \begin{vmatrix} 6 & -9 & -20 \\ 4 & -15 & 10 \\ 2 & -3 & -5 \end{vmatrix} \Rightarrow |A| = 6(75 + 30) + 9(-20 - 20) - 20(-12 + 30) = 6(105) + 9(-40) - 20(18)$$

$$\Rightarrow |A| = 630 - 360 - 360 = -90. \text{ Hence, } A^{-1} \text{ exist.}$$

$$\therefore \text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}'$$

So the cofactor of the element of matrix A are, $\begin{bmatrix} C_{11} = 105 & C_{12} = 40 & C_{13} = 18 \\ C_{21} = 15 & C_{22} = 10 & C_{23} = 0 \\ C_{31} = -390 & C_{32} = -140 & C_{33} = -54 \end{bmatrix}$

$$\text{adj } A = \begin{bmatrix} 105 & 40 & 18 \\ 15 & 10 & 0 \\ -390 & -140 & -54 \end{bmatrix}' \Rightarrow (\text{adj } A) = \begin{bmatrix} 105 & 15 & -390 \\ 40 & 10 & -140 \\ 18 & 0 & -54 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{|A|} (\text{adj } A) = \frac{1}{-90} \begin{bmatrix} 105 & 15 & -390 \\ 40 & 10 & -140 \\ 18 & 0 & -54 \end{bmatrix}$$

Now, $AX = B$, $X = A^{-1}B$

$$\Rightarrow X = -\frac{1}{90} \begin{bmatrix} 105 & 15 & -390 \\ 40 & 10 & -140 \\ 18 & 0 & -54 \end{bmatrix} \begin{bmatrix} -4 \\ -1 \\ -1 \end{bmatrix} \Rightarrow X = -\frac{1}{90} \begin{bmatrix} -420 - 15 + 390 \\ -160 - 10 + 140 \\ -72 + 0 + 54 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = -\frac{1}{90} \begin{bmatrix} -45 \\ -30 \\ -18 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1/2 \\ 1/3 \\ 1/5 \end{bmatrix} \text{ Hence } x = \frac{1}{2}, y = \frac{1}{3}, z = \frac{1}{5}$$

28. $3x - 4y + 2z = -1$; $2x + 3y + 5z = 7$; $x + z = 2$

Sol. $A = \begin{bmatrix} 3 & -4 & 2 \\ 2 & 3 & 5 \\ 1 & 0 & 1 \end{bmatrix}$, $X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$, $B = \begin{bmatrix} -1 \\ 7 \\ 2 \end{bmatrix}$

$$|A| = \begin{vmatrix} 3 & -4 & 2 \\ 2 & 3 & 5 \\ 1 & 0 & 1 \end{vmatrix} \Rightarrow |A| = 3(3 - 0) + 4(2 - 5) + 2(0 - 3) \Rightarrow |A| = 9 - 12 - 6 \Rightarrow |A| = (-9)$$

Hence, A^{-1} exist.

$$\therefore \text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}$$

So the cofactor of the element of matrix A are, $\begin{bmatrix} C_{11}=3 & C_{12}=3 & C_{13}=-3 \\ C_{21}=4 & C_{22}=1 & C_{23}=-4 \\ C_{31}=-26 & C_{32}=-11 & C_{33}=17 \end{bmatrix}$

$$\text{adj } A = \begin{bmatrix} 3 & 3 & -3 \\ 4 & 1 & -4 \\ -26 & -11 & 17 \end{bmatrix}' \Rightarrow \text{adj } A = \begin{bmatrix} 3 & 4 & -26 \\ 3 & 1 & -11 \\ -3 & -4 & 17 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{|A|} (\text{adj } A) = \frac{1}{(-9)} \begin{bmatrix} 3 & 4 & -26 \\ 3 & 1 & -11 \\ -3 & -4 & 17 \end{bmatrix} = \frac{1}{9} \begin{bmatrix} -3 & -4 & 26 \\ -3 & -1 & 11 \\ 3 & 4 & -17 \end{bmatrix}$$

Now, $AX = B \Rightarrow X = A^{-1}B$

$$\text{Now, } X = A^{-1}B = \frac{1}{9} \begin{bmatrix} -3 & 2 & -1 \\ 1 & 0 & -1 \\ -4 & 2 & -2 \end{bmatrix} \begin{bmatrix} 1 \\ 3 \\ -1 \end{bmatrix} \Rightarrow X = \frac{1}{9} \begin{bmatrix} -3+6+1 \\ 1+0+1 \\ -4+6+2 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{9} \begin{bmatrix} 4 \\ 2 \\ 4 \end{bmatrix} \quad \text{Hence, } x = \frac{4}{9}, y = \frac{2}{9}, z = \frac{4}{9}$$

29. $x + y - z = 1; 3x + y - 2z = 3; x - y - z = -1.$

Sol. $A = \begin{bmatrix} 1 & 1 & -1 \\ 3 & 1 & -2 \\ 1 & -1 & -1 \end{bmatrix}$ $X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$ $B = \begin{bmatrix} 1 \\ 3 \\ -1 \end{bmatrix}$ and $AX = B \Rightarrow X = A^{-1}B$

$$|A| = \begin{vmatrix} 1 & 1 & -1 \\ 3 & 1 & -2 \\ 1 & -1 & -1 \end{vmatrix} = 1 \begin{vmatrix} 1 & -2 \\ -1 & -1 \end{vmatrix} - 1 \begin{vmatrix} 3 & -2 \\ 1 & -1 \end{vmatrix} - 1 \begin{vmatrix} 3 & 1 \\ 1 & -1 \end{vmatrix}$$

$$= (-1-2) - (-3+2) - (-3-1) = -3+1+4 = 2. \therefore |A| \neq 0. \text{ Hence, } A^{-1} \text{ exist}$$

$$\therefore \text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}'$$

So the cofactor of the element of matrix A are,

$$C_{11} = (-1-2) = -3, \quad C_{21} = -(-1-1) = 2, \quad C_{31} = (-2+1) = -1$$

$$C_{12} = -(-3+2) = 1, \quad C_{22} = (-1+1) = 0, \quad C_{32} = -(-2+3) = -1$$

$$C_{13} = (-3-1) = -4, \quad C_{23} = -(-1-1) = 2, \quad C_{33} = (1-3) = -2$$

$$\therefore \text{adj } A = \begin{bmatrix} -3 & 1 & -4 \\ 2 & 0 & 2 \\ -1 & -1 & -2 \end{bmatrix}' = \begin{bmatrix} -3 & 2 & -1 \\ 1 & 0 & -1 \\ -4 & 2 & -2 \end{bmatrix}, \quad A^{-1} = \frac{\text{adj } A}{|A|} = \frac{1}{2} \begin{bmatrix} -3 & 2 & -1 \\ 1 & 0 & -1 \\ -4 & 2 & -2 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} -3 & 2 & -1 \\ 1 & 0 & -1 \\ -4 & 2 & -2 \end{bmatrix}$$

$$X = A^{-1}B = \frac{1}{2} \begin{bmatrix} -3 & 2 & 1 \\ 1 & 0 & -1 \\ -4 & 2 & -2 \end{bmatrix} \begin{bmatrix} 1 \\ 3 \\ -1 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 1+0+1 \\ 1+0+1 \\ -4+6+2 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 2 \\ 2 \\ 4 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}. \text{ So, } x=1, y=1, z=2$$

30. $2x + y - z = 1; x - y + z = 2; 3x + y - 2z = -1$

Sol. $A = \begin{bmatrix} 2 & 1 & -1 \\ 1 & -1 & 1 \\ 3 & 1 & -2 \end{bmatrix}, X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, B = \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix}$

$$|A| = \begin{vmatrix} 2 & 1 & -1 \\ 1 & -1 & 1 \\ 3 & 1 & -2 \end{vmatrix} \Rightarrow |A| = 2(2-1) - 1(-2-3) - 1(1+3) \Rightarrow |A| = 2+5-4 \Rightarrow |A| = 3$$

Hence, A^{-1} exist.

$$\therefore \text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}$$

So the cofactor of the element of matrix A are, $\begin{bmatrix} C_{11}=1 & C_{12}=5 & C_{13}=4 \\ C_{21}=1 & C_{22}=-1 & C_{23}=1 \\ C_{31}=0 & C_{32}=-3 & C_{33}=-3 \end{bmatrix}$

$$\text{adj } A = \begin{bmatrix} 1 & 5 & 4 \\ 1 & -1 & 1 \\ 0 & -3 & -3 \end{bmatrix} \Rightarrow \text{adj } A = \begin{bmatrix} 1 & 1 & 0 \\ 5 & -1 & -3 \\ 4 & 1 & -3 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{|A|} (\text{adj } A) = \frac{1}{3} \begin{bmatrix} 1 & 1 & 0 \\ 5 & -1 & -3 \\ 4 & 1 & -3 \end{bmatrix}. \text{ Now } AX = B, X = A^{-1}B$$

$$\Rightarrow X = \frac{1}{3} \begin{bmatrix} 1 & 1 & 0 \\ 5 & -1 & -3 \\ 4 & 1 & -3 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix} \Rightarrow X = \frac{1}{3} \begin{bmatrix} 1+2-0 \\ 5-2-3 \\ 4+2-3 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{3} \begin{bmatrix} 3 \\ 0 \\ 3 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}$$

Hence, $x=1, y=0$ and $z=1$

31. $x + 2y + z = 4; -x + y + z = 0; x - 3y + z = 4$

Sol. $x + 2y + z = 4$ (i)

$-x + y + z = 0$ (ii)

$x - 3y + z = 4$ (iii)

Let $A = \begin{bmatrix} 1 & 2 & 1 \\ -1 & 1 & 1 \\ 1 & -3 & 1 \end{bmatrix}$, $X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$ and $B = \begin{bmatrix} 5 \\ 0 \\ 4 \end{bmatrix}$

Here $AX = B$,

$$\Rightarrow A^{-1}AX = A^{-1}B$$

$$\Rightarrow IX = A^{-1}B$$

$$\Rightarrow X = A^{-1}B \quad \dots\dots (iv)$$

$$\begin{aligned} \text{Now } |A| &= \begin{vmatrix} 1 & 2 & 1 \\ -1 & 1 & 1 \\ 1 & -3 & 1 \end{vmatrix} = \begin{vmatrix} 0 & 2 & 1 \\ -2 & 1 & 1 \\ 0 & -3 & 1 \end{vmatrix} \{C_1 \rightarrow C_1 - C_3\} \\ &= -(-2) \begin{vmatrix} 2 & 1 \\ -3 & 1 \end{vmatrix} = 2(2+3) = 10 \end{aligned}$$

Since $|A| \neq 0 \Rightarrow A^{-1}$ exists

Now, co-factors of A,

$$A_{11} = \begin{vmatrix} 1 & 1 \\ -3 & 1 \end{vmatrix} = 1+3 = 4$$

$$A_{12} = -\begin{vmatrix} -1 & 1 \\ 1 & 1 \end{vmatrix} = -(-1-1) = 2$$

$$A_{13} = \begin{vmatrix} -1 & 1 \\ 1 & -3 \end{vmatrix} = 3-1 = 2$$

$$A_{21} = -\begin{vmatrix} 2 & 1 \\ -3 & 1 \end{vmatrix} = -(2+3) = -5$$

$$A_{22} = \begin{vmatrix} 1 & 1 \\ 1 & 1 \end{vmatrix} = (1-1) = 0$$

$$A_{23} = -\begin{vmatrix} 1 & 2 \\ 1 & -3 \end{vmatrix} = -(-3-2) = 5$$

$$A_{31} = \begin{vmatrix} 2 & 1 \\ 1 & 1 \end{vmatrix} = 2-1 = 1$$

$$A_{32} = -\begin{vmatrix} 1 & 1 \\ -1 & 1 \end{vmatrix} = -(1+1) = -2$$

$$A_{33} = \begin{vmatrix} 1 & 2 \\ -1 & 1 \end{vmatrix} = (1+2) = 3$$

$$\text{Now } \text{adj}(A) = \begin{bmatrix} A_{11} & A_{21} & A_{31} \\ A_{12} & A_{22} & A_{32} \\ A_{13} & A_{23} & A_{33} \end{bmatrix} = \begin{bmatrix} 4 & -5 & 1 \\ 2 & 0 & -2 \\ 2 & 5 & 3 \end{bmatrix}$$

$$A^{-1} = \frac{1}{|A|} \text{Adj}(A) = \frac{1}{10} \begin{bmatrix} 4 & -5 & 1 \\ 2 & 0 & -2 \\ 2 & 5 & 3 \end{bmatrix}$$

Now from (iv),

$$X = A^{-1}B$$

$$X = \frac{1}{10} \begin{bmatrix} 4 & -5 & 1 \\ 2 & 0 & -2 \\ 2 & 5 & 3 \end{bmatrix} \begin{bmatrix} 4 \\ 0 \\ 4 \end{bmatrix} = \frac{1}{10} \begin{bmatrix} 16+0+4 \\ 8+0-8 \\ 8+0+12 \end{bmatrix} = \frac{1}{10} \begin{bmatrix} 20 \\ 0 \\ 20 \end{bmatrix} = \begin{bmatrix} 2 \\ 0 \\ 2 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 2 \\ 0 \\ 2 \end{bmatrix}$$

Hence $x = 2, y = 0, z = 2$

32. $x - y - 2z = 3; x + y = 1; x + z = -6$

Sol. $A = \begin{bmatrix} 1 & -1 & -2 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix}, x = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, B = \begin{bmatrix} 3 \\ 1 \\ -6 \end{bmatrix}$

$$|A| = \begin{vmatrix} 1 & -1 & -2 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \end{vmatrix} \Rightarrow |A| = 1(1-0) + 1(1-0) - 2(0-1) \Rightarrow |A| = 1+1+2 \Rightarrow |A| = 4$$

Hence, A^{-1} exist.

$$\therefore \text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}$$

So the cofactor of the element of matrix A are, $\begin{bmatrix} C_{11}=1 & C_{12}=-1 & C_{13}=-1 \\ C_{21}=1 & C_{22}=3 & C_{23}=-1 \\ C_{31}=2 & C_{32}=-2 & C_{33}=2 \end{bmatrix}$

$$\therefore \text{adj } A = \begin{bmatrix} 1 & -1 & -1 \\ 1 & 3 & -1 \\ 2 & -2 & 2 \end{bmatrix} \Rightarrow \text{adj } A = \begin{bmatrix} 1 & 1 & 2 \\ -1 & 3 & -2 \\ -1 & -1 & 2 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{|A|} (\text{adj } A) = \frac{1}{4} \begin{bmatrix} 1 & 1 & 2 \\ -1 & 3 & -2 \\ -1 & -1 & 2 \end{bmatrix}$$

Now, $AX = B, X = A^{-1}B \Rightarrow X = \frac{1}{4} \begin{bmatrix} 1 & 1 & 2 \\ -1 & 3 & -2 \\ -1 & -1 & 2 \end{bmatrix} \begin{bmatrix} 3 \\ 1 \\ -6 \end{bmatrix}$

$$\Rightarrow X = \frac{1}{4} \begin{bmatrix} 3+1-12 \\ -3+3+12 \\ -3-1-12 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{4} \begin{bmatrix} -8 \\ 12 \\ -16 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} -2 \\ 3 \\ -4 \end{bmatrix} \text{ Hence, } x = -2, y = 3 \text{ and } z = -4$$



Show that each one of the following systems of equations is inconsistent.

$$5x - y = -7; 2x + 3z = 1; 3y - z = 5.$$

Sol. $A = \begin{bmatrix} 5 & -1 & 0 \\ 2 & 0 & 3 \\ 0 & 3 & -1 \end{bmatrix}, X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, B = \begin{bmatrix} -7 \\ 1 \\ 5 \end{bmatrix}$

$$|A| = \begin{vmatrix} 5 & -1 & 0 \\ 2 & 0 & 3 \\ 0 & 3 & -1 \end{vmatrix} \Rightarrow |A| = 5(0-9) + 1(-2-0) + 0 \Rightarrow |A| = -45 - 2 \Rightarrow |A| = (-47)$$

Hence, A^{-1} exist.

$$\therefore \text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}'$$

So the cofactor of the element of matrix A are, $\begin{bmatrix} C_{11} = -9 & C_{12} = 2 & C_{13} = 6 \\ C_{21} = -1 & C_{22} = -5 & C_{23} = -15 \\ C_{31} = -3 & C_{32} = -15 & C_{33} = 2 \end{bmatrix}$

$$\text{adj } A = \begin{bmatrix} -9 & 2 & 6 \\ -1 & -5 & -15 \\ -3 & -15 & 2 \end{bmatrix}' \Rightarrow \text{adj } A = \begin{bmatrix} -9 & -1 & -3 \\ 2 & -5 & -15 \\ 6 & -15 & 2 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{|A|}(\text{adj } A) = \frac{1}{(-47)} \begin{bmatrix} -9 & -1 & -3 \\ 2 & -5 & -15 \\ 6 & -15 & 2 \end{bmatrix} = \frac{1}{47} \begin{bmatrix} 9 & 1 & 3 \\ -2 & 5 & 15 \\ -6 & 15 & -2 \end{bmatrix}$$

Now $AX = B, X = A^{-1}B \Rightarrow X = \frac{1}{47} \begin{bmatrix} 9 & 1 & 3 \\ -2 & 5 & 15 \\ -6 & 15 & -2 \end{bmatrix} \begin{bmatrix} -7 \\ 1 \\ 5 \end{bmatrix}$

$$\Rightarrow X = \frac{1}{47} \begin{bmatrix} -63 + 1 + 15 \\ 14 + 5 + 75 \\ 42 + 15 - 10 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{47} \begin{bmatrix} -47 \\ 94 \\ 47 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} -1 \\ 2 \\ 1 \end{bmatrix}. \text{ Hence, } x = -1, y = 2 \text{ and } z = 1$$

34. $x - 2y + z = 0; y - z = 2; 2x - 3z = 10$

Sol. $A = \begin{bmatrix} 1 & -2 & 1 \\ 0 & 1 & -1 \\ 2 & 0 & -3 \end{bmatrix}, x = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, B = \begin{bmatrix} 0 \\ 2 \\ 10 \end{bmatrix}$

$$|A| = \begin{vmatrix} 1 & -2 & 1 \\ 0 & 1 & -1 \\ 2 & 0 & -3 \end{vmatrix} \Rightarrow |A| = 1(-3+0) + 2(0+2) + 1(0-2) \Rightarrow |A| = -3 + 4 - 2 \Rightarrow |A| = (-1)$$

Hence, A^{-1} exist.

$$\therefore \text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}'$$

MillionStars edu
Think, Learn & Practice

<https://millionstar.godaddy.com/>

So the cofactor of the element of matrix A are,

$$\begin{bmatrix} C_{11} = -3 & C_{12} = -2 & C_{13} = -2 \\ C_{21} = -6 & C_{22} = -5 & C_{23} = -4 \\ C_{31} = 1 & C_{32} = 1 & C_{33} = 1 \end{bmatrix}$$

$$\text{adj } A = \begin{bmatrix} -3 & -2 & -2 \\ -6 & -5 & -4 \\ 1 & 1 & 1 \end{bmatrix} \Rightarrow \text{adj } A = \begin{bmatrix} -3 & -6 & 1 \\ -2 & -5 & 1 \\ -2 & -4 & 1 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{|A|}(\text{adj } A) = \frac{1}{(-1)} \begin{bmatrix} -3 & -6 & 1 \\ -2 & -5 & 1 \\ -2 & -4 & 1 \end{bmatrix} = \begin{bmatrix} 3 & 6 & -1 \\ 2 & 5 & -1 \\ 2 & 4 & -1 \end{bmatrix}$$

$$\text{Now, } AX = B, X = A^{-1}B = \begin{bmatrix} 3 & 6 & -1 \\ 2 & 5 & -1 \\ 2 & 4 & -1 \end{bmatrix} \begin{bmatrix} 0 \\ 2 \\ 10 \end{bmatrix} \Rightarrow X = \begin{bmatrix} 0+12-10 \\ 0+10-10 \\ 0+8-10 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 2 \\ 0 \\ -2 \end{bmatrix}$$

Hence, $x = 2$, $y = 0$ and $z = -2$

35. $x - y = 3$; $2x + 3y + 4z = 17$; $y + 2z = 7$

Sol. $A = \begin{bmatrix} 1 & -1 & 0 \\ 2 & 3 & 4 \\ 0 & 1 & 2 \end{bmatrix}$, $X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$, $B = \begin{bmatrix} 3 \\ 17 \\ 7 \end{bmatrix}$

$$|A| = \begin{vmatrix} 1 & -1 & 0 \\ 2 & 3 & 4 \\ 0 & 1 & 2 \end{vmatrix} \Rightarrow |A| = 1(6-4) + 1(4-0) + 0 \Rightarrow |A| = 2+4 \Rightarrow |A| = 6$$

Hence, A^{-1} exist.

$$\therefore \text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}$$

So the cofactor of the element of matrix A are,

$$\begin{bmatrix} C_{11} = 2 & C_{12} = -4 & C_{13} = 2 \\ C_{21} = 2 & C_{22} = 2 & C_{23} = -1 \\ C_{31} = -4 & C_{32} = -4 & C_{33} = 5 \end{bmatrix}$$

$$\text{adj } A = \begin{bmatrix} 2 & -4 & 2 \\ 2 & 2 & -1 \\ -4 & -4 & 5 \end{bmatrix} \Rightarrow \text{adj } A = \begin{bmatrix} 2 & 2 & -4 \\ -4 & 2 & -1 \\ 2 & -1 & 5 \end{bmatrix}$$

$$A^{-1} = \frac{1}{|A|}(\text{adj } A) = \frac{1}{6} \begin{bmatrix} 2 & 2 & -4 \\ -4 & 2 & -1 \\ 2 & -1 & 5 \end{bmatrix} \text{ Now, } AX = B, X = A^{-1}B = \frac{1}{6} \begin{bmatrix} 2 & 2 & -4 \\ -4 & 2 & -1 \\ 2 & -1 & 5 \end{bmatrix} \begin{bmatrix} 3 \\ 17 \\ 7 \end{bmatrix}$$

$$\Rightarrow X = \frac{1}{6} \begin{bmatrix} 6+34-28 \\ -12+34-28 \\ 6-17+35 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{6} \begin{bmatrix} 12 \\ -6 \\ 24 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \\ 4 \end{bmatrix} \text{ Hence, } x = 2, y = -1 \text{ and } z = 4$$

36. $4x + 3y + 2z = 60$; $x + 2y + 3z = 45$; $6x + 2y + 3z = 70$

Sol. The system can be written as $AX = B \Rightarrow X = A^{-1}B$ (i)

Where $A = \begin{bmatrix} 4 & 3 & 2 \\ 1 & 2 & 3 \\ 6 & 2 & 3 \end{bmatrix}$, $X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$ and $B = \begin{bmatrix} 60 \\ 45 \\ 70 \end{bmatrix}$

$$|A| = 4(6-6) - 3(3-18) + 2(2-12) = 0 + 45 - 20 = 25 \neq 0$$

For adj A

$$A_{11} = 6-6=0 \quad A_{21} = -(9-4)=-5 \quad A_{31} = (9-4)=5$$

$$A_{12} = -(3-18)=15 \quad A_{22} = (12-12)=0 \quad A_{32} = -(12-2)=-10$$

$$A_{13} = (2-12)=-10 \quad A_{23} = -(8-18)=10 \quad A_{33} = (8-3)=5$$

$$\therefore \text{adj } A = \begin{bmatrix} 0 & 15 & -10 \\ -5 & 0 & 10 \\ 5 & -10 & 5 \end{bmatrix}^T = \begin{bmatrix} 0 & -5 & 5 \\ 15 & 0 & -10 \\ -10 & 10 & 5 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{25} \begin{bmatrix} 0 & -5 & 5 \\ 15 & 0 & -10 \\ -10 & 10 & 5 \end{bmatrix} = \frac{5}{25} \begin{bmatrix} 0 & -1 & 1 \\ 3 & 0 & -2 \\ -2 & 2 & 1 \end{bmatrix} = \frac{1}{5} \begin{bmatrix} 0 & -1 & 1 \\ 3 & 0 & -2 \\ -2 & 2 & 1 \end{bmatrix}$$

Now putting values in (i) we get

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{5} \begin{bmatrix} 0 & -1 & 1 \\ 3 & 0 & -2 \\ -2 & 2 & 1 \end{bmatrix} \begin{bmatrix} 60 \\ 45 \\ 70 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{5} \begin{bmatrix} 0-45+70 \\ 180+0-140 \\ -120+90+70 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{5} \begin{bmatrix} 25 \\ 40 \\ 40 \end{bmatrix} = \begin{bmatrix} 5 \\ 8 \\ 8 \end{bmatrix}$$

Hence $x=5, y=8, z=8$

37. If $A = \begin{bmatrix} 2 & -3 & 5 \\ 3 & 2 & -4 \\ 1 & 1 & -2 \end{bmatrix}$. Find A^{-1} .

Using A^{-1} , solve the following system of equations : $2x-3y+5z=11$; $3x+2y-4z=-5$; $x+y-2z=-3$

Sol. Let $|A| = \begin{vmatrix} 2 & -3 & 5 \\ 3 & 2 & -4 \\ 1 & 1 & -2 \end{vmatrix} = 2(-4+4) + 3(-6+4) + 5(3-2) = 2(0) + 3(-2) + 5(1) = -6+5 = -1$

$$\therefore |A| \neq 0, A^{-1} \text{ exist. } \therefore \text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}^T$$

<https://millionstar.godaddy.com/>

So the cofactor of the element of matrix A are

$$\begin{bmatrix} C_{11}=0 & C_{12}=2 & C_{13}=1 \\ C_{21}=-1 & C_{22}=-9 & C_{23}=-5 \\ C_{31}=2 & C_{32}=23 & C_{33}=13 \end{bmatrix}$$

$$\text{adj } A = \begin{bmatrix} 0 & 2 & 1 \\ -1 & -9 & -5 \\ 2 & 23 & 13 \end{bmatrix}^T = \begin{bmatrix} 0 & -1 & 2 \\ 2 & -9 & 23 \\ 1 & -5 & 13 \end{bmatrix} \Rightarrow A^{-1} = \frac{1}{|A|}(\text{adj } A) = \frac{1}{-1} \begin{bmatrix} 0 & -1 & 2 \\ 2 & -9 & 23 \\ 1 & -5 & 13 \end{bmatrix}$$

$$\text{Now, } AX = B, X = A^{-1} \cdot B \Rightarrow X = - \begin{bmatrix} 0 & -1 & 2 \\ 2 & -9 & 23 \\ 1 & -5 & 13 \end{bmatrix} \begin{bmatrix} 11 \\ -5 \\ -3 \end{bmatrix} \Rightarrow X = - \begin{bmatrix} 0+5-6 \\ 22+45-69 \\ 11+25-39 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = - \begin{bmatrix} -1 \\ -2 \\ -3 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \text{ Hence, } x=1, y=2 \text{ \& } z=3$$

38. If $A = \begin{bmatrix} 2 & 1 & 1 \\ 1 & -2 & -1 \\ 0 & 3 & -5 \end{bmatrix}$, find A^{-1} . Using A^{-1} , solve the following system of linear equations :

$$2x + y + z = 1; x - 2y - z = \frac{3}{2}; 3y - 5z = 9$$

Sol. $A = \begin{bmatrix} 2 & 1 & 1 \\ 1 & -2 & -1 \\ 0 & 3 & -5 \end{bmatrix}$, $x = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$, $B = \begin{bmatrix} 1 \\ 3/2 \\ 9 \end{bmatrix}$

$$|A| = \begin{vmatrix} 2 & 1 & 1 \\ 1 & -2 & -1 \\ 0 & 3 & -5 \end{vmatrix} \Rightarrow |A| = 2(10+3) - 1(-5+0) + 1(3+0) \Rightarrow |A| = 26+5+3 \Rightarrow |A| = 34$$

Hence, A^{-1} exist. $\therefore \text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}^T$

So the cofactor of the element of matrix A are,

$$\begin{bmatrix} C_{11}=13 & C_{12}=5 & C_{13}=3 \\ C_{21}=8 & C_{22}=-10 & C_{23}=-6 \\ C_{31}=1 & C_{32}=3 & C_{33}=-5 \end{bmatrix}$$

$$\text{adj } A = \begin{bmatrix} 13 & 5 & 3 \\ 8 & -10 & -6 \\ 1 & 3 & -5 \end{bmatrix}^T \Rightarrow \text{adj } A = \begin{bmatrix} 13 & 8 & 1 \\ 5 & -10 & 3 \\ 3 & -6 & -5 \end{bmatrix}$$

$$A^{-1} = \frac{1}{|A|}(\text{adj } A) = \frac{1}{34} \begin{bmatrix} 13 & 8 & 1 \\ 5 & -10 & 3 \\ 3 & -6 & -5 \end{bmatrix} \text{ Now, } AX = B, X = A^{-1}B = \frac{1}{34} \begin{bmatrix} 13 & 8 & 1 \\ 5 & -10 & 3 \\ 3 & -6 & -5 \end{bmatrix} \begin{bmatrix} 1 \\ 3/2 \\ 9 \end{bmatrix}$$

$$\Rightarrow X = \frac{1}{34} \begin{bmatrix} 13+12+9 \\ 5-15+27 \\ 3-9-45 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{34} \begin{bmatrix} 34 \\ 17 \\ -51 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 1/2 \\ -3/2 \end{bmatrix} \text{ Hence, } x=1, y=\frac{1}{2} \text{ and } z=-\frac{3}{2}$$

39. If $A = \begin{bmatrix} 1 & -2 & 0 \\ 2 & 1 & 3 \\ 0 & -2 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 7 & 2 & -6 \\ -2 & 1 & -3 \\ -4 & 2 & 5 \end{bmatrix}$, find AB

Hence solve the system of equations $x - 2y = 10$, $2x + y + 3z = 8$ and $-2y + z = 7$

Sol. Given $A = \begin{bmatrix} 1 & -2 & 0 \\ 2 & 1 & 3 \\ 0 & -2 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 7 & 2 & -6 \\ -2 & 1 & -3 \\ -4 & 2 & 5 \end{bmatrix}$

$$A \cdot B = \begin{bmatrix} 1 & -2 & 0 \\ 2 & 1 & 3 \\ 0 & -2 & 1 \end{bmatrix} \begin{bmatrix} 7 & 2 & -6 \\ -2 & 1 & -3 \\ -4 & 2 & 5 \end{bmatrix}$$

$$= \begin{bmatrix} 7+4+0 & 2-2+0 & -6+6+0 \\ 14-2-12 & 4+1+6 & -12-3+15 \\ 0+4-4 & 0-2+2 & 0+6+5 \end{bmatrix} = \begin{bmatrix} 11 & 0 & 0 \\ 0 & 11 & 0 \\ 0 & 0 & 11 \end{bmatrix} = 11 \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = 11I$$

$$\Rightarrow A \cdot \left(\frac{1}{11} B \right) = I$$

$$\Rightarrow A^{-1} = \frac{1}{11} B = \frac{1}{11} \begin{bmatrix} 7 & 2 & -6 \\ -2 & 1 & -3 \\ -4 & 2 & 5 \end{bmatrix}$$

Given system of equations are

$$x - 2y = 10$$

$$2x + y + 3z = 8$$

$$-2y + z = 7$$

This system may be written as $AX = C$ where

Where $A = \begin{bmatrix} 1 & -2 & 0 \\ 2 & 1 & 3 \\ 0 & -2 & 1 \end{bmatrix}$, $X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$ & $C = \begin{bmatrix} 10 \\ 8 \\ 7 \end{bmatrix}$

Now $AX = C$

$$\Rightarrow X = A^{-1}C$$

$$\Rightarrow X = \frac{1}{11} \begin{bmatrix} 7 & 2 & -6 \\ -2 & 1 & -3 \\ -4 & 2 & 5 \end{bmatrix} \begin{bmatrix} 10 \\ 8 \\ 7 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{11} \begin{bmatrix} 70+16-42 \\ -20+8-21 \\ -40+16+35 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{11} \begin{bmatrix} 44 \\ -33 \\ 11 \end{bmatrix} = \begin{bmatrix} 4 \\ -3 \\ 1 \end{bmatrix}$$

Hence $x = 4$, $y = -3$ & $z = 1$

40. Using matrices, solve the following system of equations :

$$\frac{2}{x} - \frac{3}{y} + \frac{3}{z} = 10, \frac{1}{x} + \frac{1}{y} + \frac{1}{z} = 10, \frac{3}{x} - \frac{1}{y} + \frac{2}{z} = 13$$



$$A = \begin{bmatrix} 2 & -3 & 3 \\ 1 & 1 & 1 \\ 3 & -1 & 2 \end{bmatrix}, X = \begin{bmatrix} 1/x \\ 1/y \\ 1/z \end{bmatrix}, B = \begin{bmatrix} 18 \\ 10 \\ 13 \end{bmatrix}$$

$$|A| = \begin{vmatrix} 2 & -3 & 3 \\ 1 & 1 & 1 \\ 3 & -1 & 2 \end{vmatrix} \Rightarrow |A| = 2(2+1) + 3(2-3) + 3(-1-3) \Rightarrow |A| = 6 - 3 - 12 \Rightarrow |A| = (-9)$$

Hence, A^{-1} exist. $\therefore \text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}'$

So the cofactor of the element of matrix A are, $\begin{bmatrix} C_{11}=3 & C_{12}=1 & C_{13}=-4 \\ C_{21}=3 & C_{22}=-5 & C_{23}=-7 \\ C_{31}=-6 & C_{32}=1 & C_{33}=5 \end{bmatrix}$

$$\text{adj } A = \begin{bmatrix} 3 & 1 & -4 \\ 3 & -5 & -7 \\ -6 & 1 & 5 \end{bmatrix}' \Rightarrow \text{adj } A = \begin{bmatrix} 3 & 3 & -6 \\ 1 & -5 & 1 \\ -4 & -7 & 5 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{|A|} (\text{adj } A) = \frac{1}{(-9)} \begin{bmatrix} 3 & 3 & -6 \\ 1 & -5 & 1 \\ -4 & -7 & 5 \end{bmatrix} = \frac{1}{9} \begin{bmatrix} -3 & -3 & 6 \\ -1 & 5 & -1 \\ 4 & 7 & -5 \end{bmatrix}$$

Now, $AX = B, X = A^{-1}B = \frac{1}{9} \begin{bmatrix} -3 & -3 & 6 \\ -1 & 5 & -1 \\ 4 & 7 & -5 \end{bmatrix} \begin{bmatrix} 18 \\ 10 \\ 13 \end{bmatrix} \Rightarrow X = \frac{1}{9} \begin{bmatrix} -30 - 30 + 78 \\ -10 + 50 - 13 \\ 40 + 70 - 65 \end{bmatrix}$

$$\Rightarrow \begin{bmatrix} 1/x \\ 1/y \\ 1/z \end{bmatrix} = \frac{1}{9} \begin{bmatrix} 18 \\ 27 \\ 45 \end{bmatrix} \Rightarrow \begin{bmatrix} 1/x \\ 1/y \\ 1/z \end{bmatrix} = \begin{bmatrix} 2 \\ 3 \\ 5 \end{bmatrix}. \text{ Hence, } x = \frac{1}{2}, y = \frac{1}{3} \text{ and } z = \frac{1}{5}$$

41. $\frac{1}{x} - \frac{1}{y} + \frac{1}{z} = 4; \frac{2}{x} + \frac{1}{y} - \frac{3}{z} = 0; \frac{1}{x} + \frac{1}{y} + \frac{1}{z} = 2 \quad (x, y, z \neq 0)$

Sol. Given equations are

$$\frac{1}{x} - \frac{1}{y} + \frac{1}{z} = 4 \quad \dots (i), \quad \frac{2}{x} + \frac{1}{y} - \frac{3}{z} = 0 \quad \dots (ii), \quad \frac{1}{x} + \frac{1}{y} + \frac{1}{z} = 2 \quad \dots (iii)$$

Let $A = \begin{bmatrix} 1 & -1 & 1 \\ 2 & 1 & -3 \\ 1 & 1 & 1 \end{bmatrix}, X = \begin{bmatrix} 1/x \\ 1/y \\ 1/z \end{bmatrix}$ and $B = \begin{bmatrix} 4 \\ 0 \\ 2 \end{bmatrix}$

Here $A \cdot X = B$

$$\Rightarrow X = A^{-1}B \quad \dots (i)$$

Now, $|A| = \begin{vmatrix} 1 & -1 & 1 \\ 2 & 1 & -3 \\ 1 & 1 & 1 \end{vmatrix}$

$$= \begin{vmatrix} 0 & 0 & 1 \\ 5 & -2 & -3 \\ 0 & 2 & 1 \end{vmatrix} \begin{cases} C_1 \rightarrow C_1 - C_3 \\ C_2 \rightarrow C_2 + C_3 \end{cases}$$

$$= \begin{vmatrix} 5 & -2 \\ 0 & 2 \end{vmatrix} \quad \{\text{expanding by } R_1\}$$

$$= 10$$

$$\because |A| \neq 0 \quad \therefore A^{-1} \text{ exists}$$

Co-factors of A :

$$A_{11} = \begin{vmatrix} 1 & -3 \\ 1 & 1 \end{vmatrix} = 1 + 3 = 4$$

$$A_{12} = - \begin{vmatrix} 2 & -3 \\ 1 & 1 \end{vmatrix} = -(2 + 3) = -5$$

$$A_{13} = \begin{vmatrix} 2 & 1 \\ 1 & 1 \end{vmatrix} = 2 - 1 = 1$$

$$A_{21} = - \begin{vmatrix} -1 & 1 \\ 1 & 1 \end{vmatrix} = -(1 - 1) = 0$$

$$A_{22} = \begin{vmatrix} 1 & 1 \\ 1 & 1 \end{vmatrix} = (1 - 1) = 0$$

$$A_{23} = - \begin{vmatrix} 1 & -1 \\ 1 & 1 \end{vmatrix} = -(1 + 1) = -2$$

$$A_{31} = \begin{vmatrix} -1 & 1 \\ 1 & -3 \end{vmatrix} = 3 - 1 = 2$$

$$A_{32} = - \begin{vmatrix} 1 & 1 \\ 2 & -3 \end{vmatrix} = -(-3 - 2) = 5$$

$$A_{33} = \begin{vmatrix} 1 & -1 \\ 2 & 1 \end{vmatrix} = 1 + 2 = 3$$

$$\text{Now, } \text{Adj}(A) = \begin{bmatrix} A_{11} & A_{21} & A_{31} \\ A_{12} & A_{22} & A_{32} \\ A_{13} & A_{23} & A_{33} \end{bmatrix} = \begin{bmatrix} 4 & 0 & 2 \\ -5 & 0 & 5 \\ 1 & -2 & 3 \end{bmatrix}$$

$$A^{-1} = \frac{1}{|A|} \cdot \text{adj}(A) = \frac{1}{10} \begin{bmatrix} 4 & 0 & 2 \\ -5 & 0 & 5 \\ 1 & -2 & 3 \end{bmatrix}$$

$$\text{Now, } X = A^{-1}B$$

$$\Rightarrow \begin{bmatrix} 1/x \\ 1/y \\ 1/z \end{bmatrix} = \frac{1}{10} \begin{bmatrix} 4 & 0 & 2 \\ -5 & 0 & 5 \\ 1 & -2 & 3 \end{bmatrix} \begin{bmatrix} 4 \\ 0 \\ 2 \end{bmatrix} = \frac{1}{10} \begin{bmatrix} 16 + 0 + 4 \\ -20 + 0 + 10 \\ 4 + 0 + 6 \end{bmatrix} = \frac{1}{10} \begin{bmatrix} 20 \\ -10 \\ 10 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \\ 1 \end{bmatrix}$$

$$\therefore \frac{1}{x} = 2, y = -1 \text{ \& } \frac{1}{z} = 1$$

$$\Rightarrow x = \frac{1}{2}, y = 1 \text{ \& } z = 1 \quad \text{https://millionstar.godaddysites.com/}$$

42. The sum of three numbers is 2. If twice the second number is added to the sum of first and third, we get 1. on adding the sum of second and third numbers to five times the first, we get 6. Find the three numbers by using matrices.

Sol. Let these three number, first number be x , second number by y and third number be z .

$$x + y + z = 2 \quad \dots(1)$$

$$x + 2y + z = 1 \quad \dots(2)$$

$$5x + y + z = 6 \quad \dots(3)$$

$$A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 1 \\ 5 & 1 & 1 \end{bmatrix}, X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, B = \begin{bmatrix} 2 \\ 1 \\ 6 \end{bmatrix}$$

$$|A| = 1 \begin{vmatrix} 2 & 1 \\ 1 & 1 \end{vmatrix} - 1 \begin{vmatrix} 1 & 1 \\ 5 & 1 \end{vmatrix} + 1 \begin{vmatrix} 1 & 2 \\ 5 & 1 \end{vmatrix} \Rightarrow |A| = (2-1) - 1(1-5) + 1(1-10) \Rightarrow |A| = 1 + 4 - 9 = -4$$

Hence, A^{-1} exist. $\therefore \text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}'$

So the cofactor of the element of matrix A are, $\begin{bmatrix} C_{11}=1 & C_{12}=4 & C_{13}=-9 \\ C_{21}=0 & C_{22}=-4 & C_{23}=4 \\ C_{31}=-1 & C_{32}=0 & C_{33}=1 \end{bmatrix}$

$$(\text{adj } A) = \begin{bmatrix} 1 & 4 & -9 \\ 0 & -4 & 4 \\ -1 & 0 & 1 \end{bmatrix} \Rightarrow (\text{adj } A) = \begin{bmatrix} 1 & 0 & -1 \\ 4 & -4 & 0 \\ -9 & 4 & 1 \end{bmatrix}$$

$$\text{Now, } AX = B \Rightarrow X = A^{-1}B \Rightarrow X = -\frac{1}{4} \begin{bmatrix} 1 & 0 & -1 \\ 4 & -4 & 0 \\ -9 & 4 & 1 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \\ 6 \end{bmatrix} \Rightarrow X = -\frac{1}{4} \begin{bmatrix} 2+0-6 \\ 8-4+0 \\ -18+4+6 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = -\frac{1}{4} \begin{bmatrix} -4 \\ 4 \\ -8 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix} \quad \text{Hence } x = 1, y = -1, z = 2$$

43. The cost of 4 kg potato, 3 kg wheat and 2 kg rice is Rs 60. The cost of 1 kg potato, 2 kg wheat and 3 kg rice is Rs 45. The cost of 6 kg potato, 2 kg wheat and 3 kg rice is Rs 70. Find the cost of each item per kg by matrix method.

Sol. Let these three item, potato, wheat and rice be x, y, z respectively.

$$4x + 3y + 2z = 60 \quad \dots(1)$$

$$x + 2y + 3z = 45 \quad \dots(2)$$

$$6x + 2y + 3z = 70 \quad \dots(3)$$

$$A = \begin{bmatrix} 4 & 3 & 2 \\ 1 & 2 & 3 \\ 6 & 2 & 3 \end{bmatrix}, X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, B = \begin{bmatrix} 60 \\ 45 \\ 70 \end{bmatrix}$$

$$|A| = 4 \begin{vmatrix} 2 & 3 \\ 2 & 3 \end{vmatrix} - 3 \begin{vmatrix} 1 & 5 \\ 6 & 3 \end{vmatrix} + 2 \begin{vmatrix} 1 & 2 \\ 6 & 2 \end{vmatrix} \Rightarrow |A| = 4(6-6) - 3(3-18) + 2(2-12)$$

$$\Rightarrow |A| = 4(0) - 3(-15) + 2(-10) \Rightarrow |A| = 45 - 20 = 25. \text{ Hence, } A^{-1} \text{ exist.}$$

$$\therefore \text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}$$

So the cofactor of the element of matrix A are, $\begin{bmatrix} C_{11}=0 & C_{12}=15 & C_{13}=-10 \\ C_{21}=-5 & C_{22}=0 & C_{23}=10 \\ C_{31}=5 & C_{32}=-10 & C_{33}=5 \end{bmatrix}$

$$\text{adj } A = \begin{bmatrix} 0 & 15 & -10 \\ -5 & 0 & 10 \\ 5 & -10 & 5 \end{bmatrix} \Rightarrow \text{adj } A = \begin{bmatrix} 0 & -5 & 5 \\ 15 & 0 & -10 \\ -10 & 10 & 5 \end{bmatrix}$$

$$A^{-1} = \frac{1}{|A|} (\text{adj } A) = \frac{1}{25} \begin{bmatrix} 0 & -5 & 5 \\ 15 & 0 & -10 \\ -10 & 10 & 5 \end{bmatrix}$$

Now $AX = B, X = A^{-1}B \Rightarrow X = \frac{1}{25} \begin{bmatrix} 0 & -5 & 5 \\ 15 & 0 & -10 \\ -10 & 10 & 5 \end{bmatrix} \begin{bmatrix} 60 \\ 45 \\ 70 \end{bmatrix}$

$$\Rightarrow x = \frac{1}{25} \begin{bmatrix} 0 - 225 + 350 \\ 900 + 0 - 700 \\ -600 + 450 + 350 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{25} \begin{bmatrix} 125 \\ 200 \\ 200 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 5 \\ 8 \\ 8 \end{bmatrix}$$

$$\therefore x = \text{Rs. } 5, y = \text{Rs. } 8, z = \text{Rs. } 8$$

44. An amount of Rs. 5000 is put into three investments at 6%, 7% and 8% per annum respectively. The total annual income from these investments is Rs 358. If the total annual income from first two investments is Rs 70 more than the income from the third, find the amount of each investment by the matrix method.

Sol. Let these investment be Rs x , Rs y and Rs. z respectively.

$$x + y + z = 5000 \quad \dots(1) \quad \Rightarrow \frac{6x}{100} + \frac{7y}{100} + \frac{8z}{100} = 358$$

$$6x + 7y + 8z = 35800 \quad \dots(2) \quad \text{and} \quad \frac{6x}{100} + \frac{7y}{100} = \frac{8z}{100} + 70$$

$$6x + 7y - 8z = 7000 \quad \dots(3)$$

$$A = \begin{bmatrix} 1 & 1 & 1 \\ 6 & 7 & 8 \\ 6 & 7 & -8 \end{bmatrix}, X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, B = \begin{bmatrix} 5000 \\ 35800 \\ 7000 \end{bmatrix}$$

$$|A| = 1 \begin{vmatrix} 7 & 8 \\ 7 & -8 \end{vmatrix} - 1 \begin{vmatrix} 6 & 8 \\ 6 & -8 \end{vmatrix} + 1 \begin{vmatrix} 6 & 7 \\ 6 & 7 \end{vmatrix} \Rightarrow |A| = 1(-56 - 56) - 1(-48 - 48) + 1(42 - 42)$$

$$\Rightarrow |A| = (-112) - 1(-96) + 1(0) \Rightarrow |A| = -112 + 96 \Rightarrow |A| = -16$$

<https://millionstar.godaddysites.com/>

$$\text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}$$

So the cofactor of the element of matrix A are, $\begin{bmatrix} C_{11} = -112 & C_{12} = 96 & C_{13} = 0 \\ C_{21} = 15 & C_{22} = -14 & C_{23} = -1 \\ C_{31} = 1 & C_{32} = -2 & C_{33} = 1 \end{bmatrix}$

$$\text{adj } A = \begin{bmatrix} -112 & 96 & 0 \\ 15 & -14 & -1 \\ 1 & -2 & 1 \end{bmatrix} \Rightarrow \text{adj } A = \begin{bmatrix} -112 & 15 & 1 \\ 96 & -14 & -2 \\ 0 & -1 & 1 \end{bmatrix}$$

$$A^{-1} = \frac{1}{|A|} (\text{adj } A) \Rightarrow A^{-1} = -\frac{1}{16} \begin{bmatrix} -112 & 15 & 1 \\ 96 & -14 & -2 \\ 0 & -1 & 1 \end{bmatrix}$$

$$\text{Now, } AX = B \Rightarrow X = A^{-1}B \Rightarrow X = \frac{1}{-16} \begin{bmatrix} -112 & 15 & 1 \\ 96 & -14 & -2 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} 5000 \\ 35800 \\ 7000 \end{bmatrix}$$

$$\Rightarrow X = -\frac{1}{16} \begin{bmatrix} -560000 + 537000 + 7000 \\ 480000 - 501200 - 14000 \\ 0 - 35800 + 7000 \end{bmatrix} \Rightarrow X = -\frac{1}{16} \begin{bmatrix} -16000 \\ -35200 \\ -28800 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1000 \\ 2200 \\ 1800 \end{bmatrix}$$

Hence $x = 1000$, $y = 2200$, $z = 1800$

45. Two schools A and B want to award their selected students on the values of sincerity, truthfulness and helpfulness. The school A wants to award Rs. X each rs. Y each and rs. Z each for the three respective values to 3, 2 and 1 students respectively with a total award money of Rs. 1600. school b wants to spend Rs. 2300 to award its 4, 1 and 3 students on the respective values (by giving the same award money to the three values as before). If the total amount of award for one prize on each value of Rs. 900, using matrices, find the award money for each value. Apart from these three values suggest one more value which should be considered for award

Sol.

Number of student of school	Sincerity	Truthfulness	Helpfulness
A	3	2	1
B	4	1	3
One student for each value	1	1	1

$$X = \begin{bmatrix} x \\ y \\ z \end{bmatrix} \text{ where } x, y \text{ and } z \text{ are rupees mentioned as it is the question, for sincerity truthfulness and}$$

Helpfulness respectively

$$E = \begin{bmatrix} 1600 \\ 2300 \\ 900 \end{bmatrix} \text{ is a matrix representing total award money for school A, B and for one prize for each value}$$



We can represent the given system as : $DX = E$

$$\text{Or } \begin{bmatrix} 3 & 2 & 1 \\ 4 & 1 & 3 \\ 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1600 \\ 2300 \\ 900 \end{bmatrix}$$

Solution of the matrix equation exist if $|D| \neq 0$

$$\begin{aligned} \text{i.e. } \begin{bmatrix} 3 & 2 & 1 \\ 4 & 1 & 3 \\ 1 & 1 & 1 \end{bmatrix} &= 3[1-3] - 2[4-3] + 1[4-1] \\ &= -6 - 2 + 3 = -5 \end{aligned}$$

Therefore, the solution of the matrix equation is $X = D^{-1}E$

$$\text{To find } D^{-1}; D^{-1} = \frac{1}{|D|} \text{adj}(D)$$

Cofactor matrix of D

$$= \begin{bmatrix} -2 & -1 & 3 \\ -1 & 2 & -1 \\ 5 & -5 & -5 \end{bmatrix}$$

Adjoint of $D = \text{adj}(D)$

$$= \begin{bmatrix} -2 & -1 & 5 \\ -1 & 2 & -5 \\ 3 & -1 & -5 \end{bmatrix} \quad \{\text{transpose of cofactor matrix}\}$$

$$D^{-1} = \frac{1}{-5} \begin{bmatrix} -2 & -1 & 5 \\ -1 & 2 & -5 \\ 3 & -1 & -5 \end{bmatrix}$$

Now, $X = D^{-1}E$

$$= \frac{1}{-5} \begin{bmatrix} -2 & -1 & 5 \\ -1 & 2 & -5 \\ 3 & -1 & -5 \end{bmatrix} \begin{bmatrix} 1600 \\ 2300 \\ 900 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 200 \\ 300 \\ 400 \end{bmatrix}$$

$$x = 200, y = 300, z = 400$$

Award can also be given for punctuality