


Exercise 11A

Question 1:

Let AB be a chord of the given circle with centre O and radius 10 cm.Then, OA = 10 cm and AB = 16 cm. From O, draw OL \perp AB. We know that the perpendicular from the centre of a circle to a chord bisects the chord.

$$AL = \frac{1}{2} \times AB$$
$$= \left(\frac{1}{2} \times 16\right) \text{ cm} = 8 \text{ cm}.$$




$$OA^{2} = OL^{2} + AL^{2}$$
⇒
$$OL^{2} = OA^{2} - AL^{2}$$

$$= 10^{2} - 8^{2}$$

$$= 100 - 64 = 36$$
∴
$$OL = \sqrt{36} = 6 \text{ cm}.$$

... The distance of the chord from the centre is 6 cm.




Question 2:

Let AB be the chord of the given circle with centre O and radius 5 cm.

From O, draw OL ⊥ AB

Then, OA = 5 cm and OL = 3 cm [given]

We know that the perpendicular from the centre of a circle to a chord bisects the chord.



Now, in right angled Δ OLA, we have

$$OA^{2} = AL^{2} + OL^{2}$$

$$\Rightarrow AL^{2} = OA^{2} - OL^{2}$$

$$\Rightarrow AL^{2} = 5^{2} - 3^{2}$$

$$= 25 - 9 = 16$$

$$\therefore AL = \sqrt{16} = 4 \text{ cm}$$
So, $AB = 2 \text{ AL}$

$$= (2 \times 4) \text{ cm} = 8 \text{ cm}$$


$$\therefore \text{ the length of the chord is 8 cm}$$

:, the length of the chord is 8 cm.





Let AB be the chord of the given airde with centre O.Draw OL \perp AB.



Then, OLis the distance from the centre to the chord. So, we have AB = 30 cm and AC = 8 cm

We know that the perpendicular from the centre of a circle to a circle bisects the chord.

$$AL = \frac{1}{2} \times AB$$
$$= \left(\frac{1}{2} \times 30\right) \text{ cm} = 15 \text{ cm}$$

Now, in right angled △OLA we have,


$$OA^2 = OL^2 + AL^2$$

= $8^2 + 15^2$
= $64 + 225 = 289$
 $OA = \sqrt{289} = 17 \text{ cm}$

... the radius of the circle is 17 cm.

Question 4:

(i)Let AB and CD be two chords of a circle such that AB || CD which are on the same side of the circle Also AB = 8 cm and CD = 6 cm OB = OD = 5 cm Join OL and LM Since the perpendicular from the centre of a circle to a chord bisects the chord.



We have
$$LB = \frac{1}{2} \times AB$$
$$= \left(\frac{1}{2} \times 8\right) \text{ cm} = 4 \text{ cm}$$

and
$$MD = \frac{1}{2} \times CD$$

$$=\left(\frac{1}{2}\times 6\right)$$
 cm $=3$ cm

Now in right angled Δ BLO

$$OB^2 = LB^2 + LO^2$$

$$LO^2 = OB^2 - LB^2$$

$$\Rightarrow LO^{2} = OB^{2} - LB^{2}$$

$$\Rightarrow = 5^{2} - 4^{2}$$

∴ LO =
$$\sqrt{9}$$
 = 3 cm.

Again in right angled ∆DMO

$$OD^2 = MD^2 + MO^2$$

$$\Rightarrow$$
 MO² = OD² - MD²

$$=5^2-3^2$$


$$= 25-9 = 16$$

MO $= \sqrt{16} = 4 \text{ cm}$

... The distance between the chords =
$$(4-3)$$
 cm = 1 cm.



(ii)Let AB and CD be two chords of a circle such that AB || CD and they are on the opposite sides of the centre.AB = 8 cm and $CD = 6 \text{ cm.Draw OL} \perp AB \text{ and } OM \perp CD.$



Join OA and OC

Then OA = OC = 5cm(radius)

Since the perpendicular from the centre of a circle to a chord bisects the chord, we have,

$$AL = \frac{1}{2}AB$$
$$= \left(\frac{1}{2} \times 8\right) cm = 4 cm.$$

Also

$$\begin{aligned} \mathsf{CM} &= \frac{1}{2}\mathsf{CD} \\ &= \left(\frac{1}{2} \times 6\right) \mathsf{cm} = 3 \; \mathsf{cm} \end{aligned}$$

Now in right angled △ OLA, we have

$$OA^{2} = AL^{2} + OL^{2}$$

$$\Rightarrow OL^{2} = OA^{2} - AL^{2}$$

$$= 5^{2} - 4^{2}$$

$$= 25 - 16 = 9 \text{ cm}$$

$$\therefore OL = \sqrt{9} = 3 \text{ cm}$$

Again in right angled Δ OMC, we have

$$OC^{2} = OM^{2} + CM^{2}$$

$$OM^{2} = OC^{2} - CM^{2}$$

$$= 5^{2} - 3^{2}$$

$$= 25 - 9 = 16$$

$$OM = \sqrt{16} = 4 \text{ cm}$$


:, the distance between the chords = (4+3)cm = 7 cm

Question 5:





Let AB and CD be two chords of a circle having centre O. $AB = 30 \, \text{cm}$ and $CD = 16 \, \text{cm}$.



Join AO and OC which are its radii. So AO = 17 cm and $\rm CO = 17$ cm.

Draw OM ⊥ CD and OL ⊥ AB.

Since the perpendicular from the centre of a circle to a chord bisects the chord, we have

$$AL = \frac{1}{2} \times AB$$

$$= \left(\frac{1}{2} \times 30\right) \text{cm} = 15 \text{ cm}$$

$$CM = \frac{1}{2} \times CD$$

$$= \left(\frac{1}{2} \times 16\right) \text{cm} = 8 \text{ cm}$$

Now, in right angled Δ ALO, we have

$$AO^2 = OL^2 + AL^2$$

$$\Rightarrow LO^2 = AO^2 - AL^2$$

$$= 17^2 - 15^2$$

$$= 289 - 225 = 64$$

$$\Rightarrow LO = \sqrt{64} = 8 \text{ cm}$$
Again, in right angled \triangle CMO, we have
$$CO^2 = CM^2 + OM^2$$

$$CO^2 = CM^2 + OM^2$$

⇒ $OM^2 = CO^2 - CM^2$
 $= 17^2 - 8^2$
 $= 289 - 64 = 225$
⇒ $OM = \sqrt{225} = 15 \text{ cm}$


 \therefore Distance between the chords = OM+OL =(8+15)cm = 23 cm.

Question 6:





CD is the diameter of a circle with centre O, and is perpendicular to chord AB. Join OA.



$$AB = 12$$
 cm and $CE = 3$ cm

[Given]

Let OA = OC = r cm

Then,

$$OE = (r-3) cm$$

Since the perpendicular from the centre of a circle to a chord bisects the chord, we have

$$AE = \frac{1}{2} \times AB$$
$$= \left(\frac{1}{2} \times 12\right) cm = 6 cm$$

Now, in right angled △OEA,

$$OA^{2} = OE^{2} + AE^{2}$$

$$\Rightarrow r^{2} = (r - 3)^{2} + 6^{2}$$

$$\Rightarrow r^{2} - 6r + 9 + 36$$

$$\Rightarrow r^{2} - r^{2} + 6r = 45$$

$$\Rightarrow 6r = 45$$

$$\Rightarrow r = \frac{45}{6} = 7.5 \text{ cm}$$

.. OA, the radius of the circle is 7.5 cm.


Question 7:

AB is the diameter of a circle with centre O which bisects the chord CD at point E. CE = ED = 8cm and EB = 4cm. Join OC.

Let OC = OB = r am.

Then,

$$OE = (r - 4) cm$$



Now, in right angled \triangle OEC

$$OC^{2} = OE^{2} + EC^{2}$$
 $r^{2} = (r - 4)^{2} + 8^{2}$
 $\Rightarrow r^{2} = r^{1} - 8r + 16 + 64$
 $\Rightarrow r^{2} = r^{2} - 8r + 80$
 $\Rightarrow r^{2} - r^{2} + 8r = 80$
 $\Rightarrow 8r = 80$
 $\Rightarrow r = \frac{80}{8} = 10 \text{ cm}$

.. the radius of the circle is 10 cm.

Question 8:



Given: OD \perp AB of a circle with centre O. BC is a diameter.

To Prove: AC || OD and AC= 2xOD

Construction: Join AC.

Proof: We know that the perpendicular from the centre of the circle to a chord bisects the chord.

Here OD ⊥AB

⇒D is the mid - point of AB

AD = BD

Also, O is the mid -point of BC

OC = OB

Now, in \triangle ABC, Dis the midpoint of AB and O is

the midpoint of BC.

Midpoint Theorem: The line segment joining the midpoints of any two sides of a triangle is parallel to the third side and equal to half of it.

∴ OD|| AC and OD=
$$\frac{1}{2}$$
AC

Question 9:

Sol.9. Given: O is the centre in which chords AB and CD intersects at P such that PO bisects ∠BPD.

AB = CD

Construction:Draw OE ⊥ AB and OF ⊥ CD



Proof: In \triangle OEP and \triangle OFP

∠ OEP = ∠ OFP Each equal to 90°

OP = OP common

∠ OPE=∠ OPF [Since OP bisects \(BPD \)]

Thus, by Angle-Side-Angle criterion of congruence, have,

 Δ OEP $\cong \Delta$ OFP [By ASA]

The corresponding the parts of the congruent triangles are equal

OE = OFC.P.C.T.

⇒ Chords AB and CD are equidistant from the centre O.

· chords equidistant AB = CD

from the centre are equal AB = CD

Question 10:

Given: AB and CD are two parallel chords of a circle with centre O.POQ is a diameter which is perpendicular to AB. To Prove: PF \(\text{CD} \) and CF = FD



Proof: AB | CD and POQ is a diameter.

∠PEB=90° Gven

Then, ∠PFD= ∠PEB [AB||CD, Corresponding angles]

PF _ CD Thus, OF L CD

We know that, the perpendicular from the centre of a circle to chord, bisects the chord.

CF = FD.






Question 11:

If possible let two different circles intersect at three distinct point A, B and C.

Then, these points are noncollinear. So a unique dirde can be drawn to pass through these points. This is a contradiction.

Question 12:



OA = 10 cm and AB = 12 cm

$$\therefore AD = \frac{1}{2} \times AB$$

$$AD = \left(\frac{1}{2} \times 12\right) \text{cm} = 6 \text{ cm}$$

Now in right angled \triangle ADO,

Again, we have O'A = 8 cm. In right angle △ ADO'

$$O'A^{2} = AD^{2} + O'D^{2}$$

$$O'D^{2} = O'A^{2} - AD^{2}$$

$$= 8' - 6^{2}$$

$$= 64 - 36 = 28$$

$$O'D = \sqrt{28} = 2\sqrt{7} \text{ cm}$$

$$OO' = (OD + O'D)$$

$$= (8 + 2\sqrt{7}) \text{ cm}$$


 \therefore the distance between their centres is $(8 + 2\sqrt{7})$ cm.

Question 13:

Given: Two equal cirles intersect at points P and Q.A straight

line through P meets the circles in Aand B.

To Prove: QA = QB Construction: Join PQ



Proof: Two circles will be congruent if and only if they have equal radii.

If two chords of a circle are equal then their corresponding arcs are congruent.

Here PQ is the common chord to both the circles.

Thus, their corresponding arcs are equal.

So,
$$arc PCQ = arc PDQ$$

[isosceles triangle,

QA = QBbase angles are equal]

· Million State & Practice



Question 14:

Given: AB and CD are the two chords of a circle with centre O. Diameter POQ bisects them at L and M.

To Prove :AB || CD.



Proof: AB and CD are two chords of a circle with centre O. Diameter POQ bisects them at L and M.

Then, OL \perp AB and, OM \perp CD \therefore \angle ALM = \angle LMD

∴ AB || CD [alternate angles are equal]

Question 15:

Two circles with centres A and B, having radii 5 cm and 3 cm touch each otherint ernally.

The perpendicular bisector of AB meets the bigger circle in P and Q.


Join AP.

Let PQ intersect AB at L.

Then, AB = (5-3) cm = 2 cm

Since PQ is the perpendicular bisector of AB, we have

$$AL = \frac{1}{2} \times AB$$
$$= \left(\frac{1}{2} \times 2\right) am = 1 am$$




Now,in right angle △PLA

 \therefore the length of PQ = $4\sqrt{6}$ cm

Question 16:



Given: AB is a chord of a circle with centre O.AB is produced to C such that BC = OB.Also, CO is joined to meet the circle in $D.\angle ACD = y^{\circ}$ and $\angle AOD = x^{\circ}$.



To Prove: x = 3y

Proof: OB=BC [Given]

∴ ∠BOC=∠BCO=y° [isosceles triangle]

Ext. \angle OBA = \angle BOC + \angle BCO = (2y)°


Again, OA = OB [radii of same circle] \therefore $\angle OAB = \angle OBA = (2\gamma)^{\circ}$ [isosceles triangle]

Ext. $\angle AOD = \angle OAC + \angle ACO$ = $\angle OAB + \angle BCO = 3y^{\circ}$

 $\therefore \qquad \qquad \mathsf{X}^{\circ} = \mathsf{3}\mathsf{y}^{\circ} \qquad \qquad [\because \angle \mathsf{AOD} = \mathsf{X} \; (\mathsf{given})]$

Question 17:

Given: AB and AC are chords of the circle with centre O. AB = AC, $OP \perp AB$ and $OQ \perp AC$.



To Prove: PB= QC

Proof: AB = AC [Given]

 $\therefore \frac{1}{2}AB = \frac{1}{2}AC \qquad [Divide by 2]$

The perpendicular from the centre of a circle to a chord bisects the chord.

⇒ MB =NC....(1)

Equal chords of a circle are equidistant from the centre.

→ OM = ON

Also, OP=OQ [Radii]

 $\Rightarrow OP - OM = OQ - ON$ $\Rightarrow PM = QN.....(2)$

Now consider the triangles, Δ MPB and Δ NQC:

MB = NC [from (1)]

∠PMB=∠QNC [right angle, given]

PM=QN [from (2)]

Thus, by Side-Angle-Side criterion of congruence, we have

∴ ΔMPB≅ΔNQC [S.A.S]

The corresponding parts of the congruent triangles are equal. $% \label{eq:congruent} % \label{eq:corresponding} % \label{eq:cor$

∴ PB = QC [by c.p.c.t]

Question 18:



Given: BC is a diameter of a circle with centre 0.AB and CD are two chords such that AB \parallel CD.

To Prove: AB = CD

Construction: Draw OL LAB and OM LCD.

Proof: In △ OLB and △OMC


 \angle OLB = \angle OMC [Perpendicular bisector, angle = 90°] \angle OBL = \angle OCD [AB | CD,BC is a transversal, thus alternate interior angles are equal]

OB=OC [Radii]

Thus by Angle-Angle-Side criterion of congruence, we have

 \triangle OLB \cong \triangle OMC [By AAS]

The corresponding parts of the congruent triangle are equal.




But the chords equidistant from the centre are equal.

Question 19:

Let ΔABC be an equilateral triangle of side 9 cm.

Let AD be one of its medians.

Then, AD \perp BC and BD= $\frac{1}{2}\times$ BC = $\left(\frac{1}{2}\times 9\right)$ cm = 4.5 cm.



∴ In right angled △ADB,

$$AB^{2} = AD^{2} + BD^{2}$$

$$\Rightarrow AD^{2} = AB^{2} - BD^{2}$$

$$\Rightarrow AD = \sqrt{AB^{2} - BD^{2}}$$

$$= \sqrt{(9)^{2} - \left(\frac{9}{2}\right)^{2}} \text{ cm } = \frac{9\sqrt{3}}{2} \text{ cm}$$

In an equilateral triangle, the centroid and discumcentre coincide and AG: ${\rm GD}=2:1$

$$\therefore \qquad \text{radius AG} = \frac{2}{3} \text{AD}$$

$$= \left(\frac{2}{3} \times \frac{9\sqrt{3}}{2}\right) \text{cm} = 3\sqrt{3} \text{ cm}$$

 \therefore The radius of the circle is $3\sqrt{3}$ cm.

Question 20:





Given: AB and AC are two equal chords of a circle with

centre O

To Prove: ZOAB = ZOAC
Construction: Join OA, OB and OC.



Proof:In ∆OAB and ∆OAC,

AB = AC [Given]

OA = OA [common]

OB = OC [Radii]

Thus by Side-Side-Side criterion of congruence, we have

∴ ΔOAB≅OAC [by SSS]


The corresponding parts of the congruent triangles are equal.

⇒ ∠OAB=∠OAC [by C.P.C.T.] Therefore, O lies on the bisector of ∠BAC

Question 21:

Given: OPQR is a square. A circle with centre O cuts the

square in X and Y.
To Prove: QX=QY



Construction: Join OX and OY.

Proof: In Δ OXP and Δ OYR

ZOPX = ZORY [Each equal to 90°]

OX = OY [Radii]

OP = OR [Sides of a square]

Thus by Right Angle-Hypotenuse-Side criterion of congruence, we have,

 \triangle OXP \cong \triangle OYR [by RHS]

The corresponding parts of the congruent triangles are equal.

 \Rightarrow PX = RY [by c.p.c.T.]

 \Rightarrow $\Gamma Q - \Gamma X = Q N - N$ [, $\Gamma Q = Q N$

QX = QY.

Exercise 11B

Question 1:

(i) Join BO.

In $\triangle BOC$ we have


OC = OB [Each equal to theradius]

→ ∠OBC=∠OCB : base angles of an isosceles triangle are equal

∠OBC=30° [∵∠OCB=30°]

Thus, we have,

∠OBC=30°(1)





Now, in ABOA, we have

OB=OC [Each equal to the radius]

→ ZOAB = ZOBA [Triangle are equal]

 \angle OBA = 40° [:: \angle OAB = 40°, given]

Thus, we have,

$$\Rightarrow$$
 =30° + 40° [from (1) and (2)]

The angle subtended by an arc of a circle at the centre is double the angle subtended by the arc at any point on the circumference.

(II)
$$\angle BOC = 360^{\circ} - (\angle AOB + \angle AOC)$$

= $360^{\circ} - (90^{\circ} + 110^{\circ})$
= $360^{\circ} - 200^{\circ} = 160^{\circ}$

We know that ∠BOC= 2∠BAC



$$\Rightarrow$$
 $\angle BAC = \frac{160^{\circ}}{2} = 80^{\circ}$

Question 2:

(i)

The angle subtended by an arc of a circle at the centre is double the angle subtended by the arc at any point on the circumference.

$$\Rightarrow$$
 $\angle OCA = \frac{70}{2} = 35^{\circ}$



(ii) The radius of the circle is

OA = OC


⇒ ∠OAC = ∠OCA [base angles of an

isosceles triangle are equal

⇒ ∠OAC = 35° [as∠OCA = 35°]


Question 3:





It is clear that $\angle ACB = \angle PCB$ Consider the triangle $\triangle PCB$. Applying the angle sum property, we have, $\angle PCB = 180^{\circ} - (\angle BPC + \angle PBC)$ $= 180^{\circ} - (180^{\circ} - 110^{\circ} + 25^{\circ})$ [$\angle APB$ and $\angle BPC$ are linear pair; $\angle PBC = 25^{\circ}$, given] $= 180^{\circ} - (70^{\circ} + 25^{\circ})$ $\angle PCB = 180^{\circ} - 95^{\circ} = 85^{\circ}$ Angles in the same segment of a circle are equal. $\therefore \angle ADB = \angle ACB = 85^{\circ}$

Question 4:



It is clear that, BD is the diameter of the circle.

Also we know that, the angle in a semicircle is a right angle.

∴ ∠BAD= 90°

Now consider the triangle, $\triangle BAD$

⇒ ∠ADB = 180° - (∠BAD + ∠ABD) [Angle sum property]

 \Rightarrow = 180° - (90° + 35°) [∠BAD = 90° and ∠ABD = 35°]

⇒ = 180° −125°

⇒ ∠ADB = 55°


Angles in the same segment of a circle are equal.

:. ZACB=ZADB=55°

∴ ∠ACB=55°

Question 5:





The angle subtended by an arc of a circle at the centre is double the angle subtended by the arc at any point on the circumference.

=2×50° [Given] ∠AOB=100°(1)

Consider the triangle △OAB

OA = OB [radius of the circle] $\angle OAB = \angle OBA$ [base angles of an

isosceles triangle are equal]

Thus we have

∠OAB = ∠OBA(2)

By angle sum property, we have

Now ZAOB + ZOAB + ZOBA = 180°


⇒ 100° + 2∠OAB =180° [from (1) and (2)]

⇒ 2∠OAB = 180° - 100° = 80°

 \Rightarrow $\angle OAB = \frac{80^{\circ}}{2} = 40^{\circ}$

∴ ∠OAB = 40°

Question 6:



(i) Angles in the same segment of a circle are equal. ∠ABD and ∠ACD are in the segment AD.

ZACD=ZABD

= 54° [Given]

(ii) Angles in the same segment of a circle are equal.

∠BAD and ∠BCD are in the segment BD.

∠BAD= ∠BCD

= 43° [Given]

(iii) Consider the △ABD.

By Angle sum property we have

∠BAD+∠ADB+∠DBA=180°

⇒ 43° +∠ADB + 54° = 180°

⇒ ∠ADB = 180° - 97° = 83°

⇒ ∠BDA = 83°

Question 7:





Angles in the same segment of a circle are equal.

∠CAD and ∠CBD are in the segment CD.

We know that an angle in a semi circle is a right angle.

$$\angle ACD = 180^{\circ} - (\angle ADC + \angle CAD)$$

= $180^{\circ} - (90^{\circ} + 60^{\circ})$

AC || DE and CD is a transversal, thus alternate angles are equal

Question 8:



Join CO and DO, \angle BCD= \angle ABC=25° [alternate interior angles] The angle subtended by an arc of a circle at the centre

is double the angle subtended by the arc at any point on the circumference.

Similarly,


AB is a straight line passing through the centre

$$\Rightarrow$$
 $\angle COD = 180^{\circ} - 100^{\circ} = 80^{\circ}$

$$\angle CED = \frac{1}{2} \angle COD$$

$$=\frac{80^{\circ}}{2}=40^{\circ}$$


Question 9:



$$= 180^{\circ} - (100^{\circ} + 50^{\circ})$$
 [from (1) and (2)]

Question 10:





The angle subtended by an arc of a circle at the centre is double the angle subtended by the arc at any point on the circumference.

∴
$$\angle AOB = 2\angle ACB$$

⇒ $= 2\angle DCB \ [\because \angle ACB = \angle DCB]$
⇒ $\angle DCB = \frac{1}{2}\angle AOB$
 $= \left(\frac{1}{2} \times 40\right) = 20^{\circ}$

Consider the △DBC;

Byangle sum property, we have

Question 11:

Join OB.

Now in △OAB, we have

$$\Rightarrow \angle OAB + \angle OBA + \angle AOB = 180^{\circ}$$

$$\Rightarrow 25^{\circ} + 25^{\circ} + \angle AOB = 180^{\circ}$$

$$\Rightarrow \angle AOB = 180^{\circ} - 50^{\circ} = 130^{\circ}$$

The angle subtended by an arc of a circle at the centre is double the angle subtended by the arc at any point on the circumference.



Consider the right triangle $\triangle BEC$.

We know that the sum of three angles in a triangle is 180°.

Question 12:




OB = OC[Radius] ∠OBC=∠OCB=55° [base angles in an isosceles triangle are equal]

Consider the triangle ABOC. By angle sum property, we have

$$\angle BOC = 180^{\circ} - (\angle OCB + \angle OBC)$$

= $180^{\circ} - (55^{\circ} + 55^{\circ})$
= $180^{\circ} - 110^{\circ} = 70^{\circ}$
 $\angle BOC = 70^{\circ}$



Again, OA = OB

Consider the triangle $\triangle AOB$.

By angle sum property, we have

Question 13:



Join OB and OC.

The angle subtended by an arc of a circle at the centre is double the angle subtended by the arc at any point on the circumference.

$$\angle BOC = 2\angle BAC$$

$$= 2 \times 30^{\circ} \qquad \left[\because \angle BAC = 30^{\circ}\right]$$

$$= 60^{\circ} \qquad \dots \dots \dots (1)$$

Now consider the triangle ABOC.

[base angles in an isosceles triangle] are equal

Now, in ABOC, we have

$$\Rightarrow \qquad \angle OCB = \frac{120^{\circ}}{2} = 60^{\circ}$$

$$\Rightarrow \qquad \angle OBC = 60^{\circ} \quad [from (2)]$$

Thus, we have, $\angle OBC = \angle OCB = \angle BOC = 60^{\circ}$ So, ΔBOC is an equilateral triangle

OB = OC = BC

.. BC is the radius of the circumference.

Question 14:



```
Consider the triangle, \trianglePRQ.

PQ is the diameter.

The angle in a semicircle is a right angle.

\Rightarrow \anglePRQ = 90°

By the angle sum property in \trianglePRQ, we have,

\angleQPR + \anglePRQ + \anglePQR = 180°

\Rightarrow \angleQPR + 90° + 65° = 180°

\Rightarrow \angleQPR = 180° - 155° = 25° ......(1)
```



Now consider the triangle \triangle PQM. Since PQ is the diameter, \angle PMQ = 90° Again applying the angle sum property in \triangle PQM, we have \angle QPM + \angle PMQ + \angle PQM = 180° \Rightarrow \angle QPM + 90° + 50° = 180° \Rightarrow \angle QPM = 180° - 140° = 40° Now in quadrilateral PQRS \angle QPS + \angle SRQ = 180° \Rightarrow \angle QPR + \angle RPS + \angle PRQ + \angle PRS = 180° [from (1)] \Rightarrow 25° + 40° + 90° + \angle PRS = 180° \Rightarrow \angle PRS = 25° \Rightarrow \angle PRS = 25°

Exercise 110

Question 1:

```
∠BDC = ∠BAC = 40° [angles in the same segment]


In∆BCD, we have

∠BCD + ∠BDC + ∠DBC = 180^{\circ}

∴ ∠BCD + 40^{\circ} + 60^{\circ} = 180^{\circ}

⇒ ∠BCD = 180^{\circ} - 100^{\circ} = 80^{\circ}

∴ ∠BCD = 80^{\circ}
```



(ii) Also $\angle CAD = \angle CBD$ [angles in the same segment] $\angle CAD = 60^{\circ}$ [$\therefore \angle CBD = 60^{\circ}$]

Question 2:



In cyclic quadrilateral PQRS

$$\angle$$
PSR + \angle PQR = 180°
 \Rightarrow 150° + \angle PQR = 180°
 \Rightarrow \angle PQR = 180° - 150° = 30°(i)
Also, \angle PRQ = 90°(ii)
[angle ina semi circle]



Now in $\triangle PRQ$ we have

$$\Rightarrow$$
 30° + 90° + \angle RPQ = 180° [from (i)and(ii)]

$$\Rightarrow \qquad \angle RPQ = 180^{\circ} - 120^{\circ} = 60^{\circ}$$

Question 3:

In cyclic quadrilateral ABCD, AB | DC and \(\text{BAD} = 100\)



Question 4:


Take a point D on the major arc CA and join AD and DC

Angle subtended by an arc is twice the angle subtended by it on the circumference in the alternate segment.

$$\angle PBC = \angle$$

[. exterior angle of a cyclic quadrilateral interior opposite angle]

Question 5:








ABCD is a cyclic quadrilateral

∴ ∠ABC + ∠ADC = 180° ⇒ 92° + ∠ADC = 180°

⇒ ∠ADC =180° - 92° = 88°



Also, AE | CD

∴ ∠EAD = ∠ADC = 88°
 ∴ ∠BCD = ∠DAF

[: exterior angle of a cyclic quadrilateral =int.opp.angle]

∴ ∠BCD = ∠EAD + ∠EAF

 $=88^{\circ} + 20^{\circ}$ [:: \angle FAE = 20° (given)]

=108°

∴ ∠BCD = 108°

Question 6:

BD = DC

∴ ∠BCD =∠CBD=30°



In △BCD, we have

 $\angle BCD + \angle CBD + \angle CDB = 180^{\circ}$ $\Rightarrow 30^{\circ} + 30^{\circ} + \angle CDB = 180^{\circ}$ $\Rightarrow \angle CDB = 180^{\circ} - 60^{\circ}$ $= 120^{\circ}$

The opposite angles of a cyclic quadrilateral are supplementary.

ABCD is a cyclic quadrilateral and thus, ∠CDB + ∠BAC =180°

= $180^{\circ} - 120^{\circ} [\because \angle CDB = 120^{\circ}]$ = 60°

∠BAC=60°

Question 7:



Angle subtended by an arc is twice the angle subtended by it on the circumference in the alternate segment.

Here arc ABC makes ∠AOC =100° at the centre of the circle and ZADC on the circumference of the circle

⇒
$$\angle ADC = \frac{1}{2}(\angle AOC)$$

$$\Rightarrow = \frac{1}{2} \times 100^{\circ} \left[\angle AOC = 100^{\circ} \right]$$



The opposite angles of a cyclic quadrilateral are supplementary ABCD is a cyclic quadrilateral and thus,

$$\angle ADC + \angle ABC = 180^{\circ}$$

= $180^{\circ} - 50^{\circ} [\because \angle ADC = 50^{\circ}]$

Question 8:

- Δ ABC is an equilateral triangle.
- ... Each of its angle is equal to 60°
- ⇒ ∠BAC = ∠ABC = ∠ACB = 60°



(i) Angle s in the same segment of a circle are equal.

(ii) The opposite angles of a cyclic quadrilateral are supplementary ABCE is a cyclic quadrilateral and thus,

Question 9:

ABCD is a cyclic quadrilateral.

Million State & Practice opp.angle of a cyclic quadrilateral are supplementary

$$\Rightarrow$$
 $\angle A + 100^{\circ} = 180^{\circ}$



Now in AABD, we have

Question 10:



O is the centre of the circle and ∠BOD = 150° Reflex $\angle BOD = (360^{\circ} - \angle BOD)$ $=(360^{\circ}-150^{\circ})=210^{\circ}$



Now,
$$x = \frac{1}{2} (\text{reflex} \angle BOD)$$

 $= \frac{1}{2} \times 210^{\circ} = 105^{\circ}$
 $\therefore \qquad x = 105^{\circ}$
Again, $x + y = 180^{\circ}$
 $\Rightarrow \qquad 105^{\circ} + y = 180^{\circ}$
 $\Rightarrow \qquad y = 180^{\circ} - 105^{\circ} = 75^{\circ}$
 $\therefore \qquad y = 75^{\circ}$

Question 11:

O is the centre of the circle and $\angle DAB = 50^{\circ}$

In △OAB we have

$$\angle$$
OAB + \angle OBA + \angle AOB = 180°
 \Rightarrow 50° + 50° + \angle AOB = 180°
 \Rightarrow \angle AOB = 180° - 100° = 80°

Since, AOD is a straight line, x=180° - ∠AOB.

The opposite angles of a cyclic quadrilateral are supplementary.

ABCD is a cyclic quadrilateral and thus,

$$\angle DAB + \angle BCD = 180^{\circ}$$

 $\angle BCD = 180^{\circ} - 50^{\circ} [\because \angle DAB = 50^{\circ}, given]$
 $= 130^{\circ}$
 $\Rightarrow \qquad y = 130^{\circ}$
Thus, $\times = 100^{\circ}$ and $y = 130^{\circ}$

Question 12:

ABCD is a cyclic quadrilateral.


We know that in a cyclic quadrilateral exterior angle = interior opposite angle.

$$\angle CBF = \angle CDA = (180^{\circ} - \times)$$

$$\Rightarrow 130^{\circ} = 180^{\circ} - \times$$

$$\Rightarrow \times = 180^{\circ} - 130^{\circ} = 50^{\circ}$$

$$\times = 50^{\circ}$$



Question 13:




AB is a diameter of a circle with centre O and DO || CB, ∠BCD = 120°

(i) Since ABCD is a cyclic quadrilateral

∴ ∠BCD + ∠BAD = 180°⇒ 120° + ∠BAD = 180°

⇒ ∠BAD = 180° - 120° = 60°



∠BDA + ∠BAD + ∠ABD = 180°

(iii) OD = OA.

∠ODA = ∠OAD = ∠BAD = 60°

$$\angle ODB = 90^{\circ} - \angle ODA$$

= $90^{\circ} - 60^{\circ} = 30^{\circ}$

Since DO | CB, alternate angles are equal

⇒ ∠CBD = ∠ODB

$$= 90^{\circ} + 30^{\circ} = 120^{\circ}$$

Also, in AAOD, we have

Since all the angles of AAOD are of 60° each


∴ △ AOD is an equilateral triangle.

Question 14:

AB and CD are two chords of a circle which interect each other at P, outside the circle. AB = 6cm, BP = 2 cm and PD = 2.5 cm

Therefore, $AP \times BP = CP \times DP$

Or,
$$8 \times 2 = (CD + 2.5) \times 2.5 \text{ cm}$$
 [as $CP = CD + DP$]



Thus,
$$8 \times 2 = (x + 2.5) \times 2.5$$

⇒ $16 \text{ cm} = 2.5 \times + 6.25 \text{ cm}$
⇒ $2.5 \times = (16 - 6.25) \text{ cm}$
⇒ $2.5 \times = 9.75 \text{ cm}$
⇒ $\times = \frac{9.75}{2.5} = 3.9 \text{ cm}$
∴ $\times = 3.9 \text{ cm}$

Therefore, CD = 3.9 cm

Question 15:



```
O is the centre of a circle having \angle AOD = 140^\circ and \angle CAB = 50^\circ

(i) \angle BOD = 180^\circ - \angle AOD

= 180^\circ - 140^\circ = 40^\circ

OB = OD


\angle OBD = \angle ODB
```

```
In ∆OBD, we have
  ∠BOD + ∠OBD + ∠ODB = 180°
 ⇒ ∠BOD + ∠OBD + ∠OBD = 180°
                                           [∵∠OBD = ∠ODB]
               40° + 2∠OBD = 180°
                                           [:: ∠BOD = 40°]
                     2∠OBD =180° - 40° =140°
                       \angleOBD = \angleODB = \frac{140}{2} = 70°
                ∠CAB + ∠BDC = 180°
                                            [: ABCD is cyclic]
 Also.
         ∠CAB + ∠ODB + ∠ODC = 180°
          50° + 70° + ∠ODC = 180°
\Rightarrow
                       ∠ODC = 180° - 120° = 60°
                       ∠ODC = 60°
                        \angle EDB = 180^{\circ} - (\angle ODC + \angle ODB)
                              =180^{\circ} - (60^{\circ} + 70^{\circ})
                               = 180° -130° = 50°
  (ii)
                  ∠EBD = 180° - ∠OBD
                         = 180^{\circ} - 70^{\circ} = 110^{\circ}
```

Question 16:

Consider the triangles, ΔEBC and ΔEDA
Side AB of the cyclic quadrilateral ABCD is produced to E

∴ ∠EBC = ∠CDA



Again, side DC of the cyclic quadrilateral ABCD isproduced

....(ii)

∴ ∠ECB=∠BAD ⇒ ∠ECB=∠EAD


and $\angle BEC = \angle DEA$ [each equal to $\angle E$]....(iii)

Thus from (i), (ii) and (iii), we have $\triangle EBC \cong \triangle EDA$

Question 17:

 Δ ABC is an isosceles triangle in which AB = AC and a drde passing through B and C intersects AB and AC at D and E.


Since AB = AC
∴ ∠ACB = ∠ABC
So, ext. ∠ADE = ∠ACB = ∠ABC
∴ ∠ADE = ∠ABC
⇒ DE || BC.



Question 18:



 \triangle ABC is an isosceles trianglein which AB = AC. D and E are the mid points of AB and AC respectively.



.*.	DE BC	
\Rightarrow	$\angle ADE = \angle ABC$	(i)
Also,	AB = AC	[Given]
\Rightarrow	$\angle ABC = \angle ACB$	(ii)
<i>:</i> .	$\angle ADE = \angle ACB$	[From (i) and(ii)]

Now, $\angle ADE + \angle EDB = 180^{\circ}$ [: ADBis a straightline]

∴ ∠ACB + ∠EDB = 180°

- ⇒ The opposite angles are supplementary.
- ⇒ D,B,C and E are concyclic i.e. D,B,C and E is a cyclic quadrilateral.

Question 19:

Let ABCD be a cyclic quadrilateral and let O be the centre of the circle passing through A, B, C, D.

Then each of AB,BC,CD and DA being a chord of the circle, its right bisector must pass through O.

∴ the right bisectors of AB,BC,CD andDA pass through and are concurrent.



Question 20:


ABCD is a rhombus.

Let the diagonals AC and BD of the rhombus ABCD intersect at O.

But, we know, that the diagonals of a rhombus bisect each other at right angles.

So,∠BOC = 90°

.. ∠BOC lies in a drde.



Thus the circle drawn with BC as diameter will pass through O

Similarly, all the circles described with AB, AD and CD as diameters will pass through O.

Question 21:





ABCD is a rectangle.

Let O be the point of intersection of the diagonals AC and BD of rectangle ABCD.



Since the diagonals of a rectangle are equal and bisecteach other.

: OA = OB = OC = OD

Thus, O is the centre of the circle through A, B, C, D.


Question 22:

Let A, B, C be the given points.

With B as centre and radius equal to AC draw an arc.

With C as centre and AB as radius draw another arc,

which cuts the previous arcat D.



Then D is the required point BD and CD.

In ∆ABC and ∆DCB

AB = DC

AC = DB

BC = CB [common]

ABC≅ADCB [by SSS]

⇒ ∠BAC = ∠CDB [CPCIT]

Thus, BC subtends equal angles, ∠BAC and ∠CDB on the same side of it.

.. Point's A,B,C,D are concyclic.

Question 23:

ABCD is a cydic quadrilateral

$$\angle B - \angle D = 60^{\circ}$$
(i)

and $\angle B + \angle D = 180^{\circ}$ (ii)

Adding (i) and (ii) we get,

$$\angle B = \frac{240}{2} = 120^{\circ}$$

Substituting the value of $\angle B = 120^{\circ}$ in (i) we get

$$\Rightarrow$$
 $\angle D = 120^{\circ} - 60^{\circ} = 60^{\circ}$

The smaller of the two angles i.e. $\angle D = 60^{\circ}$

Question 24:



ABCD is a quadrilateral in which AD = BC and \angle ADC = \angle BCD Draw DE \perp AB and CF \perp AB



Now, in \triangle ADE and \triangle BCF, we have

$$\angle AED = \angle BFC$$
 [each equal to 90°]
 $\angle ADE = \angle ADC - 90^\circ = \angle BCD - 90^\circ = \angle BCF$
 $AD = BC$ [given]

Thus, by Angle-Angle-Side criterionof congruence, we have $\therefore \qquad \triangle \, \mathsf{ADE} \cong \Delta \, \mathsf{BCF} \qquad \text{[by AAS congruence]}$

The corresponding parts of the congruent triangles are equal.

$$\angle A = \angle B$$
Now, $\angle A + \angle B + \angle C + \angle D = 360^{\circ}$

$$\Rightarrow 2\angle B + 2\angle D = 360^{\circ}$$

$$\Rightarrow 2(\angle B + \angle D) = 360^{\circ}$$

$$\Rightarrow \angle B + \angle D = \frac{360}{2} = 180^{\circ}$$

$$\therefore ABCD \text{ is a cyclic quadrilateral.}$$


Question 25:

If one side of a cyclic quadrilateral is produced then the exterior angle is equal to the interior opposite angle.

$$\Rightarrow \angle BAD = \angle DCF = 75^{\circ}$$

$$\angle DCF = x = 75^{\circ}$$

$$x = 75^{\circ}$$



The opposite angles of the opposite angles of a cyclic quadrilateral is 180°

$$\Rightarrow \qquad \angle DCF + \angle DEF = 180^{\circ}$$

$$\Rightarrow \qquad 75^{\circ} + \angle DEF = 180^{\circ}$$

$$\Rightarrow \qquad \angle DEF = 180^{\circ} - 75^{\circ} = 105^{\circ}$$

$$As \qquad \angle DEF = y^{\circ} = 105^{\circ}$$

$$\therefore \times = 75^{\circ} \text{ and } y = 105^{\circ}$$

Question 26:

Given: Let ABCD be a cyclic quadrilateral whose diagonals AC and BD intersect at O at right angles.

Let OL ⊥ AB such that LO produced meets CD at M.



To Pr ove: CM = MD

Pr oof:
$$\angle 1 = \angle 2$$
 [angles in the same segment]

 $\angle 2 + \angle 3 = 90^{\circ}$ [: $\angle OLB = 90^{\circ}$]

 $\angle 3 + \angle 4 = 90^{\circ}$ [: LOM is a straight line] and $\angle BOC = 90^{\circ}$


segment]



Question 27:

Chord AB of a circle is produced to E.

If one side of a cyclic quadrilateral is produced then the exterior angle is equal to the interior opposite angle.




Chord CD of a circle is produced to E \therefore Ext. \angle DBE = \angle ACD = \angle ACE....(2)

Consider the triangles \triangle EDB and \triangle EAC. \angle BDE = \angle CAE [from(1)] \angle DBE = \angle ACE [from(2)] \angle E = \angle E [common] \triangle EDB \sim \triangle EAC.

Question 28:

Given: AB and CD are two parallel chords of a circle BDE and ACE are straight lines which intersect at E.

If one side of a cyclic quadrilateral is produced then the exterior angle is equal to the interior opposite angle.




Also, AB | CD

 \Rightarrow $\angle EDC = \angle B$ and $\angle DCE = \angle A$ \therefore $\angle A = \angle B$ \therefore \triangle AEB is isosceles.

Question 29:

AB is a diameter of a circle with centre O. ADE and CBE are straight lines, meeting at E, such that ZBAD = 35° and ZBED = 25°.

Join BD and AC.



```
(i) Now,
                   \angle BDA = 90^{\circ} = \angle EDB
                                                         [angle in a semi circle]
                        ∠EBD = 180° - (∠EDB + ∠BED)
     \Rightarrow
                                =180^{\circ} - (90^{\circ} + 25^{\circ})
                                 =180^{\circ} - 115^{\circ} = 65^{\circ}
                       \angle DBC = (180^{\circ} - \angle EBD)
                                =180^{\circ} - 65^{\circ} = 115^{\circ}
                        \angle DBC = 115^{\circ}
(ii) Again,
                   \angle DCB = \angle BAD
                                             [angle in the same segment]
                    ∠BAD = 35°
     Since,
                    ∠DCB = 35°
(iii)
                    \angle BDC = 180^{\circ} - (\angle DBC + \angle DCB)
                             = 180^{\circ} - (\angle DBC + \angle BAD)
                            =180°-(115°+35°)
                            =180^{\circ} - 150^{\circ} = 30^{\circ}
                   ∠BDC =30°
```