

# Chapter 5 Continuity and Differentiability

## EXERCISE 5.1

### Question 1:

Prove that the function  $f(x) = 5x - 3$  is continuous at  $x = 0$ ,  $x = -3$  and at  $x = 5$ .

### Solution:

The given function is  $f(x) = 5x - 3$

$$\text{At } x = 0, f(0) = 5(0) - 3 = -3$$

$$\lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} (5x - 3) = 5(0) - 3 = -3$$

$$\therefore \lim_{x \rightarrow 0} f(x) = f(0)$$

Therefore,  $f$  is continuous at  $x = 0$ .

$$\text{At } x = -3, f(-3) = 5(-3) - 3 = -18$$

$$\lim_{x \rightarrow -3} f(x) = \lim_{x \rightarrow -3} (5x - 3) = 5(-3) - 3 = -18$$

$$\therefore \lim_{x \rightarrow -3} f(x) = f(-3)$$

Therefore,  $f$  is continuous at  $x = -3$ .

$$\text{At } x = 5, f(5) = 5(5) - 3 = 22$$

$$\lim_{x \rightarrow 5} f(x) = \lim_{x \rightarrow 5} (5x - 3) = 5(5) - 3 = 22$$

$$\therefore \lim_{x \rightarrow 5} f(x) = f(5)$$

Therefore,  $f$  is continuous at  $x = 5$ .

### Question 2:

Examine the continuity of the function  $f(x) = 2x^2 - 1$  at  $x = 3$ .

### Solution:

The given function is  $f(x) = 2x^2 - 1$

$$\text{At } x = 3, f(3) = 2(3)^2 - 1 = 17$$

$$\lim_{x \rightarrow 3} f(x) = \lim_{x \rightarrow 3} (2x^2 - 1) = 2(3^2) - 1 = 17$$

$$\therefore \lim_{x \rightarrow 3} f(x) = f(3)$$

Therefore,  $f$  is continuous at  $x = 3$ .

### Question 3:

Examine the following functions for continuity.

(i)  $f(x) = x - 5$

- (ii)  $f(x) = \frac{1}{x-5}, x \neq 5$   
 (iii)  $f(x) = \frac{x^2-25}{x+5}, x \neq -5$   
 (iv)  $f(x) = |x-5|, x \neq 5$

### Solution:

- (i) The given function is  $f(x) = x - 5$

It is evident that  $f$  is defined at every real number  $k$  and its value at  $k$  is  $k - 5$ .

It is also observed that

$$\lim_{x \rightarrow k} f(x) = \lim_{x \rightarrow k} (x - 5) = k - 5 = f(k)$$

$$\therefore \lim_{x \rightarrow k} f(x) = f(k)$$

Hence,  $f$  is continuous at every real number and therefore, it is a continuous function.

- (ii) The given function is  $f(x) = \frac{1}{x-5}, x \neq 5$

For any real number  $k \neq 5$ , we obtain

$$\lim_{x \rightarrow k} f(x) = \lim_{x \rightarrow k} \frac{1}{x-5} = \frac{1}{k-5}$$

Also,

$$f(k) = \frac{1}{k-5} \quad (\text{As } k \neq 5)$$

$$\therefore \lim_{x \rightarrow k} f(x) = f(k)$$

Hence,  $f$  is continuous at every point in the domain of  $f$  and therefore, it is a continuous function.

- (iii) The given function is  $f(x) = \frac{x^2-25}{x+5}, x \neq -5$

For any real number  $c \neq -5$ , we obtain

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} \frac{x^2-25}{x+5} = \lim_{x \rightarrow c} \frac{(x+5)(x-5)}{x+5} = \lim_{x \rightarrow c} (x-5) = (c-5)$$

Also,

$$f(c) = \frac{(c+5)(c-5)}{c+5} = (c-5)$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Hence,  $f$  is continuous at every point in the domain of  $f$  and therefore, it is a continuous function.

(iv) The given function is  $f(x) = |x - 5| = \begin{cases} 5 - x, & \text{if } x < 5 \\ x - 5, & \text{if } x \geq 5 \end{cases}$

This function  $f$  is defined at all points of the real line. Let  $c$  be a point on a real line. Then,  $c < 5$ ,  $c = 5$  or  $c > 5$

Case I:  $c < 5$

Then,  $f(c) = 5 - c$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (5 - x) = 5 - c$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all real numbers less than 5.

Case II:  $c = 5$

Then,  $f(c) = f(5) = (5 - 5) = 0$

$$\lim_{x \rightarrow 5^-} f(x) = \lim_{x \rightarrow 5^-} (5 - x) = (5 - 5) = 0$$

$$\lim_{x \rightarrow 5^+} f(x) = \lim_{x \rightarrow 5^+} (x - 5) = 0$$

$$\therefore \lim_{x \rightarrow c^-} f(x) = \lim_{x \rightarrow c^+} f(x) = f(c)$$

Therefore,  $f$  is continuous at  $x = 5$

Case III:  $c > 5$

Then,  $f(c) = f(5) = c - 5$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (x - 5) = c - 5$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all real numbers greater than 5.

Hence,  $f$  is continuous at every real number and therefore, it is a continuous function.

#### Question 4:

Prove that the function  $f(x) = x^n$  is continuous at  $x = n$ , where  $n$  is a positive integer.

### Solution:

The given function is  $f(x) = x^n$

It is observed that  $f$  is defined at all positive integers,  $n$ , and its value at  $n$  is  $n^n$ .

Then,

$$\lim_{x \rightarrow n} f(n) = \lim_{x \rightarrow n} (x^n) = x^n$$

$$\therefore \lim_{x \rightarrow n} f(x) = f(n)$$

Therefore,  $f$  is continuous at  $n$ , where  $n$  is a positive integer.

### Question 5:

Is the function  $f$  defined by  $f(x) = \begin{cases} x, & \text{if } x \leq 1 \\ 5, & \text{if } x > 1 \end{cases}$  continuous at  $x = 0$ ? At  $x = 1$ ? At  $x = 2$ ?

### Solution:

The given function is  $f(x) = \begin{cases} x, & \text{if } x \leq 1 \\ 5, & \text{if } x > 1 \end{cases}$

At  $x = 0$ ,

It is evident that  $f$  is defined at 0 and its value at 0 is 0.

Then,

$$\lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} (x) = 0$$

$$\therefore \lim_{x \rightarrow 0} f(x) = f(0)$$

Therefore,  $f$  is continuous at  $x = 0$ .

At  $x = 1$ ,

It is evident that  $f$  is defined at 1 and its value at 1 is 1.

The left hand limit of  $f$  at  $x = 1$  is,

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} (x) = 1$$

The right hand limit of  $f$  at  $x = 1$  is,

$$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} (5) = 5$$

$$\therefore \lim_{x \rightarrow 1^-} f(x) \neq \lim_{x \rightarrow 1^+} f(x)$$

Therefore,  $f$  is not continuous at  $x = 1$ .

At  $x = 2$ ,

It is evident that  $f$  is defined at 2 and its value at 2 is 5.

$$\lim_{x \rightarrow 2} f(x) = \lim_{x \rightarrow 2} (5) = 5$$

$$\therefore \lim_{x \rightarrow 2} f(x) = f(2)$$

Therefore,  $f$  is continuous at  $x = 2$ .

### Question 6:

Find all points of discontinuity of  $f$ , where  $f$  is defined by  $f(x) = \begin{cases} 2x+3, & \text{if } x \leq 2 \\ 2x-3, & \text{if } x > 2 \end{cases}$ .

#### Solution:

The given function is  $f(x) = \begin{cases} 2x+3, & \text{if } x \leq 2 \\ 2x-3, & \text{if } x > 2 \end{cases}$

It is evident that the given function  $f$  is defined at all the points of the real line.

Let  $c$  be a point on the real line. Then, three cases arise.

$$c < 2$$

$$c > 2$$

$$c = 2$$

Case I:  $c < 2$

$$f(c) = 2c + 3$$

Then,

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (2x + 3) = 2c + 3$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x < 2$ .

Case II:  $c > 2$

Then,

$$f(c) = 2c - 3$$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (2x - 3) = 2c - 3$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x > 2$ .

Case III:  $c = 2$

Then, the left hand limit of  $f$  at  $x = 2$  is,

$$\lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^-} (2x + 3) = 2(2) + 3 = 7$$

The right hand limit of  $f$  at  $x = 2$  is,

$$\lim_{x \rightarrow 2^+} f(x) = \lim_{x \rightarrow 2^+} (2x - 3) = 2(2) - 3 = 1$$

It is observed that the left and right hand limit of  $f$  at  $x = 2$  do not coincide.

Therefore,  $f$  is not continuous at  $x = 2$ .

Hence,  $x = 2$  is the only point of discontinuity of  $f$ .

### Question 7:

$$f(x) = \begin{cases} |x| + 3, & \text{if } x \leq -3 \\ -2x, & \text{if } -3 < x < 3 \\ 6x + 2, & \text{if } x \geq 3 \end{cases}$$

Find all points of discontinuity of  $f$ , where  $f$  is defined by

### Solution:

$$f(x) = \begin{cases} |x| + 3, & \text{if } x \leq -3 \\ -2x, & \text{if } -3 < x < 3 \\ 6x + 2, & \text{if } x \geq 3 \end{cases}$$

The given function is

The given function  $f$  is defined at all the points of the real line.

Let  $c$  be a point on the real line.

Case I:

If  $c < -3$ , then  $f(c) = -c + 3$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (-x + 3) = -c + 3$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x < -3$ .

Case II:

If  $c = -3$ , then  $f(-3) = -(-3) + 3 = 6$

$$\lim_{x \rightarrow -3^-} f(x) = \lim_{x \rightarrow -3^-} (-x + 3) = -(-3) + 3 = 6$$

$$\lim_{x \rightarrow -3^+} f(x) = \lim_{x \rightarrow -3^+} (-2x) = -2(-3) = 6$$

$$\therefore \lim_{x \rightarrow -3} f(x) = f(-3)$$

Therefore,  $f$  is continuous at  $x = -3$ .

Case III:

If  $-3 < c < 3$ , then  $f(c) = -2c$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (-2x) = -2c$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous in  $(-3, 3)$ .

Case IV:

If  $c = 3$ , then the left hand limit of  $f$  at  $x = 3$  is,

$$\lim_{x \rightarrow 3^-} f(x) = \lim_{x \rightarrow 3^-} (-2x) = -2(3) = -6$$

The right hand limit of  $f$  at  $x = 3$  is,

$$\lim_{x \rightarrow 3^+} f(x) = \lim_{x \rightarrow 3^+} (6x + 2) = 6(3) + 2 = 20$$

It is observed that the left and right hand limit of  $f$  at  $x = 3$  do not coincide.

Therefore,  $f$  is not continuous at  $x = 3$ .

Case V:

If  $c > 3$ , then  $f(c) = 6c + 2$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (6x + 2) = 6c + 2$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x > 3$ .

Hence,  $x = 3$  is the only point of discontinuity of  $f$ .

**Question 8:**

Find all points of discontinuity of  $f$ , where  $f$  is defined by

$$f(x) = \begin{cases} \frac{|x|}{x}, & \text{if } x \neq 0 \\ 0, & \text{if } x = 0 \end{cases}$$

**Solution:**

$$f(x) = \begin{cases} \frac{|x|}{x}, & \text{if } x \neq 0 \\ 0, & \text{if } x = 0 \end{cases}$$

The given function is

It is known that,  $x < 0 \Rightarrow |x| = -x$  and  $x > 0 \Rightarrow |x| = x$

Therefore, the given function can be rewritten as

$$f(x) = \begin{cases} \frac{|x|}{x} = \frac{-x}{x} = -1, & \text{if } x < 0 \\ 0, & \text{if } x = 0 \\ \frac{|x|}{x} = \frac{x}{x} = 1, & \text{if } x > 0 \end{cases}$$

The given function  $f$  is defined at all the points of the real line.

Let  $c$  be a point on the real line.

Case I:

If  $c < 0$ , then  $f(c) = -1$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (-1) = -1$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x < 0$ .

Case II:

If  $c = 0$ , then the left hand limit of  $f$  at  $x = 0$  is,

$$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} (-1) = -1$$

The right hand limit of  $f$  at  $x = 0$  is,

$$\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} (1) = 1$$

It is observed that the left and right hand limit of  $f$  at  $x = 0$  do not coincide.

Therefore,  $f$  is not continuous at  $x = 0$ .



Case III:

If  $c > 0$ , then  $f(c) = 1$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (1) = 1$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x > 0$ .

Hence,  $x = 0$  is the only point of discontinuity of  $f$ .

### Question 9:

$$f(x) = \begin{cases} \frac{x}{|x|}, & \text{if } x < 0 \\ -1, & \text{if } x \geq 0 \end{cases}$$

Find all points of discontinuity of  $f$ , where  $f$  is defined by

**Solution:**

$$f(x) = \begin{cases} \frac{x}{|x|}, & \text{if } x < 0 \\ -1, & \text{if } x \geq 0 \end{cases}$$

The given function is

It is known that  $x < 0 \Rightarrow |x| = -x$

Therefore, the given function can be rewritten as

$$f(x) = \begin{cases} \frac{x}{|x|} = \frac{x}{-x} = -1, & \text{if } x < 0 \\ -1, & \text{if } x \geq 0 \end{cases}$$

$$\Rightarrow f(x) = -1 \forall x \in \mathbb{R}$$

Let  $c$  be any real number.

$$\text{Then, } \lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (-1) = -1$$

$$\text{Also, } f(c) = -1 = \lim_{x \rightarrow c} f(x)$$

Therefore, the given function is a continuous function.

Hence, the given function has no point of discontinuity.

### Question 10:

Find all points of discontinuity of  $f$ , where  $f$  is defined by

$$f(x) = \begin{cases} x+1, & \text{if } x \geq 1 \\ x^2+1, & \text{if } x < 1 \end{cases}$$

**Solution:**

The given function is  $f(x) = \begin{cases} x+1, & \text{if } x \geq 1 \\ x^2+1, & \text{if } x < 1 \end{cases}$

The given function  $f$  is defined at all the points of the real line.

Let  $c$  be a point on the real line.

Case I:

If  $c < 1$ , then  $f(c) = c^2 + 1$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (x^2 + 1) = c^2 + 1$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x < 1$ .

Case II:

If  $c = 1$ , then  $f(c) = f(1) = 1 + 1 = 2$

The left hand limit of  $f$  at  $x = 1$  is,

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} (x^2 + 1) = 1^2 + 1 = 2$$

The right hand limit of  $f$  at  $x = 1$  is,

$$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} (x + 1) = 1 + 1 = 2$$

$$\therefore \lim_{x \rightarrow 1} f(x) = f(1)$$

Therefore,  $f$  is continuous at  $x = 1$ .

Case III:

If  $c > 1$ , then  $f(c) = c + 1$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (x + 1) = c + 1$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x > 1$ .

Hence, the given function  $f$  has no point of discontinuity.

**Question 11:**

Find all points of discontinuity of  $f$ , where  $f$  is defined by  $f(x) = \begin{cases} x^3 - 3, & \text{if } x \leq 2 \\ x^2 + 1, & \text{if } x > 2 \end{cases}$

**Solution:**

The given function is  $f(x) = \begin{cases} x^3 - 3, & \text{if } x \leq 2 \\ x^2 + 1, & \text{if } x > 2 \end{cases}$

The given function  $f$  is defined at all the points of the real line.

Let  $c$  be a point on the real line.

Case I:

If  $c < 2$ , then  $f(c) = c^3 - 3$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (x^3 - 3) = c^3 - 3$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x < 2$ .

Case II:

If  $c = 2$ , then  $f(c) = f(2) = 2^3 - 3 = 5$

$$\lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^-} (x^3 - 3) = 2^3 - 3 = 5$$

$$\lim_{x \rightarrow 2^+} f(x) = \lim_{x \rightarrow 2^+} (x^2 + 1) = 2^2 + 1 = 5$$

$$\therefore \lim_{x \rightarrow 2} f(x) = f(2)$$

Therefore,  $f$  is continuous at  $x = 2$ .

Case III:

If  $c > 2$ , then  $f(c) = c^2 + 1$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (x^2 + 1) = c^2 + 1$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x > 2$ .

Thus, the given function  $f$  is continuous at every point on the real line.

Hence,  $f$  has no point of discontinuity.

### Question 12:

Find all points of discontinuity of  $f$ , where  $f$  is defined by  $f(x) = \begin{cases} x^{10} - 1, & \text{if } x \leq 1 \\ x^2, & \text{if } x > 1 \end{cases}$ .

**Solution:**

The given function is  $f(x) = \begin{cases} x^{10} - 1, & \text{if } x \leq 1 \\ x^2, & \text{if } x > 1 \end{cases}$

The given function  $f$  is defined at all the points of the real line.

Let  $c$  be a point on the real line.

Case I:

If  $c < 1$ , then  $f(c) = c^{10} - 1$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (x^{10} - 1) = c^{10} - 1$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x < 1$ .

Case II:

If  $c = 1$ , then the left hand limit of  $f$  at  $x = 1$  is,

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} (x^{10} - 1) = 1^{10} - 1 = 1 - 1 = 0$$

The right hand limit of  $f$  at  $x = 1$  is,

$$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} (x^2) = 1^2 = 1$$

It is observed that the left and right hand limit of  $f$  at  $x = 1$  do not coincide.

Therefore,  $f$  is not continuous at  $x = 1$ .

Case III:

If  $c > 1$ , then  $f(c) = c^2$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (x^2) = c^2$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x > 1$ .

Thus from the above observation, it can be concluded that  $x = 1$  is the only point of discontinuity of  $f$ .

### Question 13:

Is the function defined by  $f(x) = \begin{cases} x+5, & \text{if } x \leq 1 \\ x-5, & \text{if } x > 1 \end{cases}$  a continuous function?

**Solution:**

The given function is  $f(x) = \begin{cases} x+5, & \text{if } x \leq 1 \\ x-5, & \text{if } x > 1 \end{cases}$

The given function  $f$  is defined at all the points of the real line.

Let  $c$  be a point on the real line.

Case I:

If  $c < 1$ , then  $f(c) = c + 5$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (x + 5) = c + 5$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x < 1$ .

Case II:

If  $c = 1$ , then  $f(1) = 1 + 5 = 6$

The left hand limit of  $f$  at  $x = 1$  is,

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} (x + 5) = 1 + 5 = 6$$

The right hand limit of  $f$  at  $x = 1$  is,

$$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} (x - 5) = 1 - 5 = -4$$

It is observed that the left and right hand limit of  $f$  at  $x = 1$  do not coincide.

Therefore,  $f$  is not continuous at  $x = 1$ .

Case III:

If  $c > 1$ , then  $f(c) = c - 5$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (x - 5) = c - 5$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x > 1$ .

From the above observation it can be concluded that,  $x = 1$  is the only point of discontinuity of  $f$ .

#### Question 14:

$$f(x) = \begin{cases} 3, & \text{if } 0 \leq x \leq 1 \\ 4, & \text{if } 1 < x < 3 \\ 5, & \text{if } 3 \leq x \leq 10 \end{cases}$$

Discuss the continuity of the function  $f$ , where  $f$  is defined by

**Solution:**

$$f(x) = \begin{cases} 3, & \text{if } 0 \leq x \leq 1 \\ 4, & \text{if } 1 < x < 3 \\ 5, & \text{if } 3 \leq x \leq 10 \end{cases}$$

The given function is

The given function  $f$  is defined at all the points of the interval  $[0, 10]$ .

Let  $c$  be a point in the interval  $[0, 10]$ .

Case I:

If  $0 \leq c < 1$ , then  $f(c) = 3$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (3) = 3$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous in the interval  $[0, 1)$ .

Case II:

If  $c = 1$ , then  $f(3) = 3$

The left hand limit of  $f$  at  $x = 1$  is,

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} (3) = 3$$

The right hand limit of  $f$  at  $x = 1$  is,

$$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} (4) = 4$$

It is observed that the left and right hand limit of  $f$  at  $x = 1$  do not coincide.

Therefore,  $f$  is not continuous at  $x = 1$ .

Case III:

If  $1 < c < 3$ , then  $f(c) = 4$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (4) = 4$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at in the interval  $(1, 3)$ .

Case IV:

If  $c = 3$ , then  $f(c) = 5$

The left hand limit of  $f$  at  $x = 3$  is,

$$\lim_{x \rightarrow 3^-} f(x) = \lim_{x \rightarrow 3^-} (4) = 4$$

The right hand limit of  $f$  at  $x = 3$  is,

$$\lim_{x \rightarrow 3^+} f(x) = \lim_{x \rightarrow 3^+} (5) = 5$$

It is observed that the left and right hand limit of  $f$  at  $x = 3$  do not coincide.

Therefore,  $f$  is discontinuous at  $x = 3$ .

Case V:

If  $3 < c \leq 10$ , then  $f(c) = 5$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (5) = 5$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points of the interval  $(3, 10]$ .

Hence,  $f$  is discontinuous at  $x = 1$  and  $x = 3$ .

### Question 15:

$$f(x) = \begin{cases} 2x, & \text{if } x < 0 \\ 0, & \text{if } 0 \leq x \leq 1 \\ 4x, & \text{if } x > 1 \end{cases}$$

Discuss the continuity of the function  $f$ , where  $f$  is defined by

### Solution:

$$f(x) = \begin{cases} 2x, & \text{if } x < 0 \\ 0, & \text{if } 0 \leq x \leq 1 \\ 4x, & \text{if } x > 1 \end{cases}$$

The given function is

The given function  $f$  is defined at all the points of the real line.

Let  $c$  be a point on the real line.

Case I:

If  $c < 0$ , then  $f(c) = 2c$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (2x) = 2c$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x < 0$ .

Case II:

If  $c = 0$ , then  $f(c) = f(0) = 0$

The left hand limit of  $f$  at  $x = 0$  is,

$$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} (2x) = 2(0) = 0$$

The right hand limit of  $f$  at  $x = 0$  is,

$$\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} (0) = 0$$

$$\therefore \lim_{x \rightarrow 0} f(x) = f(0)$$

Therefore,  $f$  is continuous at  $x = 0$

Case III:

If  $0 < c < 1$ , then  $f(c) = 0$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (0) = 0$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous in the interval  $(0,1)$ .

Case IV:

If  $c = 1$ , then  $f(c) = f(1) = 0$

The left hand limit of  $f$  at  $x = 1$  is,

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} (0) = 0$$

The right hand limit of  $f$  at  $x = 1$  is,

$$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} (4x) = 4(1) = 4$$

It is observed that the left and right hand limit of  $f$  at  $x = 1$  do not coincide.

Therefore,  $f$  is not continuous at  $x = 1$ .

Case V:

If  $c < 1$ , then  $f(c) = 4c$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (4x) = 4c$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x > 1$ .

Hence,  $f$  is not continuous only at  $x = 1$ .

### Question 16:

$$f(x) = \begin{cases} -2, & \text{if } x \leq -1 \\ 2x, & \text{if } -1 < x \leq 1 \\ 2, & \text{if } x > 1 \end{cases}$$

Discuss the continuity of the function  $f$ , where  $f$  is defined by

**Solution:**

$$f(x) = \begin{cases} -2, & \text{if } x \leq -1 \\ 2x, & \text{if } -1 < x \leq 1 \\ 2, & \text{if } x > 1 \end{cases}$$

The given function is

The given function  $f$  is defined at all the points.

Let  $c$  be a point on the real line.

Case I:

If  $c < -1$ , then  $f(c) = -2$



$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (-2) = -2$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x < -1$ .

Case II:

$$\text{If } c = -1, \text{ then } f(c) = f(-1) = -2$$

The left hand limit of  $f$  at  $x = -1$  is,

$$\lim_{x \rightarrow -1^-} f(x) = \lim_{x \rightarrow -1^-} (-2) = -2$$

The right hand limit of  $f$  at  $x = -1$  is,

$$\lim_{x \rightarrow -1^+} f(x) = \lim_{x \rightarrow -1^+} (2x) = 2(-1) = -2$$

$$\therefore \lim_{x \rightarrow -1} f(x) = f(-1)$$

Therefore,  $f$  is continuous at  $x = -1$

Case III:

$$\text{If } -1 < c < 1, \text{ then } f(c) = 2c$$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (2x) = 2c$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous in the interval  $(-1, 1)$ .

Case IV:

$$\text{If } c = 1, \text{ then } f(c) = f(1) = 2(1) = 2$$

The left hand limit of  $f$  at  $x = 1$  is,

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} (2x) = 2(1) = 2$$

The right hand limit of  $f$  at  $x = 1$  is,

$$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} (2) = 2$$

$$\therefore \lim_{x \rightarrow 1} f(x) = f(c)$$

Therefore,  $f$  is continuous at  $x = 2$ .

Case V:

$$\text{If } c > 1, \text{ then } f(c) = 2$$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (2) = 2$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$



Therefore,  $f$  is continuous at all points  $x$ , such that  $x > 1$ .

Thus, from the above observations, it can be concluded that  $f$  is continuous at all points of the real line.

### Question 17:

Find the relationship between  $a$  and  $b$  so that the function  $f$  defined by  $f(x) = \begin{cases} ax+1, & \text{if } x \leq 3 \\ bx+3, & \text{if } x > 3 \end{cases}$  is continuous at  $x = 3$ .

**Solution:**

The given function is  $f(x) = \begin{cases} ax+1, & \text{if } x \leq 3 \\ bx+3, & \text{if } x > 3 \end{cases}$

For  $f$  to be continuous at  $x = 3$ , then

$$\lim_{x \rightarrow 3^-} f(x) = \lim_{x \rightarrow 3^+} f(x) = f(3) \quad \dots(1)$$

Also,

$$\lim_{x \rightarrow 3^-} f(x) = \lim_{x \rightarrow 3^-} (ax+1) = 3a+1$$

$$\lim_{x \rightarrow 3^+} f(x) = \lim_{x \rightarrow 3^+} (bx+3) = 3b+3$$

$$f(3) = 3a+1$$

Therefore, from (1), we obtain

$$3a+1 = 3b+3 = 3a+1$$

$$\Rightarrow 3a+1 = 3b+3$$

$$\Rightarrow 3a = 3b+2$$

$$\Rightarrow a = b + \frac{2}{3}$$

Therefore, the required relationship is given by,  $a = b + \frac{2}{3}$ .

### Question 18:

For what value of  $\lambda$  is the function defined by  $f(x) = \begin{cases} \lambda(x^2 - 2x), & \text{if } x \leq 0 \\ 4x+1, & \text{if } x > 0 \end{cases}$  is continuous at  $x = 0$ ? What about continuity at  $x = 1$ ?

**Solution:**

$$f(x) = \begin{cases} \lambda(x^2 - 2x), & \text{if } x \leq 0 \\ 4x + 1, & \text{if } x > 0 \end{cases}$$

The given function is

If  $f$  is continuous at  $x = 0$ , then

$$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^+} f(x) = f(0)$$

$$\Rightarrow \lim_{x \rightarrow 0^-} \lambda(x^2 - 2x) = \lim_{x \rightarrow 0^+} (4x + 1) = \lambda(0^2 - 2 \times 0)$$

$$\Rightarrow \lambda(0^2 - 2 \times 0) = 4(0) + 1 = 0$$

$$\Rightarrow 0 = 1 = 0 \quad [\text{which is not possible}]$$

Therefore, there is no value of  $\lambda$  for which  $f$  is continuous at  $x = 0$ .

At  $x = 1$

$$f(1) = 4x + 1 = 4(1) + 1 = 5$$

$$\lim_{x \rightarrow 1} (4x + 1) = 4(1) + 1 = 5$$

$$\therefore \lim_{x \rightarrow 1} f(x) = f(1)$$

Therefore, for any values of  $\lambda$ ,  $f$  is continuous at  $x = 1$ .

**Question 19:**

Show that the function defined by  $g(x) = x - [x]$  is discontinuous at all integral point. Here  $[x]$  denotes the greatest integer less than or equal to  $x$ .

**Solution:**

The given function is  $g(x) = x - [x]$

It is evident that  $g$  is defined at all integral points.

Let  $n$  be an integer.

Then,

$$g(n) = n - [n] = n - n = 0$$

The left hand limit of  $g$  at  $x = n$  is,

$$\lim_{x \rightarrow n^-} g(x) = \lim_{x \rightarrow n^-} (x - [x]) = \lim_{x \rightarrow n^-} (x) - \lim_{x \rightarrow n^-} [x] = n - (n-1) = 1$$

The right hand limit of  $g$  at  $x = n$  is,

$$\lim_{x \rightarrow n^+} g(x) = \lim_{x \rightarrow n^+} (x - [x]) = \lim_{x \rightarrow n^+} (x) - \lim_{x \rightarrow n^+} [x] = n - n = 0$$

It is observed that the left and right hand limit of  $g$  at  $x = n$  do not coincide.

Therefore,  $g$  is not continuous at  $x = n$ .

Hence,  $g$  is discontinuous at all integral points.

### Question 20:

Is the function defined by  $f(x) = x^2 - \sin x + 5$  continuous at  $x = \pi$ ?

### Solution:

The given function is  $f(x) = x^2 - \sin x + 5$

It is evident that  $f$  is defined at  $x = \pi$ .

$$\text{At } x = \pi, f(x) = f(\pi) = \pi^2 - \sin \pi + 5 = \pi^2 - 0 + 5 = \pi^2 + 5$$

$$\text{Consider } \lim_{x \rightarrow \pi} f(x) = \lim_{x \rightarrow \pi} (x^2 - \sin x + 5)$$

Put  $x = \pi + h$ , it is evident that if  $x \rightarrow \pi$ , then  $h \rightarrow 0$

$$\begin{aligned} \therefore \lim_{x \rightarrow \pi} f(x) &= \lim_{x \rightarrow \pi} (x^2 - \sin x) + 5 \\ &= \lim_{h \rightarrow 0} [(\pi + h)^2 - \sin(\pi + h) + 5] \\ &= \lim_{h \rightarrow 0} (\pi + h)^2 - \lim_{h \rightarrow 0} \sin(\pi + h) + \lim_{h \rightarrow 0} 5 \\ &= (\pi + 0)^2 - \lim_{h \rightarrow 0} [\sin \pi \cos h + \cos \pi \sin h] + 5 \\ &= \pi^2 - \lim_{h \rightarrow 0} \sin \pi \cos h - \lim_{h \rightarrow 0} \cos \pi \sin h + 5 \\ &= \pi^2 - \sin \pi \cos 0 - \cos \pi \sin 0 + 5 \\ &= \pi^2 - 0(1) - (-1)0 + 5 \\ &= \pi^2 + 5 \\ &= f(\pi) \end{aligned}$$

Therefore, the given function  $f$  is continuous at  $x = \pi$ .

### Question 21:

Discuss the continuity of the following functions.

(i)  $f(x) = \sin x + \cos x$

(ii)  $f(x) = \sin x - \cos x$

(iii)  $f(x) = \sin x \times \cos x$

### Solution:

It is known that if  $g$  and  $h$  are two continuous functions, then  $g + h$ ,  $g - h$  and  $g \cdot h$  are also continuous.

Let  $g(x) = \sin x$  and  $h(x) = \cos x$  are continuous functions.

It is evident that  $g(x) = \sin x$  is defined for every real number.

Let  $c$  be a real number. Put  $x = c + h$

If  $x \rightarrow c$ , then  $h \rightarrow 0$

$$g(c) = \sin c$$

$$\lim_{x \rightarrow c} g(x) = \lim_{x \rightarrow c} \sin x$$

$$= \lim_{h \rightarrow 0} \sin(c + h)$$

$$= \lim_{h \rightarrow 0} [\sin c \cos h + \cos c \sin h]$$

$$= \lim_{h \rightarrow 0} (\sin c \cos h) + \lim_{h \rightarrow 0} (\cos c \sin h)$$

$$= \sin c \cos 0 + \cos c \sin 0$$

$$= \sin c(1) + \cos c(0)$$

$$= \sin c$$

$$\therefore \lim_{x \rightarrow c} g(x) = g(c)$$

Therefore,  $g(x) = \sin x$  is a continuous function.

Let  $h(x) = \cos x$

It is evident that  $h(x) = \cos x$  is defined for every real number.

Let  $c$  be a real number. Put  $x = c + h$

If  $x \rightarrow c$ , then  $h \rightarrow 0$

$$h(c) = \cos c$$

$$\begin{aligned}
 \lim_{x \rightarrow c} h(x) &= \lim_{x \rightarrow c} \cos x \\
 &= \lim_{h \rightarrow 0} \cos(c + h) \\
 &= \lim_{h \rightarrow 0} [\cos c \cos h - \sin c \sin h] \\
 &= \lim_{h \rightarrow 0} (\cos c \cos h) - \lim_{h \rightarrow 0} (\sin c \sin h) \\
 &= \cos c \cos 0 - \sin c \sin 0 \\
 &= \cos c(1) - \sin c(0) \\
 &= \cos c \\
 \therefore \lim_{x \rightarrow c} h(x) &= h(c)
 \end{aligned}$$

Therefore,  $h(x) = \cos x$  is a continuous function.

Therefore, it can be concluded that,

- (i)  $f(x) = g(x) + h(x) = \sin x + \cos x$  is a continuous function.
- (ii)  $f(x) = g(x) - h(x) = \sin x - \cos x$  is a continuous function.
- (iii)  $f(x) = g(x) \times h(x) = \sin x \times \cos x$  is a continuous function.

### Question 22:

Discuss the continuity of the cosine, cosecant, secant, and cotangent functions.

### Solution:

It is known that if  $g$  and  $h$  are two continuous functions, then

- (i)  $\frac{h(x)}{g(x)}, g(x) \neq 0$  is continuous.
- (ii)  $\frac{1}{g(x)}, g(x) \neq 0$  is continuous.
- (iii)  $\frac{1}{h(x)}, h(x) \neq 0$  is continuous.

Let  $g(x) = \sin x$  and  $h(x) = \cos x$  are continuous functions.

It is evident that  $g(x) = \sin x$  is defined for every real number.

Let  $c$  be a real number. Put  $x = c + h$

If  $x \rightarrow c$ , then  $h \rightarrow 0$

$$g(c) = \sin c$$

$$\begin{aligned}\lim_{x \rightarrow c} g(x) &= \lim_{x \rightarrow c} \sin x \\ &= \lim_{h \rightarrow 0} \sin(c + h) \\ &= \lim_{h \rightarrow 0} [\sin c \cos h + \cos c \sin h] \\ &= \lim_{h \rightarrow 0} (\sin c \cos h) + \lim_{h \rightarrow 0} (\cos c \sin h) \\ &= \sin c \cos 0 + \cos c \sin 0 \\ &= \sin c(1) + \cos c(0) \\ &= \sin c\end{aligned}$$

$$\therefore \lim_{x \rightarrow c} g(x) = g(c)$$

Therefore,  $g(x) = \sin x$  is a continuous function.

$$\text{Let } h(x) = \cos x$$

It is evident that  $h(x) = \cos x$  is defined for every real number.

Let  $c$  be a real number. Put  $x = c + h$

If  $x \rightarrow c$ , then  $h \rightarrow 0$

$$h(c) = \cos c$$

$$\begin{aligned}\lim_{x \rightarrow c} h(x) &= \lim_{x \rightarrow c} \cos x \\ &= \lim_{h \rightarrow 0} \cos(c + h) \\ &= \lim_{h \rightarrow 0} [\cos c \cos h - \sin c \sin h] \\ &= \lim_{h \rightarrow 0} (\cos c \cos h) - \lim_{h \rightarrow 0} (\sin c \sin h) \\ &= \cos c \cos 0 - \sin c \sin 0 \\ &= \cos c(1) - \sin c(0) \\ &= \cos c\end{aligned}$$

$$\therefore \lim_{x \rightarrow c} h(x) = h(c)$$

Therefore,  $h(x) = \cos x$  is a continuous function.

Therefore, it can be concluded that,

$$\operatorname{cosec} x = \frac{1}{\sin x}, \sin x \neq 0 \text{ is continuous.}$$

$$\Rightarrow \operatorname{cosec} x, x \neq n\pi (n \in \mathbb{Z}) \text{ is continuous.}$$

Therefore, cosecant is continuous except at  $x = n\pi (n \in \mathbb{Z})$

$$\sec x = \frac{1}{\cos x}, \cos x \neq 0 \text{ is continuous.}$$

$$\Rightarrow \sec x, x \neq (2n+1)\frac{\pi}{2} (n \in \mathbb{Z}) \text{ is continuous.}$$

Therefore, secant is continuous except at  $x = (2n+1)\frac{\pi}{2} (n \in \mathbb{Z})$

$\cot x = \frac{\cos x}{\sin x}, \sin x \neq 0$  is continuous.

$\Rightarrow \cot x, x \neq n\pi (n \in \mathbb{Z})$  is continuous.

Therefore, cotangent is continuous except at  $x = n\pi (n \in \mathbb{Z})$ .

### Question 23:

$$f(x) = \begin{cases} \frac{\sin x}{x}, & \text{if } x < 0 \\ x+1, & \text{if } x \geq 0 \end{cases}$$

Find the points of discontinuity of  $f$ , where

### Solution:

$$f(x) = \begin{cases} \frac{\sin x}{x}, & \text{if } x < 0 \\ x+1, & \text{if } x \geq 0 \end{cases}$$

The given function is

The given function  $f$  is defined at all the points of the real line.

Let  $c$  be a point on the real line.

Case I:

If  $c < 0$ , then  $f(c) = \frac{\sin c}{c}$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} \left( \frac{\sin x}{x} \right) = \frac{\sin c}{c}$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x < 0$ .

Case II:

If  $c > 0$ , then  $f(c) = c+1$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (x+1) = c+1$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x > 0$ .



Case III:

If  $c = 0$ , then  $f(c) = f(0) = 0 + 1 = 1$

The left hand limit of  $f$  at  $x = 0$  is,

$$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} \left( \frac{\sin x}{x} \right) = 1$$

The right hand limit of  $f$  at  $x = 0$  is,

$$\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} (x + 1) = 1$$

$$\therefore \lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^+} f(x) = f(0)$$

Therefore,  $f$  is continuous at  $x = 0$

From the above observations, it can be concluded that  $f$  is continuous at all points of the real line.

Thus,  $f$  has no point of discontinuity.

#### Question 24:

$$f(x) = \begin{cases} x^2 \sin \frac{1}{x}, & \text{if } x \neq 0 \\ 0, & \text{if } x = 0 \end{cases}$$

Determine if  $f$  defined by

is a continuous function?

**Solution:**

$$f(x) = \begin{cases} x^2 \sin \frac{1}{x}, & \text{if } x \neq 0 \\ 0, & \text{if } x = 0 \end{cases}$$

The given function is

The given function  $f$  is defined at all the points of the real line.

Let  $c$  be a point on the real line.

Case I:

If  $c \neq 0$ , then  $f(c) = c^2 \sin \frac{1}{c}$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} \left( x^2 \sin \frac{1}{x} \right) = \left( \lim_{x \rightarrow c} x^2 \right) \left( \lim_{x \rightarrow c} \sin \frac{1}{x} \right) = c^2 \sin \frac{1}{c}$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x \neq 0$ .

Case II:

If  $c = 0$ , then  $f(0) = 0$

$$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} \left( x^2 \sin \frac{1}{x} \right) = \lim_{x \rightarrow 0^-} \left( x^2 \sin \frac{1}{x} \right)$$

It is known that,  $-1 \leq \sin \frac{1}{x} \leq 1, x \neq 0$

$$\Rightarrow -x^2 \leq x^2 \sin \frac{1}{x} \leq x^2$$

$$\Rightarrow \lim_{x \rightarrow 0^-} (-x^2) \leq \lim_{x \rightarrow 0^-} \left( x^2 \sin \frac{1}{x} \right) \leq \lim_{x \rightarrow 0^-} x^2$$

$$\Rightarrow 0 \leq \lim_{x \rightarrow 0^-} \left( x^2 \sin \frac{1}{x} \right) \leq 0$$

$$\Rightarrow \lim_{x \rightarrow 0^-} \left( x^2 \sin \frac{1}{x} \right) = 0$$

$$\therefore \lim_{x \rightarrow 0^-} f(x) = 0$$

Similarly,

$$\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} \left( x^2 \sin \frac{1}{x} \right) = \lim_{x \rightarrow 0^+} \left( x^2 \sin \frac{1}{x} \right) = 0$$

$$\therefore \lim_{x \rightarrow 0^-} f(x) = f(0) = \lim_{x \rightarrow 0^+} f(x)$$

Therefore,  $f$  is continuous at  $x = 0$ .

From the above observations, it can be concluded that  $f$  is continuous at every point of the real line.

Thus,  $f$  is a continuous function.

### Question 25:

Examine the continuity of  $f$ , where  $f$  is defined by  $f(x) = \begin{cases} \sin x - \cos x, & \text{if } x \neq 0 \\ -1, & \text{if } x = 0 \end{cases}$

### Solution:

The given function is  $f(x) = \begin{cases} \sin x - \cos x, & \text{if } x \neq 0 \\ -1, & \text{if } x = 0 \end{cases}$

The given function  $f$  is defined at all the points of the real line.

Let  $c$  be a point on the real line.

Case I:

If  $c \neq 0$ , then  $f(c) = \sin c - \cos c$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (\sin x - \cos x) = \sin c - \cos c$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x \neq 0$ .

Case II:

If  $c = 0$ , then  $f(0) = -1$

$$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} (\sin x - \cos x) = \sin 0 - \cos 0 = 0 - 1 = -1$$

$$\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} (\sin x - \cos x) = \sin 0 - \cos 0 = 0 - 1 = -1$$

$$\lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} f(x) = f(0)$$

Therefore,  $f$  is continuous at  $x = 0$ .

From the above observations, it can be concluded that  $f$  is continuous at every point of the real line.

Thus,  $f$  is a continuous function.

### Question 26:

Find the values of  $k$  so that the function  $f$  is continuous at the indicated point

$$f(x) = \begin{cases} \frac{k \cos x}{\pi - 2x}, & \text{if } x \neq \frac{\pi}{2} \\ 3, & \text{if } x = \frac{\pi}{2} \end{cases} \quad \text{at } x = \frac{\pi}{2}$$

**Solution:**

$$f(x) = \begin{cases} \frac{k \cos x}{\pi - 2x}, & \text{if } x \neq \frac{\pi}{2} \\ 3, & \text{if } x = \frac{\pi}{2} \end{cases}$$

The given function is

The given function  $f$  is continuous at  $x = \frac{\pi}{2}$ , if  $f$  is defined at  $x = \frac{\pi}{2}$  and if the value of the  $f$  at  $x = \frac{\pi}{2}$  equals the limit of  $f$  at  $x = \frac{\pi}{2}$ .

It is evident that  $f$  is defined at  $x = \frac{\pi}{2}$  and  $f\left(\frac{\pi}{2}\right) = 3$

$$\lim_{x \rightarrow \frac{\pi}{2}} f(x) = \lim_{x \rightarrow \frac{\pi}{2}} \frac{k \cos x}{\pi - 2x}$$

Put  $x = \frac{\pi}{2} + h$

Then  $x \rightarrow \frac{\pi}{2} \Rightarrow h \rightarrow 0$

$$\begin{aligned}\therefore \lim_{x \rightarrow \frac{\pi}{2}} f(x) &= \lim_{x \rightarrow \frac{\pi}{2}} \frac{k \cos x}{\pi - 2x} = \lim_{h \rightarrow 0} \frac{k \cos\left(\frac{\pi}{2} + h\right)}{\pi - 2\left(\frac{\pi}{2} + h\right)} \\ &= k \lim_{h \rightarrow 0} \frac{-\sin h}{-2h} = \frac{k}{2} \lim_{h \rightarrow 0} \frac{\sin h}{h} = \frac{k}{2} \cdot 1 = \frac{k}{2} \\ \therefore \lim_{x \rightarrow \frac{\pi}{2}} f(x) &= f\left(\frac{\pi}{2}\right) \\ \Rightarrow \frac{k}{2} &= 3 \\ \Rightarrow k &= 6\end{aligned}$$

Therefore, the value of  $k = 6$ .

### Question 27:

Find the values of  $k$  so that the function  $f$  is continuous at the indicated point.

$$f(x) = \begin{cases} kx^2, & \text{if } x \leq 2 \\ 3, & \text{if } x > 2 \end{cases} \quad \text{at } x = 2$$

### Solution:

The given function is  $f(x) = \begin{cases} kx^2, & \text{if } x \leq 2 \\ 3, & \text{if } x > 2 \end{cases}$

The given function  $f$  is continuous at  $x = 2$ , if  $f$  is defined at  $x = 2$  and if the value of the  $f$  at  $x = 2$  equals the limit of  $f$  at  $x = 2$ .

It is evident that  $f$  is defined at  $x = 2$  and  $f(2) = k(2)^2 = 4k$

$$\lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^+} f(x) = f(2)$$

$$\Rightarrow \lim_{x \rightarrow 2^-} (kx^2) = \lim_{x \rightarrow 2^+} (3) = 4k$$

$$\Rightarrow k \times 2^2 = 3 = 4k$$

$$\Rightarrow 4k = 3$$

$$\Rightarrow k = \frac{3}{4}$$

Therefore, the value of  $k = \frac{3}{4}$ .

### Question 28:

Find the values of  $k$  so that the function  $f$  is continuous at the indicated point

$$f(x) = \begin{cases} kx+1, & \text{if } x \leq \pi \\ \cos x, & \text{if } x > \pi \end{cases} \text{ at } x = \pi$$

### Solution:

The given function is  $f(x) = \begin{cases} kx+1, & \text{if } x \leq \pi \\ \cos x, & \text{if } x > \pi \end{cases}$

The given function  $f$  is continuous at  $x = \pi$ , if  $f$  is defined at  $x = \pi$  and if the value of the  $f$  at  $x = \pi$  equals the limit of  $f$  at  $x = \pi$ .

It is evident that  $f$  is defined at  $x = \pi$  and  $f(\pi) = k\pi + 1$

$$\lim_{x \rightarrow \pi^-} f(x) = \lim_{x \rightarrow \pi^+} f(x) = f(\pi)$$

$$\Rightarrow \lim_{x \rightarrow \pi^-} (kx+1) = \lim_{x \rightarrow \pi^+} (\cos x) = k\pi + 1$$

$$\Rightarrow k\pi + 1 = \cos \pi = k\pi + 1$$

$$\Rightarrow k\pi + 1 = -1 = k\pi + 1$$

$$\Rightarrow k = -\frac{2}{\pi}$$

Therefore, the value of  $k = -\frac{2}{\pi}$ .

### Question 29:

Find the values of  $k$  so that the function  $f$  is continuous at the indicated point

$$f(x) = \begin{cases} kx+1, & \text{if } x \leq 5 \\ 3x-5, & \text{if } x > 5 \end{cases} \text{ at } x = 5.$$

### Solution:

The given function is  $f(x) = \begin{cases} kx+1, & \text{if } x \leq 5 \\ 3x-5, & \text{if } x > 5 \end{cases}$

The given function  $f$  is continuous at  $x = 5$ , if  $f$  is defined at  $x = 5$  and if the value of the  $f$  at  $x = 5$  equals the limit of  $f$  at  $x = 5$ .

It is evident that  $f$  is defined at  $x = 5$  and  $f(5) = kx+1 = 5k+1$

$$\begin{aligned}\lim_{x \rightarrow 5^-} f(x) &= \lim_{x \rightarrow 5^+} f(x) = f(5) \\ \Rightarrow \lim_{x \rightarrow 5^-} (kx + 1) &= \lim_{x \rightarrow 5^+} (3x - 5) = 5k + 1 \\ \Rightarrow 5k + 1 &= 3(5) - 5 = 5k + 1 \\ \Rightarrow 5k + 1 &= 15 - 5 = 5k + 1 \\ \Rightarrow 5k + 1 &= 10 = 5k + 1 \\ \Rightarrow 5k + 1 &= 10 \\ \Rightarrow 5k &= 9 \\ \Rightarrow k &= \frac{9}{5}\end{aligned}$$

Therefore, the value of  $k = \frac{9}{5}$ .

### Question 30:

Find the values of  $a$  &  $b$  such that the function defined by  $f(x) = \begin{cases} 5, & \text{if } x \leq 2 \\ ax + b, & \text{if } 2 < x < 10 \\ 21, & \text{if } x \geq 10 \end{cases}$ , is a continuous function.

### Solution:

$$f(x) = \begin{cases} 5, & \text{if } x \leq 2 \\ ax + b, & \text{if } 2 < x < 10 \\ 21, & \text{if } x \geq 10 \end{cases}$$

The given function is

It is evident that  $f$  is defined at all points of the real line.

If  $f$  is a continuous function, then  $f$  is continuous at all real numbers.

In particular,  $f$  is continuous at  $x = 2$  and  $x = 10$

Since  $f$  is continuous at  $x = 2$ , we obtain

$$\begin{aligned}\lim_{x \rightarrow 2^-} f(x) &= \lim_{x \rightarrow 2^+} f(x) = f(2) \\ \Rightarrow \lim_{x \rightarrow 2^-} (5) &= \lim_{x \rightarrow 2^+} (ax + b) = 5 \\ \Rightarrow 5 &= 2a + b = 5 \\ \Rightarrow 2a + b &= 5 \quad \dots(1)\end{aligned}$$

Since  $f$  is continuous at  $x = 10$ , we obtain

$$\begin{aligned}\lim_{x \rightarrow 10^-} f(x) &= \lim_{x \rightarrow 10^+} f(x) = f(10) \\ \Rightarrow \lim_{x \rightarrow 10^-} (ax + b) &= \lim_{x \rightarrow 10^+} (21) = 21 \\ \Rightarrow 10a + b &= 21 = 21 \\ \Rightarrow 10a + b &= 21 \quad \dots(2)\end{aligned}$$

On subtracting equation (1) from equation (2), we obtain

$$8a = 16$$

$$\Rightarrow a = 2$$

By putting  $a = 2$  in equation (1), we obtain

$$2(2) + b = 5$$

$$\Rightarrow 4 + b = 5$$

$$\Rightarrow b = 1$$

Therefore, the values of  $a$  and  $b$  for which  $f$  is a continuous function are 2 and 1 respectively.

### Question 31:

Show that the function defined by  $f(x) = \cos(x^2)$  is a continuous function.

#### Solution:

The given function is  $f(x) = \cos(x^2)$ .

This function  $f$  is defined for every real number and  $f$  can be written as the composition of two functions as,

$$f = goh, \text{ where } g(x) = \cos x \text{ and } h(x) = x^2$$

$$[\because (goh)(x) = g(h(x)) = g(x^2) = \cos(x^2) = f(x)]$$

It has to be proved first that  $g(x) = \cos x$  and  $h(x) = x^2$  are continuous functions.

It is evident that  $g$  is defined for every real number.

Let  $c$  be a real number.

Let  $g(c) = \cos c$ . Put  $x = c + h$

If  $x \rightarrow c$ , then  $h \rightarrow 0$

$$\begin{aligned} \lim_{x \rightarrow c} g(x) &= \lim_{x \rightarrow c} \cos x \\ &= \lim_{h \rightarrow 0} \cos(c + h) \\ &= \lim_{h \rightarrow 0} [\cos c \cos h - \sin c \sin h] \\ &= \lim_{h \rightarrow 0} (\cos c \cos h) - \lim_{h \rightarrow 0} (\sin c \sin h) \\ &= \cos c \cos 0 - \sin c \sin 0 \\ &= \cos c(1) - \sin c(0) \\ &= \cos c \\ \therefore \lim_{x \rightarrow c} g(x) &= g(c) \end{aligned}$$

Therefore,  $g(x) = \cos x$  is a continuous function.

Let  $h(x) = x^2$

It is evident that  $h$  is defined for every real number.

Let  $k$  be a real number, then  $h(k) = k^2$

$$\lim_{x \rightarrow k} h(x) = \lim_{x \rightarrow k} x^2 = k^2$$

$$\therefore \lim_{x \rightarrow k} h(x) = h(k)$$

Therefore,  $h$  is a continuous function.

It is known that for real valued functions  $g$  and  $h$ , such that  $(goh)$  is defined at  $c$ , if  $g$  is continuous at  $c$  and if  $f$  is continuous at  $g(c)$ , then  $(fog)$  is continuous at  $c$ .

Therefore,  $f(x) = (goh)(x) = \cos(x^2)$  is a continuous function.

### Question 32:

Show that the function defined by  $f(x) = |\cos x|$  is a continuous function.

### Solution:

The given function is  $f(x) = |\cos x|$ .

This function  $f$  is defined for every real number and  $f$  can be written as the composition of two functions as,

$$f = goh, \text{ where } g(x) = |x| \text{ and } h(x) = \cos x$$

$$[\because (goh)(x) = g(h(x)) = g(\cos x) = |\cos x| = f(x)]$$

It has to be proved first that  $g(x) = |x|$  and  $h(x) = \cos x$  are continuous functions.

$$g(x) = |x| \text{ can be written as } g(x) = \begin{cases} -x, & \text{if } x < 0 \\ x, & \text{if } x \geq 0 \end{cases}$$

It is evident that  $g$  is defined for every real number.

Let  $c$  be a real number.

Case I:

If  $c < 0$ , then  $g(c) = -c$

$$\lim_{x \rightarrow c} g(x) = \lim_{x \rightarrow c} (-x) = -c$$

$$\therefore \lim_{x \rightarrow c} g(x) = g(c)$$

Therefore,  $g$  is continuous at all points  $x$ , such that  $x < 0$ .

Case II:

If  $c > 0$ , then  $g(c) = c$



$$\lim_{x \rightarrow c} g(x) = \lim_{x \rightarrow c} (x) = c$$

$$\therefore \lim_{x \rightarrow c} g(x) = g(c)$$

Therefore,  $g$  is continuous at all points  $x$ , such that  $x > 0$ .

Case III:

If  $c = 0$ , then  $g(c) = g(0) = 0$

$$\lim_{x \rightarrow 0^-} g(x) = \lim_{x \rightarrow 0^-} (-x) = 0$$

$$\lim_{x \rightarrow 0^+} g(x) = \lim_{x \rightarrow 0^+} (x) = 0$$

$$\therefore \lim_{x \rightarrow 0^-} g(x) = \lim_{x \rightarrow 0^+} g(x) = g(0)$$

Therefore,  $g$  is continuous at all  $x = 0$ .

From the above three observations, it can be concluded that  $g$  is continuous at all points.

Let  $h(x) = \cos x$

It is evident that  $h(x) = \cos x$  is defined for every real number.

Let  $c$  be a real number. Put  $x = c + h$

If  $x \rightarrow c$ , then  $h \rightarrow 0$

$$h(c) = \cos c$$

$$\lim_{x \rightarrow c} h(x) = \lim_{x \rightarrow c} \cos x$$

$$= \lim_{h \rightarrow 0} \cos(c + h)$$

$$= \lim_{h \rightarrow 0} [\cos c \cos h - \sin c \sin h]$$

$$= \lim_{h \rightarrow 0} (\cos c \cos h) - \lim_{h \rightarrow 0} (\sin c \sin h)$$

$$= \cos c \cos 0 - \sin c \sin 0$$

$$= \cos c(1) - \sin c(0)$$

$$= \cos c$$

$$\therefore \lim_{x \rightarrow c} h(x) = h(c)$$

Therefore,  $h(x) = \cos x$  is a continuous function.

It is known that for real valued functions  $g$  and  $h$ , such that  $(goh)$  is defined at  $c$ , if  $g$  is continuous at  $c$  and if  $f$  is continuous at  $g(c)$ , then  $(fog)$  is continuous at  $c$ .

Therefore,  $f(x) = (goh)(x) = g(h(x)) = g(\cos x) = |\cos x|$  is a continuous function.

### Question 33:

Show that the function defined by  $f(x) = |\sin x|$  is a continuous function.

### Solution:

The given function is  $f(x) = |\sin x|$ .

This function  $f$  is defined for every real number and  $f$  can be written as the composition of two functions as,

$f = goh$ , where  $g(x) = |x|$  and  $h(x) = \sin x$

$$[\because (goh)(x) = g(h(x)) = g(\sin x) = |\sin x| = f(x)]$$

It has to be proved first that  $g(x) = |x|$  and  $h(x) = \sin x$  are continuous functions.

$g(x) = |x|$  can be written as  $g(x) = \begin{cases} -x, & \text{if } x < 0 \\ x, & \text{if } x \geq 0 \end{cases}$

It is evident that  $g$  is defined for every real number.

Let  $c$  be a real number.

Case I:

If  $c < 0$ , then  $g(c) = -c$

$$\lim_{x \rightarrow c} g(x) = \lim_{x \rightarrow c} (-x) = -c$$

$$\therefore \lim_{x \rightarrow c} g(x) = g(c)$$

Therefore,  $g$  is continuous at all points  $x$ , such that  $x < 0$ .

Case II:

If  $c > 0$ , then  $g(c) = c$

$$\lim_{x \rightarrow c} g(x) = \lim_{x \rightarrow c} (x) = c$$

$$\therefore \lim_{x \rightarrow c} g(x) = g(c)$$

Therefore,  $g$  is continuous at all points  $x$ , such that  $x > 0$ .

Case III:

If  $c = 0$ , then  $g(c) = g(0) = 0$

$$\lim_{x \rightarrow 0^-} g(x) = \lim_{x \rightarrow 0^-} (-x) = 0$$

$$\lim_{x \rightarrow 0^+} g(x) = \lim_{x \rightarrow 0^+} (x) = 0$$

$$\therefore \lim_{x \rightarrow 0^-} g(x) = \lim_{x \rightarrow 0^+} g(x) = g(0)$$

Therefore,  $g$  is continuous at all  $x = 0$ .

From the above three observations, it can be concluded that  $g$  is continuous at all points.

Let  $h(x) = \sin x$

It is evident that  $h(x) = \sin x$  is defined for every real number.

Let  $c$  be a real number. Put  $x = c + k$

If  $x \rightarrow c$ , then  $k \rightarrow 0$

$$h(c) = \sin c$$

$$\begin{aligned}\lim_{x \rightarrow c} h(x) &= \lim_{x \rightarrow c} \sin x \\ &= \lim_{k \rightarrow 0} \sin(c + k) \\ &= \lim_{k \rightarrow 0} [\sin c \cos k + \cos c \sin k] \\ &= \lim_{k \rightarrow 0} (\sin c \cos k) + \lim_{k \rightarrow 0} (\cos c \sin k) \\ &= \sin c \cos 0 + \cos c \sin 0 \\ &= \sin c(1) + \cos c(0) \\ &= \sin c \\ \therefore \lim_{x \rightarrow c} h(x) &= h(c)\end{aligned}$$

Therefore,  $h(x) = \sin x$  is a continuous function.

It is known that for real valued functions  $g$  and  $h$ , such that  $(goh)$  is defined at  $c$ , if  $g$  is continuous at  $c$  and if  $f$  is continuous at  $g(c)$ , then  $(fog)$  is continuous at  $c$ .

Therefore,  $f(x) = (goh)(x) = g(h(x)) = g(\sin x) = |\sin x|$  is a continuous function.

### Question 34:

Find all the points of discontinuity of  $f$  defined by  $f(x) = |x| - |x+1|$ .

### Solution:

The given function is  $f(x) = |x| - |x+1|$ .

The two functions,  $g$  and  $h$  are defined as  $g(x) = |x|$  and  $h(x) = |x+1|$ .

Then,  $f = g - h$

The continuity of  $g$  and  $h$  are examined first.

$$g(x) = |x| \text{ can be written as } g(x) = \begin{cases} -x, & \text{if } x < 0 \\ x, & \text{if } x \geq 0 \end{cases}$$

It is evident that  $g$  is defined for every real number.

Let  $c$  be a real number.

Case I:

If  $c < 0$ , then  $g(c) = -c$

$$\lim_{x \rightarrow c} g(x) = \lim_{x \rightarrow c} (-x) = -c$$

$$\therefore \lim_{x \rightarrow c} g(x) = g(c)$$

Therefore,  $g$  is continuous at all points  $x$ , such that  $x < 0$ .

Case II:

If  $c > 0$ , then  $g(c) = c$

$$\lim_{x \rightarrow c} g(x) = \lim_{x \rightarrow c} (x) = c$$

$$\therefore \lim_{x \rightarrow c} g(x) = g(c)$$

Therefore,  $g$  is continuous at all points  $x$ , such that  $x > 0$ .

Case III:

If  $c = 0$ , then  $g(c) = g(0) = 0$

$$\lim_{x \rightarrow 0^-} g(x) = \lim_{x \rightarrow 0^-} (-x) = 0$$

$$\lim_{x \rightarrow 0^+} g(x) = \lim_{x \rightarrow 0^+} (x) = 0$$

$$\therefore \lim_{x \rightarrow 0} g(x) = \lim_{x \rightarrow 0} (x) = g(0)$$

Therefore,  $g$  is continuous at all  $x = 0$ .

From the above three observations, it can be concluded that  $g$  is continuous at all points.

$$h(x) = |x+1| \text{ can be written as } h(x) = \begin{cases} -(x+1), & \text{if } x < -1 \\ x+1, & \text{if } x \geq -1 \end{cases}$$

It is evident that  $h$  is defined for every real number.

Let  $c$  be a real number.

Case I:

If  $c < -1$ , then  $h(c) = -(c+1)$

$$\lim_{x \rightarrow c} h(x) = \lim_{x \rightarrow c} [-(x+1)] = -(c+1)$$

$$\therefore \lim_{x \rightarrow c} h(x) = h(c)$$

Therefore,  $h$  is continuous at all points  $x$ , such that  $x < -1$ .

Case II:

If  $c > -1$ , then  $h(c) = c+1$

$$\lim_{x \rightarrow c} h(x) = \lim_{x \rightarrow c} (x+1) = c+1$$

$$\therefore \lim_{x \rightarrow c} h(x) = h(c)$$

Therefore,  $h$  is continuous at all points  $x$ , such that  $x > -1$ .

Case III:

If  $c = -1$ , then  $h(c) = h(-1) = -1+1 = 0$

$$\lim_{x \rightarrow -1^-} h(x) = \lim_{x \rightarrow -1^-} [-(x+1)] = -(-1+1) = 0$$

$$\lim_{x \rightarrow -1^+} h(x) = \lim_{x \rightarrow -1^+} (x+1) = (-1+1) = 0$$

$$\therefore \lim_{x \rightarrow -1} h(x) = \lim_{x \rightarrow -1} h(x) = h(-1)$$

Therefore,  $h$  is continuous at  $x = -1$ .

From the above three observations, it can be concluded that  $h$  is continuous at all points.

It concludes that  $g$  and  $h$  are continuous functions. Therefore,  $f = g - h$  is also a continuous function.

Therefore,  $f$  has no point of discontinuity.

## EXERCISE 5.2

### Question 1:

Differentiate the function with respect to  $x$ .

$$\sin(x^2 + 5)$$

### Solution:

Let  $f(x) = \sin(x^2 + 5)$ ,  $u(x) = x^2 + 5$  and  $v(t) = \sin t$

Then,  $(v \circ u)(x) = v(u(x)) = v(x^2 + 5) = \sin(x^2 + 5) = f(x)$

Thus,  $f$  is a composite of two functions.

Put  $t = u(x) = x^2 + 5$

Then, we get

$$\frac{dv}{dt} = \frac{d}{dt}(\sin t) = \cos t = \cos(x^2 + 5)$$

$$\frac{dt}{dx} = \frac{d}{dx}(x^2 + 5) = \frac{d}{dx}(x^2) + \frac{d}{dx}(5) = 2x + 0 = 2x$$

By chain rule of derivative,

$$\frac{df}{dx} = \frac{dv}{dt} \cdot \frac{dt}{dx} = \cos(x^2 + 5) \times 2x = 2x \cos(x^2 + 5)$$

### Alternate method:

$$\begin{aligned} \frac{d}{dx}[\sin(x^2 + 5)] &= \cos(x^2 + 5) \cdot \frac{d}{dx}(x^2 + 5) \\ &= \cos(x^2 + 5) \cdot \left[ \frac{d}{dx}(x^2) + \frac{d}{dx}(5) \right] \\ &= \cos(x^2 + 5) \cdot [2x + 0] \\ &= 2x \cos(x^2 + 5) \end{aligned}$$

### Question 2:

Differentiate the function with respect to  $x$

$$\cos(\sin x)$$

### Solution:

Let  $f(x) = \cos(\sin x)$ ,  $u(x) = \sin x$  and  $v(t) = \cos t$

Then,  $(v \circ u)(x) = v(u(x)) = v(\sin x) = \cos(\sin x) = f(x)$

Here,  $f$  is a composite function of two functions.

Put  $t = u(x) = \sin x$

$$\therefore \frac{dv}{dt} = \frac{d}{dt} [\cos t] = -\sin t = -\sin(\sin x)$$

$$\frac{dt}{dx} = \frac{d}{dx} (\sin x) = \cos x$$

By chain rule,

$$\frac{df}{dx} = \frac{dv}{dt} \cdot \frac{dt}{dx} = -\sin(\sin x) \cdot \cos x = -\cos x \sin(\sin x)$$

**Alternate method:**

$$\begin{aligned} \frac{d}{dx} [\cos(\sin x)] &= -\sin(\sin x) \cdot \frac{d}{dx} (\sin x) \\ &= -\sin(\sin x) \times \cos x \\ &= -\cos x \sin(\sin x) \end{aligned}$$

### Question 3:

Differentiate the function with respect to  $x$

$$\sin(ax + b)$$

**Solution:**

Let  $f(x) = \sin(ax + b)$ ,  $u(x) = ax + b$  and  $v(t) = \sin t$

Then,  $(f \circ u)(x) = v(u(x)) = v(ax + b) = \sin(ax + b) = f(x)$

Here,  $f$  is a composite function of two functions  $u$  and  $v$ .

Put,  $t = u(x) = ax + b$

Thus,

$$\frac{dv}{dt} = \frac{d}{dt} (\sin t) = \cos t = \cos(ax + b)$$

$$\frac{dt}{dx} = \frac{d}{dx} (ax + b) = \frac{d}{dx} (ax) + \frac{d}{dx} (b) = a + 0 = a$$

Hence, by chain rule, we get

$$\frac{df}{dx} = \frac{dv}{dt} \cdot \frac{dt}{dx} = \cos(ax + b) \cdot a = a \cos(ax + b)$$

**Alternate method:**

$$\begin{aligned}\frac{d}{dx}[\sin(ax+b)] &= \cos(ax+b) \cdot \frac{d}{dx}(ax+b) \\ &= \cos(ax+b) \cdot \left[ \frac{d}{dx}(ax) + \frac{d}{dx}(b) \right] \\ &= \cos(ax+b) \cdot (a+0) \\ &= a \cos(ax+b)\end{aligned}$$

#### Question 4:

Differentiate the function with respect to  $x$

$$\sec(\tan(\sqrt{x}))$$

#### Solution:

Let  $f(x) = \sec(\tan(\sqrt{x}))$ ,  $u(x) = \sqrt{x}$ ,  $v(t) = \tan t$  and  $w(s) = \sec s$

Then,  $(w \circ v \circ u)(x) = w[v(u(x))] = w[v(\sqrt{x})] = w(\tan \sqrt{x}) = \sec(\tan \sqrt{x}) = f(x)$

Here,  $f$  is a composite function of three functions  $u$ ,  $v$  and  $w$ .

Put,  $s = v(t) = \tan t$  and  $t = u(x) = \sqrt{x}$

Then,

$$\begin{aligned}\frac{dw}{ds} &= \frac{d}{ds}(\sec s) \\ &= \sec s \tan s \\ &= \sec(\tan t) \cdot \tan(\tan t) \quad [s = \tan t] \\ &= \sec(\tan \sqrt{x}) \cdot \tan(\tan \sqrt{x}) \quad [t = \sqrt{x}]\end{aligned}$$

Now,

$$\begin{aligned}\frac{ds}{dt} &= \frac{d}{dt}(\tan t) = \sec^2 t = \sec^2 \sqrt{x} \\ \frac{dt}{dx} &= \frac{d}{dx}(\sqrt{x}) = \frac{d}{dx}\left(x^{\frac{1}{2}}\right) = \frac{1}{2} \cdot x^{\frac{1}{2}-1} = \frac{1}{2\sqrt{x}}\end{aligned}$$

Hence, by chain rule, we get



$$\begin{aligned}\frac{d}{dx} \left[ \sec(\tan \sqrt{x}) \right] &= \frac{dw}{ds} \cdot \frac{ds}{dt} \cdot \frac{dt}{dx} \\ &= \sec(\tan \sqrt{x}) \cdot \tan(\tan \sqrt{x}) \cdot \sec^2 \sqrt{x} \cdot \frac{1}{2\sqrt{x}} \\ &= \frac{1}{2\sqrt{x}} \sec^2 \sqrt{x} \sec(\tan \sqrt{x}) \tan(\tan \sqrt{x}) \\ &= \frac{\sec^2 \sqrt{x} \sec(\tan \sqrt{x}) \tan(\tan \sqrt{x})}{2\sqrt{x}}\end{aligned}$$

**Alternate method:**

$$\begin{aligned}\frac{d}{dx} \left[ \sec(\tan \sqrt{x}) \right] &= \sec(\tan \sqrt{x}) \cdot \tan(\tan \sqrt{x}) \cdot \frac{d}{dx}(\tan \sqrt{x}) \\ &= \sec(\tan \sqrt{x}) \cdot \tan(\tan \sqrt{x}) \cdot \sec^2(\sqrt{x}) \cdot \frac{d}{dx}(\sqrt{x}) \\ &= \sec(\tan \sqrt{x}) \cdot \tan(\tan \sqrt{x}) \cdot \sec^2(\sqrt{x}) \cdot \frac{1}{2\sqrt{x}} \\ &= \frac{\sec(\tan \sqrt{x}) \cdot \tan(\tan \sqrt{x}) \cdot \sec^2(\sqrt{x})}{2\sqrt{x}}\end{aligned}$$

### Question 5:

Differentiate the function with respect to  $x$

$$\frac{\sin(ax+b)}{\cos(cx+d)}$$

**Solution:**

Given,  $f(x) = \frac{\sin(ax+b)}{\cos(cx+d)}$ , where  $g(x) = \sin(ax+b)$  and  $h(x) = \cos(cx+d)$

$$\therefore f = \frac{g'h - gh'}{h^2}$$

Consider  $g(x) = \sin(ax+b)$

Let  $u(x) = ax+b$ ,  $v(t) = \sin t$

Then  $(v \circ u)(x) = v(u(x)) = v(ax+b) = \sin(ax+b) = g(x)$

$\therefore g$  is a composite function of two functions,  $u$  and  $v$ .

Put,  $t = u(x) = ax+b$

$$\frac{dv}{dt} = \frac{d}{dt}(\sin t) = \cos t = \cos(ax+b)$$

$$\frac{dt}{dx} = \frac{d}{dx}(ax+b) = \frac{d}{dx}(ax) + \frac{d}{dx}(b) = a + 0 = a$$

Thus, by chain rule, we get

$$g' = \frac{dg}{dx} = \frac{dv}{dt} \cdot \frac{dt}{dx} = \cos(ax+b) \cdot a = a \cos(ax+b)$$

Consider  $h(x) = \cos(cx+d)$

Let  $p(x) = cx+d$ ,  $q(y) = \cos y$

Then,  $(q \circ p)(x) = q(p(x)) = q(cx+d) = \cos(cx+d) = h(x)$

$\therefore h$  is a composite function of two functions,  $p$  and  $q$ .

Put,  $y = p(x) = cx+d$

$$\frac{dq}{dy} = \frac{d}{dy}(\cos y) = -\sin y = -\sin(cx+d)$$

$$\frac{dy}{dx} = \frac{d}{dx}(cx+d) = \frac{d}{dx}(cx) + \frac{d}{dx}(d) = c$$

Using chain rule, we get

$$\begin{aligned} h' &= \frac{dh}{dx} = \frac{dq}{dy} \cdot \frac{dy}{dx} \\ &= -\sin(cx+d) \times c \\ &= -c \sin(cx+d) \end{aligned}$$

Therefore,

$$\begin{aligned} f' &= \frac{a \cos(ax+b) \cdot \cos(cx+d) - \sin(ax+b) \{-c \sin(cx+d)\}}{[\cos(cx+d)]^2} \\ &= \frac{a \cos(ax+b)}{\cos(cx+d)} + c \sin(ax+b) \cdot \frac{\sin(cx+d)}{\cos(cx+d)} \times \frac{1}{\cos(cx+d)} \\ &= a \cos(ax+b) \sec(cx+d) + c \sin(ax+b) \tan(cx+d) \sec(cx+d) \end{aligned}$$

### Question 6:

Differentiate the function with respect to  $x$

$$\cos x^3 \cdot \sin^2(x^5)$$

### Solution:

$$\text{Given, } \cos x^3 \cdot \sin^2(x^5)$$



$$\begin{aligned}
 \frac{d}{dx} [\cos x^3 \cdot \sin^2(x^5)] &= \sin^2(x^5) \times \frac{d}{dx} (\cos x^3) + \cos x^3 \times \frac{d}{dx} [\sin^2(x^5)] \\
 &= \sin^2(x^5) \times (-\sin x^3) \times \frac{d}{dx} (x^3) + \cos x^3 \times 2 \sin(x^5) \cdot \frac{d}{dx} [\sin x^5] \\
 &= -\sin x^3 \sin^2(x^5) \times 3x^2 + 2 \sin x^5 \cos x^3 \cdot \cos x^5 \times \frac{d}{dx} (x^5) \\
 &= -3x^2 \sin x^3 \cdot \sin^2(x^5) + 2 \sin x^5 \cos x^5 \cos x^3 \times 5x^4 \\
 &= 10x^4 \sin x^5 \cos x^5 \cos x^3 - 3x^2 \sin x^3 \sin^2(x^5)
 \end{aligned}$$

**Question 7:**Differentiate the function with respect to  $x$ 

$$2\sqrt{\cot(x^2)}$$

**Solution:**

$$\begin{aligned}
 \frac{d}{dx} [2\sqrt{\cot(x^2)}] &= 2 \cdot \frac{1}{2\sqrt{\cot(x^2)}} \times \frac{d}{dx} [\cot(x^2)] \\
 &= \frac{\sqrt{\sin(x^2)}}{\sqrt{\cos(x^2)}} \times -\operatorname{cosec}^2(x^2) \times \frac{d}{dx} (x^2) \\
 &= \frac{\sqrt{\sin(x^2)}}{\sqrt{\cos(x^2)}} \times \frac{-1}{\sin^2(x^2)} \times (2x) \\
 &= \frac{-2x}{\sin x^2 \sqrt{\cos x^2 \sin x^2}} \\
 &= \frac{-2\sqrt{2}x}{\sin x^2 \sqrt{2 \sin x^2 \cos x^2}} \\
 &= \frac{-2\sqrt{2}x}{\sin x^2 \sqrt{\sin 2x^2}}
 \end{aligned}$$

**Question 8:**Differentiate the function with respect to  $x$ 

$$\cos(\sqrt{x})$$

**Solution:**

$$\text{Let } f(x) = \cos(\sqrt{x})$$

$$\text{Also, let } u(x) = \sqrt{x} \text{ and, } v(t) = \cos t$$

Then,

$$\begin{aligned}(v \circ u)(x) &= v(u(x)) \\ &= v(\sqrt{x}) \\ &= \cos \sqrt{x} \\ &= f(x)\end{aligned}$$

Since,  $f$  is a composite function of  $u$  and  $v$ .

$$t = u(x) = \sqrt{x}$$

Then,

$$\begin{aligned}\frac{dt}{dx} &= \frac{d}{dx}(\sqrt{x}) = \frac{d}{dx}\left(x^{\frac{1}{2}}\right) = \frac{1}{2}x^{-\frac{1}{2}} \\ &= \frac{1}{2\sqrt{x}} \\ \frac{dv}{dt} &= \frac{d}{dt}(\cos t) = -\sin t\end{aligned}$$

And,  $= -\sin(\sqrt{x})$

Using chain rule, we get

$$\begin{aligned}\frac{df}{dx} &= \frac{dv}{dt} \cdot \frac{dt}{dx} \\ &= -\sin(\sqrt{x}) \cdot \frac{1}{2\sqrt{x}} \\ &= -\frac{1}{2\sqrt{x}} \sin(\sqrt{x}) \\ &= -\frac{\sin(\sqrt{x})}{2\sqrt{x}}\end{aligned}$$

**Alternate method:**

$$\begin{aligned}\frac{d}{dx}[\cos(\sqrt{x})] &= -\sin(\sqrt{x}) \cdot \frac{d}{dx}(\sqrt{x}) \\ &= -\sin(\sqrt{x}) \times \frac{d}{dx}\left(x^{\frac{1}{2}}\right) \\ &= -\sin \sqrt{x} \times \frac{1}{2}x^{-\frac{1}{2}} \\ &= \frac{-\sin \sqrt{x}}{2\sqrt{x}}\end{aligned}$$

### Question 9:

Prove that the function  $f$  given by

$f(x) = |x-1|, x \in \mathbf{R}$  is not differentiable at  $x = 1$ .

### Solution:

Given,  $f(x) = |x-1|, x \in \mathbf{R}$

It is known that a function  $f$  is differentiable at a point  $x = c$  in its domain if both

$\lim_{h \rightarrow 0^-} \frac{f(c) - f(c-h)}{h}$  and  $\lim_{h \rightarrow 0^+} \frac{f(c+h) - f(c)}{h}$  are finite and equal.

To check the differentiability of the given function at  $x = 1$ ,

Consider LHD at  $x = 1$

$$\begin{aligned} \lim_{h \rightarrow 0^-} \frac{f(1) - f(1-h)}{h} &= \lim_{h \rightarrow 0^-} \frac{f|1-1| - |1-h-1|}{h} \\ &= \lim_{h \rightarrow 0^-} \frac{0 - |h|}{h} \\ &= \lim_{h \rightarrow 0^-} \frac{-h}{h} \quad (h < 0 \Rightarrow |h| = -h) \\ &= -1 \end{aligned}$$

Consider RHD at  $x = 1$

$$\begin{aligned} \lim_{h \rightarrow 0^+} \frac{f(1+h) - f(1)}{h} &= \lim_{h \rightarrow 0^+} \frac{f|1+h-1| - |1-1|}{h} \\ &= \lim_{h \rightarrow 0^+} \frac{|h| - 0}{h} \\ &= \lim_{h \rightarrow 0^+} \frac{h}{h} \quad (h > 0 \Rightarrow |h| = h) \\ &= 1 \end{aligned}$$

Since LHD and RHD at  $x = 1$  are not equal,

Therefore,  $f$  is not differentiable at  $x = 1$ .

### Question 10:

Prove that the greatest integer function defined by  $f(x) = [x], 0 < x < 3$  is not differentiable at  $x = 1$  and  $x = 2$ .

### Solution:

Given,  $f(x) = [x], 0 < x < 3$

It is known that a function  $f$  is differentiable at a point  $x = c$  in its domain if both

$$\lim_{h \rightarrow 0^-} \frac{f(c) - f(c-h)}{h} \quad \text{and} \quad \lim_{h \rightarrow 0^+} \frac{f(c+h) - f(c)}{h} \quad \text{are finite and equal.}$$

At  $x = 1$ ,

Consider the LHD at  $x = 1$

$$\begin{aligned} \lim_{h \rightarrow 0^-} \frac{f(1) - f(1-h)}{h} &= \lim_{h \rightarrow 0^-} \frac{[1] - [1-h]}{h} \\ &= \lim_{h \rightarrow 0^-} \frac{1-0}{h} \\ &= \lim_{h \rightarrow 0^-} \frac{1}{h} \\ &= \infty \end{aligned}$$

Consider RHD at  $x = 1$

$$\begin{aligned} \lim_{h \rightarrow 0^+} \frac{f(1+h) - f(1)}{h} &= \lim_{h \rightarrow 0^+} \frac{[1+h] - [1]}{h} \\ &= \lim_{h \rightarrow 0^+} \frac{1-1}{h} \\ &= \lim_{h \rightarrow 0^+} 0 \\ &= 0 \end{aligned}$$

Since LHD and RHD at  $x = 1$  are not equal,

Hence,  $f$  is not differentiable at  $x = 1$ .

To check the differentiability of the given function at  $x = 2$ ,

Consider LHD at  $x = 2$

$$\begin{aligned} \lim_{h \rightarrow 0^-} \frac{f(2) - f(2-h)}{h} &= \lim_{h \rightarrow 0^-} \frac{[2] - [2-h]}{h} \\ &= \lim_{h \rightarrow 0^-} \frac{2-1}{h} \\ &= \lim_{h \rightarrow 0^-} \frac{1}{h} \\ &= \infty \end{aligned}$$

Now, consider RHD at  $x = 2$



$$\begin{aligned}\lim_{h \rightarrow 0^+} \frac{f(2+h) - f(2)}{h} &= \lim_{h \rightarrow 0^+} \frac{[2+h] - [2]}{h} \\ &= \lim_{h \rightarrow 0^+} \frac{2-2}{h} \\ &= \lim_{h \rightarrow 0^+} 0 \\ &= 0\end{aligned}$$

Since, LHD and RHD at  $x = 2$  are not equal.

Hence,  $f$  is not differentiable at  $x = 2$ .



MillionStars edu  
Think, Learn & Practice

## EXERCISE 5.3

### Question 1:

Find  $\frac{dy}{dx}$  :  $2x + 3y = \sin x$

### Solution:

Given,  $2x + 3y = \sin x$

Differentiating with respect to  $x$ , we get

$$\frac{d}{dy}(2x + 3y) = \frac{d}{dx}(\sin x)$$

$$\Rightarrow \frac{d}{dx}(2x) + \frac{d}{dx}(3y) = \cos x$$

$$\Rightarrow 2 + 3 \frac{dy}{dx} = \cos x$$

$$\Rightarrow 3 \frac{dy}{dx} = \cos x - 2$$

$$\therefore \frac{dy}{dx} = \frac{\cos x - 2}{3}$$

### Question 2:

Find  $\frac{dy}{dx}$  :  $2x + 3y = \sin y$

### Solution:

Given,  $2x + 3y = \sin y$

Differentiating with respect to  $x$ , we get

$$\frac{d}{dx}(2x) + \frac{d}{dx}(3y) = \frac{d}{dy}(\sin y)$$

$$\Rightarrow 2 + 3 \frac{dy}{dx} = \cos y \frac{dy}{dx} \quad [\text{By using chain rule}]$$

$$\Rightarrow 2 = (\cos y - 3) \frac{dy}{dx}$$

$$\therefore \frac{dy}{dx} = \frac{2}{\cos y - 3}$$

### Question 3:

Find  $\frac{dy}{dx}$  :  $ax + by^2 = \cos y$



### Solution:

Given,  $ax + by^2 = \cos y$

Differentiating with respect to  $x$ , we get

$$\frac{d}{dx}(ax) + \frac{d}{dx}(by^2) = \frac{d}{dx}(\cos y)$$

$$\Rightarrow a + b \frac{d}{dx}(y^2) = \frac{d}{dx}(\cos y) \quad \dots(1)$$

$$\frac{d}{dx}(y^2) = 2y \frac{dy}{dx} \text{ and } \frac{d}{dx}(\cos y) = -\sin y \frac{dy}{dx} \quad \dots(2)$$

From (1) and (2), we obtain

$$a + b \times 2y \frac{dy}{dx} = -\sin y \frac{dy}{dx}$$

$$\Rightarrow (2by + \sin y) \frac{dy}{dx} = -a$$

$$\therefore \frac{dy}{dx} = \frac{-a}{2by + \sin y}$$

### Question 4:

Find  $\frac{dy}{dx}$ :  $xy + y^2 = \tan x + y$

### Solution:

Given,  $xy + y^2 = \tan x + y$

Differentiating with respect to  $x$ , we get

$$\frac{d}{dx}(xy + y^2) = \frac{d}{dx}(\tan x + y)$$

$$\Rightarrow \frac{d}{dx}(xy) + \frac{d}{dx}(y^2) = \frac{d}{dx}(\tan x) + \frac{dy}{dx}$$

$$\Rightarrow \left[ y \cdot \frac{d}{dx}(x) + x \cdot \frac{dy}{dx} \right] + 2y \frac{dy}{dx} = \sec^2 x + \frac{dy}{dx} \quad \text{[using product rule and chain rule]}$$

$$\Rightarrow y \cdot 1 + x \frac{dy}{dx} + 2y \frac{dy}{dx} = \sec^2 x + \frac{dy}{dx} \Rightarrow (x + 2y - 1) \frac{dy}{dx} = \sec^2 x - y$$

$$\therefore \frac{dy}{dx} = \frac{\sec^2 x - y}{(x + 2y - 1)}$$

### Question 5:

Find  $\frac{dy}{dx}$ :  $x^2 + xy + y^2 = 100$

### Solution:

Given,  $x^2 + xy + y^2 = 100$

Differentiating with respect to  $x$ , we get

$$\frac{d}{dx}(x^2 + xy + y^2) = \frac{d}{dx}(100)$$

$$\Rightarrow \frac{d}{dx}(x^2) + \frac{d}{dx}(xy) + \frac{d}{dx}(y^2) = 0$$

$$\Rightarrow 2x + \left[ y \cdot \frac{d}{dx}(x) + x \cdot \frac{dy}{dx} \right] + 2y \frac{dy}{dx} = 0$$

$$\Rightarrow 2x + y \cdot 1 + x \cdot \frac{dy}{dx} + 2y \frac{dy}{dx} = 0$$

$$\Rightarrow 2x + y + (x + 2y) \frac{dy}{dx} = 0$$

$$\therefore \frac{dy}{dx} = -\frac{2x + y}{x + 2y}$$

### Question 6:

Find  $\frac{dy}{dx}$ :  $x^3 + x^2y + xy^2 + y^3 = 81$

### Solution:

Given,  $x^3 + x^2y + xy^2 + y^3 = 81$

Differentiating with respect to  $x$ , we get

$$\frac{d}{dx}(x^3 + x^2y + xy^2 + y^3) = \frac{d}{dx}(81)$$

$$\Rightarrow \frac{d}{dx}(x^3) + \frac{d}{dx}(x^2y) + \frac{d}{dx}(xy^2) + \frac{d}{dx}(y^3) = 0$$

$$\Rightarrow 3x^2 + \left[ y \frac{d}{dx}(x^2) + x^2 \frac{dy}{dx} \right] + \left[ y^2 \frac{d}{dx}(x) + x \frac{d}{dx}(y^2) \right] + 3y^2 \frac{dy}{dx} = 0$$

$$\Rightarrow 3x^2 + \left[ y \cdot 2x + x^2 \frac{dy}{dx} \right] + \left[ y^2 \cdot 1 + x \cdot 2y \cdot \frac{dy}{dx} \right] + 3y^2 \frac{dy}{dx} = 0$$

$$\Rightarrow (x^2 + 2xy + 3y^2) \frac{dy}{dx} + (3x^2 + 2xy + y^2) = 0$$

$$\therefore \frac{dy}{dx} = -\frac{(3x^2 + 2xy + y^2)}{(x^2 + 2xy + 3y^2)}$$

### Question 7:

Find  $\frac{dx}{dy}$ :  $\sin^2 y + \cos xy = \pi$

### Solution:

Given,  $\sin^2 y + \cos xy = \pi$

Differentiating with respect to  $x$ , we get

$$\begin{aligned}\frac{d}{dx}(\sin^2 y + \cos xy) &= \frac{d}{dx}(\pi) \\ \Rightarrow \frac{d}{dx}(\sin^2 y) + \frac{d}{dx}(\cos xy) &= 0 \quad \dots(1)\end{aligned}$$

Using chain rule, we obtain

$$\frac{d}{dx}(\sin^2 y) = 2 \sin y \frac{d}{dx}(\sin y) = 2 \sin y \cos y \frac{dy}{dx} \quad \dots(2)$$

$$\begin{aligned}\frac{d}{dx}(\cos xy) &= -\sin xy \frac{d}{dx}(xy) = -\sin xy \left[ y \frac{d}{dx}(x) + x \frac{dy}{dx} \right] \\ &= -\sin xy \left[ y \cdot 1 + x \frac{dy}{dx} \right] = -y \sin xy - x \sin xy \frac{dy}{dx} \quad \dots(3)\end{aligned}$$

From (1), (2) and (3), we obtain

$$2 \sin y \cos y \frac{dy}{dx} + \left( -y \sin xy - x \sin xy \frac{dy}{dx} \right) = 0$$

$$\Rightarrow (2 \sin y \cos y - x \sin xy) \frac{dy}{dx} = y \sin xy$$

$$\Rightarrow (\sin 2y - x \sin xy) \frac{dx}{dy} = y \sin xy$$

$$\therefore \frac{dx}{dy} = \frac{y \sin xy}{\sin 2y - x \sin xy}$$

### Question 8:

Find  $\frac{dy}{dx}$ :  $\sin^2 x + \cos^2 y = 1$

### Solution:

Given,  $\sin^2 x + \cos^2 y = 1$

Differentiating with respect to  $x$ , we get

$$\begin{aligned}\frac{d}{dx}(\sin^2 x + \cos^2 y) &= \frac{d}{dx}(1) \\ \Rightarrow \frac{d}{dx}(\sin^2 x) + \frac{d}{dx}(\cos^2 y) &= 0 \\ \Rightarrow 2 \sin x \cdot \frac{d}{dx}(\sin x) + 2 \cos y \cdot \frac{d}{dx}(\cos y) &= 0 \\ \Rightarrow 2 \sin x \cos x + 2 \cos y(-\sin y) \cdot \frac{dy}{dx} &= 0 \\ \Rightarrow \sin 2x - \sin 2y \frac{dy}{dx} &= 0 \\ \therefore \frac{dy}{dx} &= \frac{\sin 2x}{\sin 2y}\end{aligned}$$

### Question 9:

Find  $\frac{dy}{dx}$  :  $y = \sin^{-1}\left(\frac{2x}{1+x^2}\right)$

### Solution:

Given,

$$\begin{aligned}y &= \sin^{-1}\left(\frac{2x}{1+x^2}\right) \\ \Rightarrow \sin y &= \frac{2x}{1+x^2}\end{aligned}$$

Differentiating with respect to  $x$ , we get

$$\begin{aligned}\frac{d}{dx}(\sin y) &= \frac{d}{dx}\left(\frac{2x}{1+x^2}\right) \quad \dots(1) \\ \Rightarrow \cos y \frac{dy}{dx} &= \frac{d}{dx}\left(\frac{2x}{1+x^2}\right)\end{aligned}$$

The function  $\frac{2x}{1+x^2}$ , is of the form of  $\frac{u}{v}$

By quotient rule, we get

$$\begin{aligned}\frac{d}{dx}\left(\frac{2x}{1+x^2}\right) &= \frac{(1+x^2)\frac{d}{dx}(2x) - 2x\frac{d}{dx}(1+x^2)}{(1+x^2)^2} \\ &= \frac{(1+x^2) \cdot 2 - 2x \cdot [0+2x]}{(1+x^2)^2} \\ &= \frac{2+2x^2-4x^2}{(1+x^2)^2} \\ &= \frac{2(1-x^2)}{(1+x^2)^2}\end{aligned}$$

Also,  $\sin y = \frac{2x}{1+x^2}$

$$\begin{aligned}\cos y &= \sqrt{1-\sin^2 y} = \sqrt{1-\left(\frac{2x}{1+x^2}\right)^2} \\ &= \sqrt{\frac{(1+x^2)^2-4x^2}{(1+x^2)^2}} \\ &= \sqrt{\frac{(1-x^2)^2}{(1+x^2)^2}} \\ &= \frac{1-x^2}{1+x^2}\end{aligned}$$

From (1), (2) and (3), we get

$$\begin{aligned}\frac{1-x^2}{1+x^2} \times \frac{dy}{dx} &= \frac{2(1-x^2)}{(1+x^2)^2} \\ \Rightarrow \frac{dy}{dx} &= \frac{2}{1+x^2}\end{aligned}$$

### Question 10:

Find  $\frac{dy}{dx}$ :  $y = \tan^{-1}\left(\frac{3x-x^3}{1-3x^2}\right)$ ,  $-\frac{1}{\sqrt{3}} < x < \frac{1}{\sqrt{3}}$

**Solution:**

Given,  $y = \tan^{-1} \left( \frac{3x - x^3}{1 - 3x^2} \right)$

$$\Rightarrow \tan y = \left( \frac{3x - x^3}{1 - 3x^2} \right) \quad \dots(1)$$

Since, we know that

$$\Rightarrow \tan y = \left( \frac{3 \tan \frac{y}{3} - \tan^3 \frac{y}{3}}{1 - 3 \tan^2 \frac{y}{3}} \right) \quad (2)$$

Comparing (1) and (2) we get,

$$x = \tan \frac{y}{3}$$

Differentiating with respect to  $x$ , we get

$$\frac{d}{dx}(x) = \frac{d}{dx} \left( \tan \frac{y}{3} \right)$$

$$\Rightarrow 1 = \sec^2 \frac{y}{3} \cdot \frac{d}{dx} \left( \frac{y}{3} \right)$$

$$\Rightarrow 1 = \sec^2 \frac{y}{3} \cdot \frac{1}{3} \cdot \frac{dy}{dx}$$

$$\Rightarrow \frac{dy}{dx} = \frac{3}{\sec^2 \frac{y}{3}} = \frac{3}{1 + \tan^2 \frac{y}{3}}$$

$$\therefore \frac{dy}{dx} = \frac{3}{1 + x^2}$$

**Question 11:**

Find  $\frac{dy}{dx}$ ;  $y = \cos^{-1} \left( \frac{1 - x^2}{1 + x^2} \right)$ ,  $0 < x < 1$

**Solution:**

Given,  $y = \cos^{-1} \left( \frac{1 - x^2}{1 + x^2} \right)$

$$\Rightarrow \cos y = \left( \frac{1-x^2}{1+x^2} \right)$$

$$\Rightarrow \frac{1 - \tan^2 \frac{y}{2}}{1 + \tan^2 \frac{y}{2}} = \frac{1-x^2}{1+x^2}$$

Comparing LHS and RHS, we get

$$\tan \frac{y}{2} = x$$

Differentiating with respect to  $x$ , we get

$$\sec^2 \frac{y}{2} \cdot \frac{d}{dx} \left( \frac{y}{2} \right) = \frac{d}{dx} (x)$$

$$\Rightarrow \sec^2 \frac{y}{2} \times \frac{1}{2} \frac{dy}{dx} = 1$$

$$\Rightarrow \frac{dy}{dx} = \frac{2}{\sec^2 \frac{y}{2}}$$

$$\Rightarrow \frac{dy}{dx} = \frac{2}{1 + \tan^2 \frac{y}{2}}$$

$$\therefore \frac{dy}{dx} = \frac{2}{1+x^2}$$

### Question 12:

Find  $\frac{dy}{dx}$  :  $y = \sin^{-1} \left( \frac{1-x^2}{1+x^2} \right)$ ,  $0 < x < 1$

**Solution:**

Given,  $y = \sin^{-1} \left( \frac{1-x^2}{1+x^2} \right)$

$$y = \sin^{-1} \left( \frac{1-x^2}{1+x^2} \right)$$

$$\Rightarrow \sin y = \frac{1-x^2}{1+x^2}$$

Differentiating with respect to  $x$ , we get

$$\frac{d}{dx} (\sin y) = \frac{d}{dx} \left( \frac{1-x^2}{1+x^2} \right) \quad \dots(1)$$

Using chain rule, we get

$$\frac{d}{dx}(\sin y) = \cos y \cdot \frac{dy}{dx}$$

$$\begin{aligned}\cos y &= \sqrt{1 - \sin^2 y} = \sqrt{1 - \left(\frac{1-x^2}{1+x^2}\right)^2} \\ &= \sqrt{\frac{(1+x^2)^2 - (1-x^2)^2}{(1+x^2)^2}} = \sqrt{\frac{4x^2}{(1+x^2)^2}} = \frac{2x}{1+x^2}\end{aligned}$$

Therefore,

$$\frac{d}{dx}(\sin y) = \frac{2x}{1+x^2} \frac{dy}{dx} \quad \dots(2)$$

$$\begin{aligned}\frac{d}{dx}\left(\frac{1-x^2}{1+x^2}\right) &= \frac{(1+x^2) \cdot \frac{d}{dx}(1-x^2) - (1-x^2) \cdot \frac{d}{dx}(1+x^2)}{(1+x^2)^2} && \text{[using quotient rule]} \\ &= \frac{(1+x^2)(-2x) - (1-x^2)(2x)}{(1+x^2)^2} \\ &= \frac{-2x - 2x^3 - 2x + 2x^3}{(1+x^2)^2} \\ &= \frac{-4x}{(1+x^2)^2} && \dots(3)\end{aligned}$$

From equation (1), (2) and (3), we get

$$\begin{aligned}\frac{2x}{1+x^2} \frac{dy}{dx} &= \frac{-4x}{(1+x^2)^2} \\ \Rightarrow \frac{dy}{dx} &= \frac{-2}{1+x^2}\end{aligned}$$

### Question 13:

Find  $\frac{dy}{dx}$ :  $y = \cos^{-1}\left(\frac{2x}{1+x^2}\right)$ ,  $-1 < x < 1$

**Solution:**

Given,  $y = \cos^{-1}\left(\frac{2x}{1+x^2}\right)$

$$y = \cos^{-1}\left(\frac{2x}{1+x^2}\right)$$

$$\cos y = \left(\frac{2x}{1+x^2}\right)$$



Differentiating with respect to  $x$ , we get

$$\begin{aligned}\frac{d}{dx}(\cos y) &= \frac{d}{dx}\left(\frac{2x}{1+x^2}\right) \\ \Rightarrow -\sin y \cdot \frac{dy}{dx} &= \frac{(1+x^2) \cdot \frac{d}{dx}(2x) - 2x \cdot \frac{d}{dx}(1+x^2)}{(1+x^2)^2} \\ \Rightarrow -\sqrt{1-\cos^2 y} \frac{dy}{dx} &= \frac{(1+x^2) \times 2 - 2x \times 2x}{(1+x^2)^2} \\ \Rightarrow \left[ \sqrt{1-\left(\frac{2x}{1+x^2}\right)^2} \right] \frac{dy}{dx} &= -\left[ \frac{2(1-x^2)}{(1+x^2)^2} \right] \\ \Rightarrow \sqrt{\frac{(1+x^2)^2 - 4x^2}{(1+x^2)^2}} \cdot \frac{dy}{dx} &= \frac{-2(1-x^2)}{(1+x^2)^2} \\ \Rightarrow \sqrt{\frac{(1-x^2)^2}{(1+x^2)^2}} \frac{dy}{dx} &= \frac{-2(1-x^2)}{(1+x^2)^2} \\ \Rightarrow \frac{1-x^2}{1+x^2} \cdot \frac{dy}{dx} &= \frac{-2(1-x^2)}{(1+x^2)^2} \\ \Rightarrow \frac{dy}{dx} &= \frac{-2}{1+x^2}\end{aligned}$$

#### Question 14:

Find  $\frac{dy}{dx}$ :  $y = \sin^{-1}(2x\sqrt{1-x^2})$ ,  $-\frac{1}{\sqrt{2}} < x < \frac{1}{\sqrt{2}}$

**Solution:**

Given,  $y = \sin^{-1}(2x\sqrt{1-x^2})$

$$y = \sin^{-1}(2x\sqrt{1-x^2})$$

$$\Rightarrow \sin y = (2x\sqrt{1-x^2})$$

Differentiating with respect to  $x$ , we get

$$\begin{aligned}\cos y \cdot \frac{dy}{dx} &= 2 \left[ x \frac{d}{dx} (\sqrt{1-x^2}) + \sqrt{1-x^2} \frac{dx}{dx} \right] \\ \Rightarrow \sqrt{1-\sin^2 y} \frac{dy}{dx} &= 2 \left[ \frac{x}{2} \cdot \frac{-2x}{\sqrt{1-x^2}} + \sqrt{1-x^2} \right] \\ \Rightarrow \sqrt{1-(2x\sqrt{1-x^2})^2} \cdot \frac{dy}{dx} &= 2 \left[ \frac{-x^2+1-x^2}{\sqrt{1-x^2}} \right] \\ \Rightarrow \sqrt{1-4x^2(1-x^2)} \frac{dy}{dx} &= 2 \left[ \frac{1-2x^2}{\sqrt{1-x^2}} \right] \\ \Rightarrow \sqrt{(1-2x^2)^2} \frac{dy}{dx} &= 2 \left[ \frac{1-2x^2}{\sqrt{1-x^2}} \right] \\ \Rightarrow (1-2x^2) \frac{dy}{dx} &= 2 \left[ \frac{1-2x^2}{\sqrt{1-x^2}} \right] \\ \Rightarrow \frac{dy}{dx} &= \frac{2}{\sqrt{1-x^2}}\end{aligned}$$

### Question 15:

Find  $\frac{dy}{dx}$ :  $y = \sec^{-1} \left( \frac{1}{2x^2-1} \right)$ ,  $0 < x < \frac{1}{\sqrt{2}}$

### Solution:

Given,  $y = \sec^{-1} \left( \frac{1}{2x^2-1} \right)$

$$\Rightarrow y = \sec^{-1} \left( \frac{1}{2x^2 - 1} \right)$$

$$\Rightarrow \sec y = \left( \frac{1}{2x^2 - 1} \right)$$

$$\Rightarrow \cos y = 2x^2 - 1$$

$$\Rightarrow 2x^2 = 1 + \cos y$$

$$\Rightarrow 2x^2 = 2 \cos^2 \frac{y}{2}$$

$$\Rightarrow x = \cos \frac{y}{2}$$

Differentiating with respect to  $x$ , we get

$$\frac{d}{dx}(x) = \frac{d}{dx} \left( \cos \frac{y}{2} \right)$$

$$\Rightarrow 1 = \sin \frac{y}{2} \cdot \frac{d}{dx} \left( \frac{y}{2} \right)$$

$$\Rightarrow \frac{-1}{\sin \frac{y}{2}} = \frac{1}{2} \frac{dy}{dx}$$

$$\Rightarrow \frac{dy}{dx} = \frac{-2}{\sin \frac{y}{2}}$$

$$\Rightarrow \frac{dy}{dx} = \frac{-2}{\sqrt{1 - \cos^2 \frac{y}{2}}}$$

$$\Rightarrow \frac{dy}{dx} = \frac{-2}{\sqrt{1 - x^2}}$$



## EXERCISE 5.4

### Question 1:

Differentiating the following wrt  $x$ :  $\frac{e^x}{\sin x}$

#### Solution:

Let  $y = \frac{e^x}{\sin x}$

By using the quotient rule, we get

$$\begin{aligned}\frac{dy}{dx} &= \frac{\sin x \frac{d}{dx}(e^x) - e^x \frac{d}{dx}(\sin x)}{\sin^2 x} \\ &= \frac{\sin x \cdot (e^x) - e^x \cdot (\cos x)}{\sin^2 x} \\ &= \frac{e^x (\sin x - \cos x)}{\sin^2 x}\end{aligned}$$

### Question 2:

Differentiating the following  $e^{\sin^{-1} x}$

#### Solution:

Let  $y = e^{\sin^{-1} x}$

By using the quotient rule, we get

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx}(e^{\sin^{-1} x}) \\ &= e^{\sin^{-1} x} \cdot \frac{d}{dx}(\sin^{-1} x) \\ &= e^{\sin^{-1} x} \cdot \frac{1}{\sqrt{1-x^2}} \\ &= \frac{e^{\sin^{-1} x}}{\sqrt{1-x^2}} \\ &= \frac{e^{\sin^{-1} x}}{\sqrt{1-x^2}}, x \in (-1, 1)\end{aligned}$$

### Question 3:

Differentiating the following wrt  $x$ :  $e^{x^3}$

### Solution:

Let  $y = e^{x^3}$

By using the quotient rule, we get

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx}(e^{x^3}) \\ &= e^{x^3} \cdot \frac{d}{dx}(x^3) \\ &= e^{x^3} \cdot 3x^2 \\ &= 3x^2 e^{x^3}\end{aligned}$$

### Question 4:

Differentiate the following wrt  $x: \sin(\tan^{-1} e^{-x})$

### Solution:

Let  $y = \sin(\tan^{-1} e^{-x})$

By using the chain rule, we get

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx}[\sin(\tan^{-1} e^{-x})] \\ &= \cos(\tan^{-1} e^{-x}) \cdot \frac{d}{dx}(\tan^{-1} e^{-x}) \\ &= \cos(\tan^{-1} e^{-x}) \cdot \frac{1}{1+(e^{-x})^2} \cdot \frac{d}{dx}(e^{-x}) \\ &= \frac{\cos(\tan^{-1} e^{-x})}{1+e^{-2x}} \cdot e^{-x} \cdot \frac{d}{dx}(-x) \\ &= \frac{e^{-x} \cos(\tan^{-1} e^{-x})}{1+e^{-2x}} \times (-1) \\ &= \frac{-e^{-x} \cos(\tan^{-1} e^{-x})}{1+e^{-2x}}\end{aligned}$$

### Question 5:

Differentiate the following wrt  $x: \log(\cos e^x)$

### Solution:

Let  $y = \log(\cos e^x)$

By using the chain rule, we get

$$\begin{aligned}
 \frac{dy}{dx} &= \frac{d}{dx} \left[ \log(\cos e^x) \right] \\
 &= \frac{1}{\cos e^x} \cdot \frac{d}{dx} (\cos e^x) \\
 &= \frac{1}{\cos e^x} \cdot (-\sin e^x) \cdot \frac{d}{dx} (e^x) \\
 &= \frac{-\sin e^x}{\cos e^x} \cdot e^x \\
 &= -e^x \tan e^x, e^x \neq (2n+1)\frac{\pi}{2}, n \in \mathbb{N}
 \end{aligned}$$

### Question 6:

Differentiate the following wrt  $x$ :  $e^x + e^{x^2} + \dots + e^{x^5}$

### Solution:

$$\frac{d}{dx} (e^x + e^{x^2} + \dots + e^{x^5})$$

Differentiating wrt  $x$ , we get

$$\begin{aligned}
 \frac{d}{dx} (e^x + e^{x^2} + \dots + e^{x^5}) &= \frac{d}{dx} (e^x) + \frac{d}{dx} (e^{x^2}) + \frac{d}{dx} (e^{x^3}) + \frac{d}{dx} (e^{x^4}) + \frac{d}{dx} (e^{x^5}) \\
 &= e^x + \left[ e^{x^2} \times \frac{d}{dx} (x^2) \right] + \left[ e^{x^3} \times \frac{d}{dx} (x^3) \right] + \left[ e^{x^4} \times \frac{d}{dx} (x^4) \right] + \left[ e^{x^5} \times \frac{d}{dx} (x^5) \right] \\
 &= e^x + (e^{x^2} \times 2x) + (e^{x^3} \times 3x^2) + (e^{x^4} \times 4x^3) + (e^{x^5} \times 5x^4) \\
 &= e^x + 2xe^{x^2} + 3x^2e^{x^3} + 4x^3e^{x^4} + 5x^4e^{x^5}
 \end{aligned}$$

### Question 7:

Differentiating the following wrt  $x$ :  $\sqrt{e^{\sqrt{x}}}$ ,  $x > 0$

### Solution:

$$\text{Let } y = \sqrt{e^{\sqrt{x}}}$$

$$\text{Then, } y^2 = e^{\sqrt{x}}$$

Differentiating wrt  $x$ , we get

$$y^2 = e^{\sqrt{x}}$$

$$\begin{aligned}\frac{d}{dx}(y^2) &= \frac{d}{dx}(e^{\sqrt{x}}) \\ \Rightarrow 2y \frac{dy}{dx} &= e^{\sqrt{x}} \frac{d}{dx}(\sqrt{x}) \\ \Rightarrow 2y \frac{dy}{dx} &= e^{\sqrt{x}} \frac{1}{2} \cdot \frac{1}{\sqrt{x}} \\ \Rightarrow \frac{dy}{dx} &= \frac{e^{\sqrt{x}}}{4y\sqrt{x}} \\ \Rightarrow \frac{dy}{dx} &= \frac{e^{\sqrt{x}}}{4\sqrt{e^{\sqrt{x}}}\sqrt{x}} \\ \Rightarrow \frac{dy}{dx} &= \frac{e^{\sqrt{x}}}{4\sqrt{x}e^{\sqrt{x}}}, x > 0\end{aligned}$$

### Question 8:

Differentiating the following wrt  $x: \log(\log x), x > 1$

#### Solution:

Let  $y = \log(\log x)$

By using the chain rule, we get

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx}[\log(\log x)] \\ &= \frac{1}{\log x} \cdot \frac{d}{dx}(\log x) \\ &= \frac{1}{\log x} \cdot \frac{1}{x} \\ &= \frac{1}{x \log x}, x > 1\end{aligned}$$

### Question 9:

Differentiating the following wrt  $x: \frac{\cos x}{\log x}, x > 0$

#### Solution:

Let  $y = \frac{\cos x}{\log x}$

By using the quotient rule, we get

$$\begin{aligned}\frac{dy}{dx} &= \frac{\frac{d}{dx}(\cos x) \cdot \log x - \cos x \cdot \frac{d}{dx}(\log x)}{(\log x)^2} \\ &= \frac{-\sin x \log x - \cos x \cdot \frac{1}{x}}{(\log x)^2} \\ &= -\left[ \frac{x \log x \cdot \sin x + \cos x}{x(\log x)^2} \right], x > 0\end{aligned}$$

### Question 10:

Differentiate the following wrt  $x$ :  $\cos(\log x + e^x)$ ,  $x > 0$

### Solution:

Let  $y = \cos(\log x + e^x)$

By using the chain rule, we get

$$\begin{aligned}\frac{dy}{dx} &= -\sin[\log x + e^x] \cdot \frac{d}{dx}(\log x + e^x) \\ &= -\sin(\log x + e^x) \cdot \left[ \frac{d}{dx}(\log x) + \frac{d}{dx}(e^x) \right] \\ &= -\sin(\log x + e^x) \cdot \left( \frac{1}{x} + e^x \right) \\ &= -\left( \frac{1}{x} + e^x \right) \sin(\log x + e^x), x > 0\end{aligned}$$



## EXERCISE 5.5

### Question 1:

Differentiate the function with respect to  $x$ :  $\cos x \cdot \cos 2x \cdot \cos 3x$

### Solution:

Let  $y = \cos x \cdot \cos 2x \cdot \cos 3x$

Taking logarithm on both the sides, we obtain

$$\log y = \log(\cos x \cdot \cos 2x \cdot \cos 3x)$$

$$\Rightarrow \log y = \log(\cos x) + \log(\cos 2x) + \log(\cos 3x)$$

Differentiating both sides with respect to  $x$ , we obtain

$$\frac{1}{y} \frac{dy}{dx} = \frac{1}{\cos x} \cdot \frac{d}{dx}(\cos x) + \frac{1}{\cos 2x} \cdot \frac{d}{dx}(\cos 2x) + \frac{1}{\cos 3x} \cdot \frac{d}{dx}(\cos 3x)$$

$$\Rightarrow \frac{dy}{dx} = y \left[ -\frac{\sin x}{\cos x} - \frac{\sin 2x}{\cos 2x} \cdot \frac{d}{dx}(2x) - \frac{\sin 3x}{\cos 3x} \cdot \frac{d}{dx}(3x) \right]$$

$$\therefore \frac{dy}{dx} = -\cos x \cdot \cos 2x \cdot \cos 3x [\tan x + 2 \tan 2x + 3 \tan 3x]$$

### Question 2:

Differentiate the function with respect to  $x$ :  $\sqrt{\frac{(x-1)(x-2)}{(x-3)(x-4)(x-5)}}$

### Solution:

$$\text{Let } y = \sqrt{\frac{(x-1)(x-2)}{(x-3)(x-4)(x-5)}}$$

Taking logarithm on both the sides, we obtain

$$\log y = \log \sqrt{\frac{(x-1)(x-2)}{(x-3)(x-4)(x-5)}}$$

$$\Rightarrow \log y = \frac{1}{2} \log \left[ \frac{(x-1)(x-2)}{(x-3)(x-4)(x-5)} \right]$$

$$\Rightarrow \log y = \frac{1}{2} [\log \{(x-1)(x-2)\} - \log \{(x-3)(x-4)(x-5)\}]$$

$$\Rightarrow \log y = \frac{1}{2} [\log(x-1) + \log(x-2) - \log(x-3) - \log(x-4) - \log(x-5)]$$

Differentiating both sides with respect to  $x$ , we obtain

$$\begin{aligned}\frac{1}{y} \frac{dy}{dx} &= \frac{1}{2} \left[ \frac{1}{x-1} \cdot \frac{d}{dx}(x-1) + \frac{1}{x-2} \cdot \frac{d}{dx}(x-2) - \frac{1}{x-3} \cdot \frac{d}{dx}(x-3) \right. \\ &\quad \left. - \frac{1}{x-4} \cdot \frac{d}{dx}(x-4) - \frac{1}{x-5} \cdot \frac{d}{dx}(x-5) \right] \\ \Rightarrow \frac{dy}{dx} &= \frac{y}{2} \left[ \frac{1}{x-1} + \frac{1}{x-2} - \frac{1}{x-3} - \frac{1}{x-4} - \frac{1}{x-5} \right] \\ \therefore \frac{dy}{dx} &= \frac{1}{2} \sqrt{\frac{(x-1)(x-2)}{(x-3)(x-4)(x-5)}} \left[ \frac{1}{x-1} + \frac{1}{x-2} - \frac{1}{x-3} - \frac{1}{x-4} - \frac{1}{x-5} \right]\end{aligned}$$

### Question 3:

Differentiate the function with respect to  $x$ :  $(\log x)^{\cos x}$

**Solution:**

Let  $y = (\log x)^{\cos x}$

Taking logarithm on both the sides, we obtain

$$\log y = \cos x \cdot \log(\log x)$$

Differentiating both sides with respect to  $x$ , we obtain

$$\begin{aligned}\frac{1}{y} \cdot \frac{dy}{dx} &= \frac{d}{dx}(\cos x) \cdot \log(\log x) + \cos x \cdot \frac{d}{dx}[\log(\log x)] \\ \Rightarrow \frac{1}{y} \cdot \frac{dy}{dx} &= -\sin x \log(\log x) + \cos x \cdot \frac{1}{\log x} \cdot \frac{d}{dx}(\log x) \\ \Rightarrow \frac{dy}{dx} &= y \left[ -\sin x \log(\log x) + \frac{\cos x}{\log x} \cdot \frac{1}{x} \right] \\ \therefore \frac{dy}{dx} &= (\log x)^{\cos x} \left[ \frac{\cos x}{x \log x} - \sin x \log(\log x) \right]\end{aligned}$$

### Question 4:

Differentiate the function with respect to  $x$ :  $x^x - 2^{\sin x}$

### Solution:

Let  $y = x^x - 2^{\sin x}$

Also, let  $x^x = u$  and  $2^{\sin x} = v$

$\therefore y = u - v$

$$\Rightarrow \frac{dy}{dx} = \frac{du}{dx} - \frac{dv}{dx}$$

$u = x^x$

Taking logarithm on both the sides, we obtain

$$\log u = x \log x$$

Differentiating both sides with respect to  $x$ , we obtain

$$\frac{1}{u} \cdot \frac{du}{dx} = \left[ \frac{d}{dx}(x) \times \log x + x \times \frac{d}{dx}(\log x) \right]$$

$$\Rightarrow \frac{du}{dx} = u \left[ 1 \times \log x + x \times \frac{1}{x} \right]$$

$$\Rightarrow \frac{du}{dx} = x^x (\log x + 1)$$

$$\Rightarrow \frac{du}{dx} = x^x (1 + \log x)$$

$v = 2^{\sin x}$

Taking logarithm on both the sides, we obtain

$$\log v = \sin x \cdot \log 2$$

Differentiating both sides with respect to  $x$ , we obtain

$$\frac{1}{v} \cdot \frac{dv}{dx} = \log 2 \cdot \frac{d}{dx}(\sin x)$$

$$\Rightarrow \frac{dv}{dx} = v \log 2 \cos x$$

$$\Rightarrow \frac{dv}{dx} = 2^{\sin x} \cos x \log 2$$

$$\therefore \frac{dy}{dx} = x^x (1 + \log x) - 2^{\sin x} \cos x \log 2$$

### Question 5:

Differentiate the function with respect to  $x$ :  $(x+3)^2 \cdot (x+4)^3 \cdot (x+5)^4$

### Solution:

$$\text{Let } y = (x+3)^2 \cdot (x+4)^3 \cdot (x+5)^4$$

Taking logarithm on both the sides, we obtain

$$\begin{aligned}\log y &= \log(x+3)^2 + \log(x+4)^3 + \log(x+5)^4 \\ \Rightarrow \log y &= 2\log(x+3) + 3\log(x+4) + 4\log(x+5)\end{aligned}$$

Differentiating both sides with respect to  $x$ , we obtain

$$\begin{aligned}\frac{1}{y} \cdot \frac{dy}{dx} &= 2 \cdot \frac{1}{x+3} \cdot \frac{d}{dx}(x+3) + 3 \cdot \frac{1}{x+4} \cdot \frac{d}{dx}(x+4) + 4 \cdot \frac{1}{x+5} \cdot \frac{d}{dx}(x+5) \\ \Rightarrow \frac{dy}{dx} &= y \left[ \frac{2}{x+3} + \frac{3}{x+4} + \frac{4}{x+5} \right] \\ \Rightarrow \frac{dy}{dx} &= (x+3)^2 (x+4)^3 (x+5)^4 \cdot \left[ \frac{2}{x+3} + \frac{3}{x+4} + \frac{4}{x+5} \right] \\ \Rightarrow \frac{dy}{dx} &= (x+3)^2 (x+4)^3 (x+5)^4 \cdot \left[ \frac{2(x+4)(x+5) + 3(x+3)(x+5) + 4(x+3)(x+4)}{(x+3)(x+4)(x+5)} \right] \\ \Rightarrow \frac{dy}{dx} &= (x+3)(x+4)^2 (x+5)^3 \cdot \left[ \frac{2(x^2 + 9x + 20) + 3(x^2 + 8x + 15) + 4(x^2 + 7x + 12)}{(x+3)(x+4)(x+5)} \right] \\ \therefore \frac{dy}{dx} &= (x+3)(x+4)^2 (x+5)^3 (9x^2 + 70x + 133)\end{aligned}$$

### Question 6:

Differentiate the function with respect to  $x$ :  $\left(x + \frac{1}{x}\right)^x + x^{\left(1 + \frac{1}{x}\right)}$

### Solution:

$$\text{Let } y = \left(x + \frac{1}{x}\right)^x + x^{\left(1 + \frac{1}{x}\right)}$$

$$\text{Also, let } u = \left(x + \frac{1}{x}\right)^x \text{ and } v = x^{\left(1 + \frac{1}{x}\right)}$$

$$\therefore y = u + v$$

$$\Rightarrow \frac{dy}{dx} = \frac{du}{dx} + \frac{dv}{dx} \quad \dots(1)$$

Then,  $u = \left(x + \frac{1}{x}\right)^x$

Taking logarithm on both the sides, we obtain

$$\Rightarrow \log u = \log \left(x + \frac{1}{x}\right)^x$$

$$\Rightarrow \log u = x \log \left(x + \frac{1}{x}\right)$$

Differentiating both sides with respect to  $x$ , we obtain

$$\frac{1}{u} \cdot \frac{du}{dx} = \frac{d}{dx}(x) \times \log \left(x + \frac{1}{x}\right) + x \times \frac{d}{dx} \left[ \log \left(x + \frac{1}{x}\right) \right]$$

$$\Rightarrow \frac{1}{u} \cdot \frac{du}{dx} = 1 \times \log \left(x + \frac{1}{x}\right) + x \times \frac{1}{\left(x + \frac{1}{x}\right)} \cdot \frac{d}{dx} \left(x + \frac{1}{x}\right)$$

$$\Rightarrow \frac{du}{dx} = u \left[ \log \left(x + \frac{1}{x}\right) + \frac{x}{\left(x + \frac{1}{x}\right)} \times \left(1 - \frac{1}{x^2}\right) \right]$$

$$\Rightarrow \frac{du}{dx} = \left(x + \frac{1}{x}\right)^x \left[ \log \left(x + \frac{1}{x}\right) + \frac{\left(x - \frac{1}{x}\right)}{\left(x + \frac{1}{x}\right)} \right]$$

$$\Rightarrow \frac{du}{dx} = \left(x + \frac{1}{x}\right)^x \left[ \log \left(x + \frac{1}{x}\right) + \frac{x^2 - 1}{x^2 + 1} \right]$$

$$\Rightarrow \frac{du}{dx} = \left(x + \frac{1}{x}\right)^x \left[ \frac{x^2 - 1}{x^2 + 1} + \log \left(x + \frac{1}{x}\right) \right] \quad \dots(2)$$

Now,  $v = x^{\left(1 + \frac{1}{x}\right)}$

Taking logarithm on both the sides, we obtain

$$\Rightarrow \log v = \log \left[ x^{\left(1 + \frac{1}{x}\right)} \right]$$

$$\Rightarrow \log v = \left(1 + \frac{1}{x}\right) \log x$$



Differentiating both sides with respect to  $x$ , we obtain

$$\frac{1}{v} \cdot \frac{dv}{dx} = \left[ \frac{d}{dx} \left( 1 + \frac{1}{x} \right) \right] \times \log x + \left( 1 + \frac{1}{x} \right) \cdot \frac{d}{dx} \log x$$

$$\Rightarrow \frac{1}{v} \cdot \frac{dv}{dx} = \left( -\frac{1}{x^2} \right) \log x + \left( 1 + \frac{1}{x} \right) \cdot \frac{1}{x}$$

$$\Rightarrow \frac{1}{v} \cdot \frac{dv}{dx} = -\frac{\log x}{x^2} + \frac{1}{x} + \frac{1}{x^2}$$

$$\Rightarrow \frac{dv}{dx} = v \left[ \frac{-\log x + x + 1}{x^2} \right]$$

$$\Rightarrow \frac{dv}{dx} = x^{\left(1+\frac{1}{x}\right)} \left[ \frac{-\log x + x + 1}{x^2} \right] \quad \dots(3)$$

Therefore, from (1), (2) and (3);

$$\frac{dy}{dx} = \left( x + \frac{1}{x} \right)^x \left[ \frac{x^2 - 1}{x^2 + 1} + \log \left( x + \frac{1}{x} \right) \right] + x^{\left(1+\frac{1}{x}\right)} \left[ \frac{x + 1 - \log x}{x^2} \right]$$

### Question 7:

Differentiate the function with respect to  $x$ :  $(\log x)^x + x^{\log x}$

### Solution:

Let  $y = (\log x)^x + x^{\log x}$

Also, let  $u = (\log x)^x$  and  $v = x^{\log x}$

$$\therefore y = u + v$$

$$\Rightarrow \frac{dy}{dx} = \frac{du}{dx} + \frac{dv}{dx} \quad \dots(1)$$

Then,  $u = (\log x)^x$

Taking logarithm on both the sides, we obtain

$$\Rightarrow \log u = \log \left[ (\log x)^x \right]$$

$$\Rightarrow \log u = x \log (\log x)$$

Differentiating both sides with respect to  $x$ , we obtain

$$\begin{aligned}
 \frac{1}{u} \cdot \frac{du}{dx} &= \frac{d}{dx}(x) \times \log(\log x) + x \cdot \frac{d}{dx}[\log(\log x)] \\
 \Rightarrow \frac{du}{dx} &= u \left[ 1 \times \log(\log x) + x \cdot \frac{1}{(\log x)} \cdot \frac{d}{dx}(\log x) \right] \\
 \Rightarrow \frac{du}{dx} &= (\log x)^x \left[ \log(\log x) + \frac{x}{(\log x)} \cdot \frac{1}{x} \right] \\
 \Rightarrow \frac{du}{dx} &= (\log x)^x \left[ \log(\log x) + \frac{1}{(\log x)} \right] \\
 \Rightarrow \frac{du}{dx} &= (\log x)^x \left[ \frac{\log(\log x) \cdot \log x + 1}{\log x} \right] \\
 \Rightarrow \frac{du}{dx} &= (\log x)^{x-1} [1 + \log x \cdot \log(\log x)] \quad \dots(2)
 \end{aligned}$$

$$v = x^{\log x}$$

Taking logarithm on both the sides, we obtain

$$\begin{aligned}
 \Rightarrow \log v &= \log(x^{\log x}) \\
 \Rightarrow \log v &= \log x \log x = (\log x)^2
 \end{aligned}$$

Differentiating both sides with respect to  $x$ , we obtain

$$\begin{aligned}
 \frac{1}{v} \cdot \frac{dv}{dx} &= \frac{d}{dx}[(\log x)^2] \\
 \Rightarrow \frac{1}{v} \cdot \frac{dv}{dx} &= 2(\log x) \cdot \frac{d}{dx}(\log x) \\
 \Rightarrow \frac{dv}{dx} &= 2v(\log x) \cdot \frac{1}{x} \\
 \Rightarrow \frac{dv}{dx} &= 2x^{\log x} \frac{\log x}{x} \\
 \Rightarrow \frac{dv}{dx} &= 2x^{\log x - 1} \cdot \log x \quad \dots(3)
 \end{aligned}$$

Therefore, from (1), (2) and (3);

$$\frac{dy}{dx} = (\log x)^{x-1} [1 + \log x \cdot \log(\log x)] + 2x^{\log x - 1} \cdot \log x$$

### Question 8:

Differentiate the function with respect to  $x$ :  $(\sin x)^x + \sin^{-1} \sqrt{x}$

### Solution:

Let  $y = (\sin x)^x + \sin^{-1} \sqrt{x}$

Also, let  $u = (\sin x)^x$  and  $v = \sin^{-1} \sqrt{x}$

$\therefore y = u + v$

$$\Rightarrow \frac{dy}{dx} = \frac{du}{dx} + \frac{dv}{dx} \quad \dots(1)$$

Then,  $u = (\sin x)^x$

Taking logarithm on both the sides, we obtain

$$\Rightarrow \log u = \log (\sin x)^x$$

$$\Rightarrow \log u = x \log (\sin x)$$

Differentiating both sides with respect to  $x$ , we obtain

$$\frac{1}{u} \cdot \frac{du}{dx} = \frac{d}{dx}(x) \times \log (\sin x) + x \cdot \frac{d}{dx}[\log (\sin x)]$$

$$\Rightarrow \frac{du}{dx} = u \left[ 1 \times \log (\sin x) + x \cdot \frac{1}{(\sin x)} \cdot \frac{d}{dx}(\sin x) \right]$$

$$\Rightarrow \frac{du}{dx} = (\sin x)^x \left[ \log (\sin x) + \frac{x}{(\sin x)} \cdot \cos x \right]$$

$$\Rightarrow \frac{du}{dx} = (\sin x)^x [x \cot x + \log \sin x] \quad \dots(2)$$

$$v = \sin^{-1} \sqrt{x}$$

Differentiating both sides with respect to  $x$ , we obtain

$$\frac{dv}{dx} = \frac{1}{\sqrt{1-(\sqrt{x})^2}} \cdot \frac{d}{dx}(\sqrt{x})$$

$$\Rightarrow \frac{dv}{dx} = \frac{1}{\sqrt{1-x}} \cdot \frac{1}{2\sqrt{x}}$$

$$\Rightarrow \frac{dv}{dx} = \frac{1}{2\sqrt{x-x^2}} \quad \dots(3)$$

Therefore, from (1), (2) and (3);

$$\frac{dy}{dx} = (\sin x)^x [x \cot x + \log \sin x] + \frac{1}{2\sqrt{x-x^2}}$$



### Question 9:

Differentiate the function with respect to  $x$ :  $x^{\sin x} + (\sin x)^{\cos x}$

### Solution:

Let  $y = x^{\sin x} + (\sin x)^{\cos x}$

Also, let  $u = x^{\sin x}$  and  $v = (\sin x)^{\cos x}$

$\therefore y = u + v$

$\Rightarrow \frac{dy}{dx} = \frac{du}{dx} + \frac{dv}{dx} \quad \dots(1)$

Then,  $u = x^{\sin x}$

Taking logarithm on both the sides, we obtain

$\Rightarrow \log u = \log(x^{\sin x})$

$\Rightarrow \log u = \sin x \log x$

Differentiating both sides with respect to  $x$ , we obtain

$\frac{1}{u} \cdot \frac{du}{dx} = \frac{d}{dx}(\sin x) \cdot \log x + \sin x \cdot \frac{d}{dx}(\log x)$

$\Rightarrow \frac{du}{dx} = u \left[ \cos x \log x + \sin x \cdot \frac{1}{x} \right]$

$\Rightarrow \frac{du}{dx} = x^{\sin x} \left[ \cos x \log x + \frac{\sin x}{x} \right] \quad \dots(2)$

$v = (\sin x)^{\cos x}$

Taking logarithm on both the sides, we obtain

$$\Rightarrow \log v = \log (\sin x)^{\cos x}$$

$$\Rightarrow \log v = \cos x \log (\sin x)$$

Differentiating both sides with respect to  $x$ , we obtain

$$\frac{1}{v} \frac{dv}{dx} = \frac{d}{dx} (\cos x) \times \log (\sin x) + \cos x \times \frac{d}{dx} [\log (\sin x)]$$

$$\Rightarrow \frac{dv}{dx} = v \left[ -\sin x \cdot \log (\sin x) + \cos x \cdot \frac{1}{\sin x} \cdot \frac{d}{dx} (\sin x) \right]$$

$$\Rightarrow \frac{dv}{dx} = (\sin x)^{\cos x} \left[ -\sin x \log (\sin x) + \frac{\cos x}{\sin x} \cos x \right]$$

$$\Rightarrow \frac{dv}{dx} = (\sin x)^{\cos x} \left[ -\sin x \log (\sin x) + \cot x \cos x \right]$$

$$\Rightarrow \frac{dv}{dx} = (\sin x)^{\cos x} [\cot x \cos x - \sin x \log (\sin x)] \quad \dots(3)$$

Therefore, from (1), (2) and (3);

$$\frac{dy}{dx} = x^{\sin x} \left[ \cos x \log x + \frac{\sin x}{x} \right] + (\sin x)^{\cos x} [\cot x \cos x - \sin x \log (\sin x)]$$

### Question 10:

Differentiate the function with respect to  $x$ :  $x^{x \cos x} + \frac{x^2 + 1}{x^2 - 1}$

**Solution:**

$$\text{Let } y = x^{x \cos x} + \frac{x^2 + 1}{x^2 - 1}$$

$$\text{Also, let } u = x^{x \cos x} \text{ and } v = \frac{x^2 + 1}{x^2 - 1}$$

$$\therefore y = u + v$$

$$\Rightarrow \frac{dy}{dx} = \frac{du}{dx} + \frac{dv}{dx} \quad \dots(1)$$

$$\text{Then, } u = x^{x \cos x}$$

Taking logarithm on both the sides, we obtain

$$\Rightarrow \log u = \log (x^{x \cos x})$$

$$\Rightarrow \log u = x \cos x \log x$$

Differentiating both sides with respect to  $x$ , we obtain

$$\begin{aligned}\frac{1}{u} \cdot \frac{du}{dx} &= \frac{d}{dx}(x) \cdot \cos x \cdot \log x + x \cdot \frac{d}{dx}(\cos x) \cdot \log x + x \cos x \cdot \frac{d}{dx}(\log x) \\ \Rightarrow \frac{du}{dx} &= u \left[ 1 \cdot \cos x \cdot \log x + x \cdot (-\sin x) \log x + x \cos x \cdot \frac{1}{x} \right] \\ \Rightarrow \frac{du}{dx} &= x^{\cos x} [\cos x \log x - x \sin x \log x + \cos x] \\ \Rightarrow \frac{du}{dx} &= x^{\cos x} [\cos x (1 + \log x) - x \sin x \log x] \quad \dots (2)\end{aligned}$$

$$v = \frac{x^2 + 1}{x^2 - 1}$$

Taking logarithm on both the sides, we obtain

$$\Rightarrow \log v = \log(x^2 + 1) - \log(x^2 - 1)$$

Differentiating both sides with respect to  $x$ , we obtain

$$\begin{aligned}\frac{1}{v} \frac{dv}{dx} &= \frac{2x}{x^2 + 1} - \frac{2x}{x^2 - 1} \\ \Rightarrow \frac{dv}{dx} &= v \left[ \frac{2x(x^2 - 1) - 2x(x^2 + 1)}{(x^2 + 1)(x^2 - 1)} \right] \\ \Rightarrow \frac{dv}{dx} &= \frac{x^2 + 1}{x^2 - 1} \times \left[ \frac{-4x}{(x^2 + 1)(x^2 - 1)} \right] \\ \Rightarrow \frac{dv}{dx} &= \frac{-4x}{(x^2 - 1)^2} \quad \dots (3)\end{aligned}$$

Therefore, from (1), (2) and (3);

$$\frac{dy}{dx} = x^{\cos x} [\cos x (1 + \log x) - x \sin x \log x] - \frac{4x}{(x^2 - 1)^2}$$

### Question 11:

Differentiate the function with respect to  $x$ :  $(x \cos x)^x + (x \sin x)^{\frac{1}{x}}$

**Solution:**

$$\text{Let } y = (x \cos x)^x + (x \sin x)^{\frac{1}{x}}$$

$$\text{Also, let } u = (x \cos x)^x \text{ and } v = (x \sin x)^{\frac{1}{x}}$$

$$\therefore y = u + v$$

$$\Rightarrow \frac{dy}{dx} = \frac{du}{dx} + \frac{dv}{dx} \quad \dots(1)$$

$$\text{Then, } u = (x \cos x)^x$$

Taking logarithm on both the sides, we obtain

$$\Rightarrow \log u = (x \cos x)^x$$

$$\Rightarrow \log u = x \log (x \cos x)$$

$$\Rightarrow \log u = x [\log x + \log \cos x]$$

$$\Rightarrow \log u = x \log x + x \log \cos x$$

Differentiating both sides with respect to  $x$ , we obtain

$$\frac{1}{u} \cdot \frac{du}{dx} = \frac{d}{dx}(x \log x) + \frac{d}{dx}(x \log \cos x)$$

$$\Rightarrow \frac{du}{dx} = u \left[ \left\{ \log x \cdot \frac{d}{dx}(x) + x \cdot \frac{d}{dx}(\log x) \right\} + \left\{ \log \cos x \cdot \frac{d}{dx}(x) + x \cdot \frac{d}{dx}(\log \cos x) \right\} \right]$$

$$\Rightarrow \frac{du}{dx} = (x \cos x)^x \left[ \left( \log x \cdot 1 + x \cdot \frac{1}{x} \right) + \left( \log \cos x \cdot 1 + x \cdot \frac{1}{\cos x} \cdot \frac{d}{dx}(\cos x) \right) \right]$$

$$\Rightarrow \frac{du}{dx} = (x \cos x)^x \left[ (\log x + 1) + \left\{ \log \cos x + \frac{x}{\cos x} \cdot (-\sin x) \right\} \right]$$

$$\Rightarrow \frac{du}{dx} = (x \cos x)^x \left[ (1 + \log x) + (\log \cos x - x \tan x) \right]$$

$$\Rightarrow \frac{du}{dx} = (x \cos x)^x \left[ (1 - x \tan x) + (\log x + \log \cos x) \right]$$

$$\Rightarrow \frac{du}{dx} = (x \cos x)^x \left[ 1 - x \tan x + \log (x \cos x) \right] \quad \dots(2)$$

$$v = (x \sin x)^{\frac{1}{x}}$$

Taking logarithm on both the sides, we obtain

$$\Rightarrow \log v = \log (x \sin x)^{\frac{1}{x}}$$

$$\Rightarrow \log v = \frac{1}{x} \log (x \sin x)$$

$$\Rightarrow \log v = \frac{1}{x} (\log x + \log \sin x)$$

$$\Rightarrow \log v = \frac{1}{x} \log x + \frac{1}{x} \log \sin x$$

Differentiating both sides with respect to  $x$ , we obtain

$$\begin{aligned} \frac{1}{v} \frac{dv}{dx} &= \frac{d}{dx} \left( \frac{1}{x} \log x \right) + \frac{d}{dx} \left[ \frac{1}{x} \log (\sin x) \right] \\ \Rightarrow \frac{1}{v} \frac{dv}{dx} &= \left[ \log x \cdot \frac{d}{dx} \left( \frac{1}{x} \right) + \frac{1}{x} \cdot \frac{d}{dx} (\log x) \right] + \left[ \log (\sin x) \cdot \frac{d}{dx} \left( \frac{1}{x} \right) + \frac{1}{x} \cdot \frac{d}{dx} \{ \log (\sin x) \} \right] \\ \Rightarrow \frac{1}{v} \frac{dv}{dx} &= \left[ \log x \cdot \left( -\frac{1}{x^2} \right) + \frac{1}{x} \cdot \frac{1}{x} \right] + \left[ \log (\sin x) \cdot \left( -\frac{1}{x^2} \right) + \frac{1}{x} \cdot \frac{1}{\sin x} \cdot \frac{d}{dx} (\sin x) \right] \\ \Rightarrow \frac{1}{v} \frac{dv}{dx} &= \frac{1}{x^2} (1 - \log x) + \left[ -\frac{\log (\sin x)}{x^2} + \frac{1}{x \sin x} \cdot \cos x \right] \\ \Rightarrow \frac{dv}{dx} &= (x \sin x)^{\frac{1}{x}} \left[ \frac{1 - \log x}{x^2} + \frac{-\log (\sin x) + x \cot x}{x^2} \right] \\ \Rightarrow \frac{dv}{dx} &= (x \sin x)^{\frac{1}{x}} \left[ \frac{1 - \log x - \log (\sin x) + x \cot x}{x^2} \right] \\ \Rightarrow \frac{dv}{dx} &= (x \sin x)^{\frac{1}{x}} \left[ \frac{1 - \log (x \sin x) + x \cot x}{x^2} \right] \quad \dots (3) \end{aligned}$$

Therefore, from (1), (2) and (3);

$$\frac{dy}{dx} = (x \cos x)^x [1 - x \tan x + \log (x \cos x)] + (x \sin x)^{\frac{1}{x}} \left[ \frac{x \cot x + 1 - \log (x \sin x)}{x^2} \right]$$

### Question 12:

Find  $\frac{dy}{dx}$  of the function  $x^y + y^x = 1$

### Solution:

The given function is  $x^y + y^x = 1$

Let,  $x^y = u$  and  $y^x = v$

$$\therefore u + v = 1$$

$$\Rightarrow \frac{du}{dx} + \frac{dv}{dx} = 0 \quad \dots(1)$$

Then,  $u = x^y$

Taking logarithm on both the sides, we obtain

$$\Rightarrow \log u = \log(x^y)$$

$$\Rightarrow \log u = y \log x$$

Differentiating both sides with respect to  $x$ , we obtain

$$\frac{1}{u} \cdot \frac{du}{dx} = \log x \cdot \frac{dy}{dx} + y \cdot \frac{d}{dx}(\log x)$$

$$\Rightarrow \frac{du}{dx} = u \left[ \log x \cdot \frac{dy}{dx} + y \cdot \frac{1}{x} \right]$$

$$\Rightarrow \frac{du}{dx} = x^y \left[ \log x \cdot \frac{dy}{dx} + \frac{y}{x} \right] \quad \dots(2)$$

Now,  $v = y^x$

Taking logarithm on both the sides, we obtain

$$\Rightarrow \log v = \log(y^x)$$

$$\Rightarrow \log v = x \log y$$

Differentiating both sides with respect to  $x$ , we obtain

$$\frac{1}{v} \frac{dv}{dx} = \log y \cdot \frac{d}{dx}(x) + x \cdot \frac{d}{dx}(\log y)$$

$$\Rightarrow \frac{dv}{dx} = v \left[ \log y \cdot 1 + x \cdot \frac{1}{y} \cdot \frac{dy}{dx} \right]$$

$$\Rightarrow \frac{dv}{dx} = y^x \left[ \log y + \frac{x}{y} \cdot \frac{dy}{dx} \right] \quad \dots(3)$$

Therefore, from (1), (2) and (3);

$$\begin{aligned}
 x^y \left[ \log x \frac{dy}{dx} + \frac{y}{x} \right] + y^x \left[ \log y + \frac{x}{y} \cdot \frac{dy}{dx} \right] &= 0 \\
 \Rightarrow (x^y \log x + xy^{x-1}) \frac{dy}{dx} &= -(yx^{y-1} + y^x \log y) \\
 \therefore \frac{dy}{dx} &= \frac{-(yx^{y-1} + y^x \log y)}{(x^y \log x + xy^{x-1})}
 \end{aligned}$$

### Question 13:

Find  $\frac{dy}{dx}$  of the function  $y^x = x^y$

#### Solution:

The given function is  $y^x = x^y$

Taking logarithm on both the sides, we obtain

$$x \log y = y \log x$$

Differentiating both sides with respect to  $x$ , we obtain

$$\begin{aligned}
 \log y \cdot \frac{d}{dx}(x) + x \cdot \frac{d}{dx}(\log y) &= \log x \cdot \frac{d}{dx}(y) + y \cdot \frac{d}{dx}(\log x) \\
 \Rightarrow \log y \cdot 1 + x \cdot \frac{1}{y} \cdot \frac{dy}{dx} &= \log x \cdot \frac{dy}{dx} + y \cdot \frac{1}{x} \\
 \Rightarrow \log y + \frac{x}{y} \cdot \frac{dy}{dx} &= \log x \cdot \frac{dy}{dx} + \frac{y}{x} \\
 \Rightarrow \left( \frac{x}{y} - \log x \right) \frac{dy}{dx} &= \frac{y}{x} - \log y \\
 \Rightarrow \left( \frac{x - y \log x}{y} \right) \frac{dy}{dx} &= \frac{y - x \log y}{x} \\
 \therefore \frac{dy}{dx} &= \frac{y}{x} \left( \frac{y - x \log y}{x - y \log x} \right)
 \end{aligned}$$

### Question 14:

Find  $\frac{dy}{dx}$  of the function  $(\cos x)^y = (\cos y)^x$

### Solution:

The given function is  $(\cos x)^y = (\cos y)^x$

Taking logarithm on both the sides, we obtain

$$y \log \cos x = x \log \cos y$$

Differentiating both sides with respect to  $x$ , we obtain

$$\begin{aligned} \log \cos x \cdot \frac{dy}{dx} + y \cdot \frac{d}{dx}(\log \cos x) &= \log \cos y \cdot \frac{d}{dx}(x) + x \cdot \frac{d}{dx}(\log \cos y) \\ \Rightarrow \log \cos x \cdot \frac{dy}{dx} + y \cdot \frac{1}{\cos x} \cdot \frac{d}{dx}(\cos x) &= \log \cos y \cdot 1 + x \cdot \frac{1}{\cos y} \cdot \frac{d}{dx}(\cos y) \\ \Rightarrow \log \cos x \cdot \frac{dy}{dx} + \frac{y}{\cos x} \cdot (-\sin x) &= \log \cos y + \frac{x}{\cos y} \cdot (-\sin y) \cdot \frac{dy}{dx} \\ \Rightarrow \log \cos x \cdot \frac{dy}{dx} - y \tan x &= \log \cos y - x \tan y \frac{dy}{dx} \\ \Rightarrow (\log \cos x + x \tan y) \frac{dy}{dx} &= y \tan x + \log \cos y \\ \therefore \frac{dy}{dx} &= \frac{y \tan x + \log \cos y}{x \tan y + \log \cos x} \end{aligned}$$

### Question 15:

Find  $\frac{dy}{dx}$  of the function  $xy = e^{(x-y)}$

### Solution:

The given function is  $xy = e^{(x-y)}$

Taking logarithm on both the sides, we obtain

$$\begin{aligned} \log(xy) &= \log(e^{x-y}) \\ \Rightarrow \log x + \log y &= (x-y) \log e \\ \Rightarrow \log x + \log y &= (x-y) \times 1 \\ \Rightarrow \log x + \log y &= (x-y) \end{aligned}$$

Differentiating both sides with respect to  $x$ , we obtain



$$\begin{aligned}\frac{d}{dx}(\log x) + \frac{d}{dx}(\log y) &= \frac{d}{dx}(x) - \frac{dy}{dx} \\ \Rightarrow \frac{1}{x} + \frac{1}{y} \frac{dy}{dx} &= 1 - \frac{dy}{dx} \\ \Rightarrow \left(1 + \frac{1}{y}\right) \frac{dy}{dx} &= 1 - \frac{1}{x} \\ \Rightarrow \left(\frac{y+1}{y}\right) \frac{dy}{dx} &= \frac{x-1}{x} \\ \therefore \frac{dy}{dx} &= \frac{y(x-1)}{x(y+1)}\end{aligned}$$

### Question 16:

Find the derivative of the function given by  $f(x) = (1+x)(1+x^2)(1+x^4)(1+x^8)$  and hence find  $f'(1)$ .

### Solution:

The given function is  $f(x) = (1+x)(1+x^2)(1+x^4)(1+x^8)$

Taking logarithm on both the sides, we obtain

$$\log f(x) = \log(1+x) + \log(1+x^2) + \log(1+x^4) + \log(1+x^8)$$

Differentiating both sides with respect to  $x$ , we obtain

$$\begin{aligned}\frac{1}{f(x)} \cdot \frac{d}{dx}[f(x)] &= \frac{d}{dx} \log(1+x) + \frac{d}{dx} \log(1+x^2) \\ &\quad + \frac{d}{dx} \log(1+x^4) + \frac{d}{dx} \log(1+x^8) \\ \Rightarrow \frac{1}{f(x)} \cdot f'(x) &= \frac{1}{1+x} \cdot \frac{d}{dx}(1+x) + \frac{1}{1+x^2} \cdot \frac{d}{dx}(1+x^2) \\ &\quad + \frac{1}{1+x^4} \cdot \frac{d}{dx}(1+x^4) + \frac{1}{1+x^8} \cdot \frac{d}{dx}(1+x^8) \\ \Rightarrow f'(x) &= f(x) \left[ \frac{1}{1+x} + \frac{1}{1+x^2} \cdot 2x + \frac{1}{1+x^4} \cdot 4x^3 + \frac{1}{1+x^8} \cdot 8x^7 \right] \\ \therefore f'(x) &= (1+x)(1+x^2)(1+x^4)(1+x^8) \left[ \frac{1}{1+x} + \frac{2x}{1+x^2} + \frac{4x^3}{1+x^4} + \frac{8x^7}{1+x^8} \right]\end{aligned}$$

Hence,

$$\begin{aligned}f'(1) &= (1+1)(1+1^2)(1+1^4)(1+1^8) \left[ \frac{1}{1+1} + \frac{2(1)}{1+1^2} + \frac{4(1)^3}{1+1^4} + \frac{8(1)^7}{1+1^8} \right] \\ &= 2 \times 2 \times 2 \times 2 \left[ \frac{1}{2} + \frac{2}{2} + \frac{4}{2} + \frac{8}{2} \right] \\ &= 16 \left( \frac{15}{2} \right) \\ &= 120\end{aligned}$$

### Question 17:

Differentiate  $(x^2 - 5x + 8)(x^3 + 7x + 9)$  in three ways mentioned below.

- By using product rule
- By expanding the product to obtain a single polynomial.
- By logarithmic differentiation.

Do they all give the same answer?

**Solution:**

Let  $y = (x^2 - 5x + 8)(x^3 + 7x + 9)$

- By using product rule

Let  $u = (x^2 - 5x + 8)$  and  $v = x^3 + 7x + 9$

$$\therefore y = uv$$

$$\Rightarrow \frac{dy}{dx} = \frac{du}{dx} \cdot v + u \cdot \frac{dv}{dx} \quad (\text{product rule})$$

$$\Rightarrow \frac{dy}{dx} = \frac{d}{dx}(x^2 - 5x + 8) \cdot (x^3 + 7x + 9) + (x^2 - 5x + 8) \cdot \frac{d}{dx}(x^3 + 7x + 9)$$

$$\Rightarrow \frac{dy}{dx} = (2x - 5)(x^3 + 7x + 9) + (x^2 - 5x + 8)(3x^2 + 7)$$

$$\Rightarrow \frac{dy}{dx} = 2x(x^3 + 7x + 9) - 5(x^3 + 7x + 9) + x^2(3x^2 + 7)$$

$$- 5x(3x^2 + 7) + 8(3x^2 + 7)$$

$$\Rightarrow \frac{dy}{dx} = (2x^4 + 14x^2 + 18x) - 5x^3 - 35x - 45 + (3x^4 + 7x^2)$$

$$\Rightarrow \frac{dy}{dx} = -15x^3 - 35x + 24x^2 + 56$$

$$\therefore \frac{dy}{dx} = 5x^4 - 20x^3 + 45x^2 - 52x + 11$$

(ii) By expanding the product to obtain a single polynomial.

$$y = (x^2 - 5x + 8)(x^3 + 7x + 9)$$

$$= x^2(x^3 + 7x + 9) - 5x(x^3 + 7x + 9) + 8(x^3 + 7x + 9)$$

$$= x^5 + 7x^3 + 9x^2 - 5x^4 - 35x^2 - 45x + 8x^3 + 56x + 72$$

$$= x^5 - 5x^4 + 15x^3 - 26x^2 + 11x + 72$$

Therefore,

$$\frac{dy}{dx} = \frac{d}{dx}(x^5 - 5x^4 + 15x^3 - 26x^2 + 11x + 72)$$

$$= \frac{d}{dx}(x^5) - 5 \frac{d}{dx}(x^4) + 15 \frac{d}{dx}(x^3) - 26 \frac{d}{dx}(x^2) + 11 \frac{d}{dx}(x) + \frac{d}{dx}(72)$$

$$= 5x^4 - 5(4x^3) + 15(3x^2) - 26(2x) + 11(1) + 0$$

$$= 5x^4 - 20x^3 + 45x^2 - 52x + 11$$

(iii) By logarithmic differentiation.

$$y = (x^2 - 5x + 8)(x^3 + 7x + 9)$$

Taking logarithm on both the sides, we obtain

$$\log y = \log(x^2 - 5x + 8) + \log(x^3 + 7x + 9)$$

Differentiating both sides with respect to  $x$ , we obtain

$$\begin{aligned}\frac{1}{y} \cdot \frac{dy}{dx} &= \frac{d}{dx} \log(x^2 - 5x + 8) + \frac{d}{dx} \log(x^3 + 7x + 9) \\ \Rightarrow \frac{1}{y} \cdot \frac{dy}{dx} &= \frac{1}{x^2 - 5x + 8} \cdot \frac{d}{dx}(x^2 - 5x + 8) + \frac{1}{x^3 + 7x + 9} \cdot \frac{d}{dx}(x^3 + 7x + 9) \\ \Rightarrow \frac{dy}{dx} &= y \left[ \frac{1}{x^2 - 5x + 8} \cdot (2x - 5) + \frac{1}{x^3 + 7x + 9} \cdot (3x^2 + 7) \right] \\ \Rightarrow \frac{dy}{dx} &= (x^2 - 5x + 8)(x^3 + 7x + 9) \left[ \frac{2x - 5}{x^2 - 5x + 8} + \frac{3x^2 + 7}{x^3 + 7x + 9} \right] \\ \Rightarrow \frac{dy}{dx} &= (x^2 - 5x + 8)(x^3 + 7x + 9) \left[ \frac{(2x - 5)(x^3 + 7x + 9) + (3x^2 + 7)(x^2 - 5x + 8)}{(x^2 - 5x + 8)(x^3 + 7x + 9)} \right] \\ \Rightarrow \frac{dy}{dx} &= 2x(x^3 + 7x + 9) - 5(x^3 + 7x + 9) + 3x^2(x^2 - 5x + 8) + 7(x^2 - 5x + 8) \\ \Rightarrow \frac{dy}{dx} &= 2x^4 + 14x^2 + 18x - 5x^3 - 35x - 45 + 3x^5 - 15x^3 + 24x^2 + 7x^2 - 35x + 56 \\ \Rightarrow \frac{dy}{dx} &= 5x^4 - 20x^3 + 45x^2 - 52x + 11\end{aligned}$$

From the above three observations, it can be concluded that all the results of  $\frac{dy}{dx}$  are same.

### Question 18:

If  $u$ ,  $v$  and  $w$  are functions of  $x$ , then show that

$$\frac{d}{dx}(u.v.w) = \frac{du}{dx}.v.w + u.\frac{dv}{dx}.w + u.v.\frac{dw}{dx}$$

in two ways - first by repeated application of product rule, second by logarithmic differentiation.

### Solution:

Let  $y = u.v.w = u.(v.w)$

By applying product rule, we get

$$\begin{aligned}\frac{dy}{dx} &= \frac{du}{dx} \cdot (v.w) + u \cdot \frac{d}{dx}(v.w) \\ \Rightarrow \frac{dy}{dx} &= \frac{du}{dx} \cdot (v.w) + u \left[ \frac{dv}{dx} \cdot w + v \cdot \frac{dw}{dx} \right] \quad (\text{Again applying product rule}) \\ \Rightarrow \frac{dy}{dx} &= \frac{du}{dx} \cdot v.w + u \cdot \frac{dv}{dx} \cdot w + u.v \cdot \frac{dw}{dx}\end{aligned}$$



Taking logarithm on both the sides of the equation  $y = u.v.w$ , we obtain

$$\log y = \log u + \log v + \log w$$

Differentiating both sides with respect to  $x$ , we obtain


$$\frac{1}{y} \cdot \frac{dy}{dx} = \frac{d}{dx}(\log u) + \frac{d}{dx}(\log v) + \frac{d}{dx}(\log w)$$

$$\Rightarrow \frac{1}{y} \cdot \frac{dy}{dx} = \frac{1}{u} \frac{du}{dx} + \frac{1}{v} \frac{dv}{dx} + \frac{1}{w} \frac{dw}{dx}$$

$$\Rightarrow \frac{dy}{dx} = y \left( \frac{1}{u} \frac{du}{dx} + \frac{1}{v} \frac{dv}{dx} + \frac{1}{w} \frac{dw}{dx} \right)$$

$$\Rightarrow \frac{dy}{dx} = u.v.w \left( \frac{1}{u} \frac{du}{dx} + \frac{1}{v} \frac{dv}{dx} + \frac{1}{w} \frac{dw}{dx} \right)$$

$$\therefore \frac{d}{dx}(u.v.w) = \frac{du}{dx} v.w + u \cdot \frac{dv}{dx} .w + u.v \cdot \frac{dw}{dx}$$



## EXERCISE 5.6

### Question 1:

If  $x$  and  $y$  are connected parametrically by the equations  $x = 2at^2$ ,  $y = at^4$ , without eliminating the parameter, find  $\frac{dy}{dx}$

### Solution:

Given,  $x = 2at^2$ ,  $y = at^4$

Then,

$$\frac{dx}{dt} = \frac{d}{dt}(2at^2) = 2a \cdot \frac{d}{dt}(t^2) = 2a \cdot 2t = 4at$$

$$\frac{dy}{dt} = \frac{d}{dt}(at^4) = a \cdot \frac{d}{dt}(t^4) = a \cdot 4t^3 = 4at^3$$

$$\therefore \frac{dy}{dx} = \frac{\left(\frac{dy}{dt}\right)}{\left(\frac{dx}{dt}\right)} = \frac{4at^3}{4at} = t^2$$

### Question 2:

If  $x$  and  $y$  are connected parametrically by the equations  $x = a \cos \theta$ ,  $y = b \cos \theta$ , without eliminating the parameter, find  $\frac{dy}{dx}$

### Solution:

Given,  $x = a \cos \theta$ ,  $y = b \cos \theta$

Then,

$$\frac{dx}{d\theta} = \frac{d}{d\theta}(a \cos \theta) = a(-\sin \theta) = -a \sin \theta$$

$$\frac{dy}{d\theta} = \frac{d}{d\theta}(b \cos \theta) = b(-\sin \theta) = -b \sin \theta$$

$$\therefore \frac{dy}{dx} = \frac{\left(\frac{dy}{d\theta}\right)}{\left(\frac{dx}{d\theta}\right)} = \frac{-b \sin \theta}{-a \sin \theta} = \frac{b}{a}$$

### Question 3:

If  $x$  and  $y$  are connected parametrically by the equations  $x = \sin t, y = \cos 2t$ , without eliminating the parameter, find  $\frac{dy}{dx}$

### Solution:

Given,  $x = \sin t, y = \cos 2t$

$$\text{Then, } \frac{dx}{dt} = \frac{d}{dt}(\sin t) = \cos t$$

$$\frac{dy}{dt} = \frac{d}{dt}(\cos 2t) = -\sin 2t \cdot \frac{d}{dt}(2t) = -2 \sin 2t$$

$$\therefore \frac{dy}{dx} = \frac{\left(\frac{dy}{dt}\right)}{\left(\frac{dx}{dt}\right)} = \frac{-2 \sin 2t}{\cos t} = \frac{-2 \cdot 2 \sin t \cos t}{\cos t} = -4 \sin t$$

### Question 4:

If  $x$  and  $y$  are connected parametrically by the equations  $x = 4t, y = \frac{4}{t}$ , without eliminating the parameter, find  $\frac{dy}{dx}$

### Solution:

Given,  $x = 4t, y = \frac{4}{t}$

$$\frac{dx}{dt} = \frac{d}{dt}(4t) = 4$$

$$\frac{dy}{dt} = \frac{d}{dt}\left(\frac{4}{t}\right) = 4 \cdot \frac{d}{dt}\left(\frac{1}{t}\right) = 4 \cdot \left(\frac{-1}{t^2}\right) = \frac{-4}{t^2}$$

$$\therefore \frac{dy}{dx} = \frac{\left(\frac{dy}{dt}\right)}{\left(\frac{dx}{dt}\right)} = \frac{\left(\frac{-4}{t^2}\right)}{4} = \frac{-1}{t^2}$$

### Question 5:

If  $x$  and  $y$  are connected parametrically by the equations  $x = \cos \theta - \cos 2\theta, y = \sin \theta - \sin 2\theta$ , without eliminating the parameter, find  $\frac{dy}{dx}$

### Solution:

Given,  $x = \cos \theta - \cos 2\theta$ ,  $y = \sin \theta - \sin 2\theta$

Then,

$$\frac{dx}{d\theta} = \frac{d}{d\theta}(\cos \theta - \cos 2\theta) = \frac{d}{d\theta}(\cos \theta) - \frac{d}{d\theta}(\cos 2\theta)$$

$$= -\sin \theta - (-2 \sin 2\theta) = 2 \sin 2\theta - \sin \theta$$

$$\frac{dy}{d\theta} = \frac{d}{d\theta}(\sin \theta - \sin 2\theta) = \frac{d}{d\theta}(\sin \theta) - \frac{d}{d\theta}(\sin 2\theta)$$

$$= \cos \theta - 2 \cos 2\theta$$

$$\therefore \frac{dy}{dx} = \frac{\left(\frac{dy}{d\theta}\right)}{\left(\frac{dx}{d\theta}\right)} = \frac{\cos \theta - 2 \cos 2\theta}{2 \sin 2\theta - \sin \theta}$$

### Question 6:

If  $x$  and  $y$  are connected parametrically by the equations  $x = a(\theta - \sin \theta)$ ,  $y = a(1 + \cos \theta)$ ,

without eliminating the parameter, find  $\frac{dy}{dx}$

### Solution:

Given,  $x = a(\theta - \sin \theta)$ ,  $y = a(1 + \cos \theta)$

Then,  $\frac{dx}{d\theta} = a \left[ \frac{d}{d\theta}(\theta) - \frac{d}{d\theta}(\sin \theta) \right] = a(1 - \cos \theta)$

$$\frac{dy}{d\theta} = a \left[ \frac{d}{d\theta}(1) + \frac{d}{d\theta}(\cos \theta) \right] = a[0 + (-\sin \theta)] = -a \sin \theta$$

$$\therefore \frac{dy}{dx} = \frac{\left(\frac{dy}{d\theta}\right)}{\left(\frac{dx}{d\theta}\right)} = \frac{-a \sin \theta}{a(1 - \cos \theta)} = \frac{-2 \sin \frac{\theta}{2} \cos \frac{\theta}{2}}{2 \sin^2 \frac{\theta}{2}} = \frac{-\cos \frac{\theta}{2}}{\sin \frac{\theta}{2}} = -\cot \frac{\theta}{2}$$

### Question 7:

If  $x$  and  $y$  are connected parametrically by the equations  $x = \frac{\sin^3 t}{\sqrt{\cos 2t}}$ ,  $y = \frac{\cos^3 t}{\sqrt{\cos 2t}}$ , without

eliminating the parameter, find  $\frac{dy}{dx}$



**Solution:**

Given,  $x = \frac{\sin^3 t}{\sqrt{\cos 2t}}, y = \frac{\cos^3 t}{\sqrt{\cos 2t}}$

Then,

$$\begin{aligned} \frac{dx}{dt} &= \frac{d}{dt} \left[ \frac{\sin^3 t}{\sqrt{\cos 2t}} \right] \\ &= \frac{\sqrt{\cos 2t} \cdot \frac{d}{dt} (\sin^3 t) - \sin^3 t \cdot \frac{d}{dt} \sqrt{\cos 2t}}{\cos 2t} \\ &= \frac{\sqrt{\cos 2t} \cdot 3 \sin^2 t \cdot \frac{d}{dt} (\sin t) - \sin^3 t \times \frac{1}{2\sqrt{\cos 2t}} \cdot \frac{d}{dt} (\cos 2t)}{\cos 2t} \\ &= \frac{3\sqrt{\cos 2t} \cdot \sin^2 t \cdot \cos t - \frac{\sin^3 t}{2\sqrt{\cos 2t}} \cdot (-2 \sin 2t)}{\cos 2t} \\ &= \frac{3 \cos 2t \cdot \sin^2 t \cos t + \sin^3 t \cdot \sin 2t}{\cos 2t \sqrt{\cos 2t}} \\ \frac{dy}{dt} &= \frac{d}{dt} \left[ \frac{\cos^3 t}{\sqrt{\cos 2t}} \right] \\ &= \frac{\sqrt{\cos 2t} \cdot \frac{d}{dt} (\cos^3 t) - \cos^3 t \cdot \frac{d}{dt} (\sqrt{\cos 2t})}{\cos 2t} \\ &= \frac{\sqrt{\cos 2t} \cdot 3 \cos^2 t \cdot \frac{d}{dt} (\cos t) - \cos^3 t \cdot \frac{1}{2\sqrt{\cos 2t}} \cdot \frac{d}{dt} (\cos 2t)}{\cos 2t} \\ &= \frac{3\sqrt{\cos 2t} \cdot \cos^2 t \cdot (-\sin t) - \cos^3 t \cdot \frac{1}{\sqrt{\cos 2t}} \cdot (-2 \sin 2t)}{\cos 2t} \\ &= \frac{-3 \cos 2t \cdot \cos^2 t \cdot \sin t + \cos^3 t \cdot \sin 2t}{\cos 2t \sqrt{\cos 2t}} \end{aligned}$$

$$\begin{aligned}
 \therefore \frac{dy}{dx} &= \frac{\left(\frac{dy}{dt}\right)}{\left(\frac{dx}{dt}\right)} = \frac{\frac{-3 \cos 2t \cdot \cos^2 t \cdot \sin t + \cos^3 t \sin 2t}{\cos 2t \cdot \sqrt{\cos 2t}}}{\frac{3 \cos 2t \cdot \sin^2 t \cdot \cos t + \sin^3 t \sin 2t}{\cos 2t \cdot \sqrt{\cos 2t}}} \\
 &= \frac{-3 \cos 2t \cdot \cos^2 t \cdot \sin t + \cos^3 t \sin 2t}{3 \cos 2t \cdot \sin^2 t \cdot \cos t + \sin^3 t \sin 2t} \\
 &= \frac{-3 \cos 2t \cdot \cos^2 t \cdot \sin t + \cos^3 t (2 \sin t \cos t)}{3 \cos 2t \cdot \sin^2 t \cdot \cos t + \sin^3 t (2 \sin t \cos t)} \\
 &= \frac{\sin t \cos t [-3 \cos 2t \cdot \cos t + 2 \cos^3 t]}{\sin t \cos t [3 \cos 2t \sin t + 2 \sin^3 t]} \\
 &= \frac{[-3(2 \cos^2 t - 1) \cos t + 2 \cos^3 t]}{[3(1 - 2 \sin^2 t) \sin t + 2 \sin^3 t]} \quad \begin{cases} \cos 2t = (2 \cos^2 t - 1) \\ \cos 2t = (1 - 2 \sin^2 t) \end{cases} \\
 &= \frac{-4 \cos^3 t + 3 \cos t}{3 \sin t - 4 \sin^3 t} \quad \begin{cases} \cos 3t = 4 \cos^3 t - 3 \cos t \\ \sin 3t = 3 \sin t - 4 \sin^3 t \end{cases} \\
 &= \frac{-\cos 3t}{\sin 3t} = -\cot 3t
 \end{aligned}$$

### Question 8:

If  $x$  and  $y$  are connected parametrically by the equations  $x = a \left( \cos t + \log \tan \frac{t}{2} \right)$ ,  $y = a \sin t$ ,  
without eliminating the parameter, find  $\frac{dy}{dx}$

### Solution:

Given,  $x = a \left( \cos t + \log \tan \frac{t}{2} \right)$ ,  $y = a \sin t$

Then,

$$\begin{aligned}\frac{dx}{dt} &= a \left[ \frac{d}{dt}(\cos t) + \frac{d}{dt} \left( \log \tan \frac{t}{2} \right) \right] \\ &= a \left[ -\sin t + \frac{1}{\tan \frac{t}{2}} \cdot \frac{d}{dt} \left( \tan \frac{t}{2} \right) \right] \\ &= a \left[ -\sin t + \cot \frac{t}{2} \cdot \sec^2 \frac{t}{2} \cdot \frac{d}{dt} \left( \frac{t}{2} \right) \right] \\ &= a \left[ -\sin t + \frac{\cos \frac{t}{2}}{\sin \frac{t}{2}} \times \frac{1}{\cos^2 \frac{t}{2}} \times \frac{1}{2} \right] \\ &= a \left[ -\sin t + \frac{1}{2 \sin \frac{t}{2} \cos \frac{t}{2}} \right] \\ &= a \left( -\sin t + \frac{1}{\sin t} \right) \\ &= a \left( \frac{-\sin^2 t + 1}{\sin t} \right) \\ &= a \left( \frac{\cos^2 t}{\sin t} \right)\end{aligned}$$

$$\frac{dy}{dt} = a \frac{d}{dt}(\sin t) = a \cos t$$

Therefore,

$$\frac{dy}{dx} = \frac{\left( \frac{dy}{dt} \right)}{\left( \frac{dx}{dt} \right)} = \frac{a \cos t}{\left( a \frac{\cos^2 t}{\sin t} \right)} = \frac{\sin t}{\cos t} = \tan t$$

### Question 9:

If  $x$  and  $y$  are connected parametrically by the equations  $x = a \sec \theta$ ,  $y = b \tan \theta$ , without

eliminating the parameter, find  $\frac{dy}{dx}$

### Solution:

Given,  $x = a \sec \theta$ ,  $y = b \tan \theta$

Then,

$$\frac{dx}{d\theta} = a \cdot \frac{d}{d\theta}(\sec \theta) = a \sec \theta \tan \theta$$

$$\frac{dy}{d\theta} = b \cdot \frac{d}{d\theta}(\tan \theta) = b \sec^2 \theta$$

Therefore,

$$\begin{aligned} \frac{dy}{dx} &= \frac{\left(\frac{dy}{d\theta}\right)}{\left(\frac{dx}{d\theta}\right)} \\ &= \frac{b \sec^2 \theta}{a \sec \theta \tan \theta} \\ &= \frac{b}{a} \sec \theta \cot \theta \\ &= \frac{b \cos \theta}{a \cos \theta \sin \theta} \\ &= \frac{b}{a} \times \frac{1}{\sin \theta} \\ &= \frac{b}{a} \operatorname{cosec} \theta \end{aligned}$$

### Question 10:

If  $x$  and  $y$  are connected parametrically by the equations

$x = a(\cos \theta + \theta \sin \theta)$ ,  $y = a(\sin \theta - \theta \cos \theta)$ , without eliminating the parameter, find  $\frac{dy}{dx}$

### Solution:

Given,  $x = a(\cos \theta + \theta \sin \theta)$ ,  $y = a(\sin \theta - \theta \cos \theta)$

Then,

$$\begin{aligned} \frac{dx}{d\theta} &= a \left[ \frac{d}{d\theta} \cos \theta + \frac{d}{d\theta} (\theta \sin \theta) \right] \\ &= a \left[ -\sin \theta + \theta \frac{d}{d\theta} (\sin \theta) + \sin \theta \frac{d}{d\theta} (\theta) \right] \\ &= a [-\sin \theta + \theta \cos \theta + \sin \theta] \\ &= a \theta \cos \theta \end{aligned}$$

$$\begin{aligned} \frac{dy}{d\theta} &= a \left[ \frac{d}{d\theta} (\sin \theta) - \frac{d}{d\theta} (\theta \cos \theta) \right] = a \left[ \cos \theta - \left\{ \theta \frac{d}{d\theta} (\cos \theta) + \cos \theta \cdot \frac{d}{d\theta} (\theta) \right\} \right] \\ &= a [\cos \theta + \theta \sin \theta - \cos \theta] \\ &= a \theta \sin \theta \end{aligned}$$

Therefore,

$$\begin{aligned}\frac{dy}{dx} &= \frac{\left(\frac{dy}{d\theta}\right)}{\left(\frac{dx}{d\theta}\right)} \\ &= \frac{a\theta \sin \theta}{a\theta \cos \theta} \\ &= \tan \theta\end{aligned}$$

### Question 11:

If  $x = \sqrt{a^{\sin^{-1} t}}$ ,  $y = \sqrt{a^{\cos^{-1} t}}$ , show that  $\frac{dy}{dx} = -\frac{y}{x}$

**Solution:**

Given,  $x = \sqrt{a^{\sin^{-1} t}}$  and  $y = \sqrt{a^{\cos^{-1} t}}$

Hence,

$$x = \sqrt{a^{\sin^{-1} t}} = \left(a^{\sin^{-1} t}\right)^{\frac{1}{2}} = a^{\frac{1}{2}\sin^{-1} t} \quad \text{and} \quad y = \sqrt{a^{\cos^{-1} t}} = \left(a^{\cos^{-1} t}\right)^{\frac{1}{2}} = a^{\frac{1}{2}\cos^{-1} t}$$

Consider  $x = a^{\frac{1}{2}\sin^{-1} t}$

Taking log on both sides, we get

$$\log x = \frac{1}{2} \sin^{-1} t \log a$$

Therefore,

$$\begin{aligned}\Rightarrow \frac{1}{x} \cdot \frac{dx}{dt} &= \frac{1}{2} \log a \cdot \frac{d}{dt}(\sin^{-1} t) \\ \Rightarrow \frac{dx}{dt} &= \frac{x}{2} \log a \cdot \frac{1}{\sqrt{1-t^2}} \\ \Rightarrow \frac{dx}{dt} &= \frac{x \log a}{2\sqrt{1-t^2}}\end{aligned}$$

Now,  $y = a^{\frac{1}{2}\cos^{-1} t}$

Taking log on both sides, we get

$$\log y = \frac{1}{2} \cos^{-1} t \log a$$



Therefore,

$$\Rightarrow \frac{1}{y} \cdot \frac{dy}{dt} = \frac{1}{2} \log a \cdot \frac{d}{dt} (\cos^{-1} t)$$

$$\Rightarrow \frac{dy}{dt} = \frac{y}{2} \log a \cdot \frac{-1}{\sqrt{1-t^2}}$$

$$\Rightarrow \frac{dy}{dt} = \frac{-y \log a}{2\sqrt{1-t^2}}$$

Hence,

$$\frac{dy}{dx} = \frac{\left(\frac{dy}{dt}\right)}{\left(\frac{dx}{dt}\right)} = \frac{\left(\frac{-y \log a}{2\sqrt{1-t^2}}\right)}{\left(\frac{x \log a}{2\sqrt{1-t^2}}\right)} = -\frac{y}{x}$$



MillionStars edu  
Think, Learn & Practice



## EXERCISE 5.7

### Question 1:

Find the second order derivative of the function  $x^2 + 3x + 2$

### Solution:

Consider,  $y = x^2 + 3x + 2$

Then,

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx}(x^2) + \frac{d}{dx}(3x) + \frac{d}{dx}(2) \\ &= 2x + 3 + 0 \\ &= 2x + 3\end{aligned}$$

Therefore,

$$\begin{aligned}\frac{d^2y}{dx^2} &= \frac{d}{dx}(2x + 3) \\ &= \frac{d}{dx}(2x) + \frac{d}{dx}(3) \\ &= 2 + 0 \\ &= 2\end{aligned}$$

### Question 2:

Find the second order derivative of the function  $x^{20}$

### Solution:

Consider,  $y = x^{20}$

Then,

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx}(x^{20}) \\ &= 20x^{19}\end{aligned}$$

Therefore,

$$\begin{aligned}\frac{d^2y}{dx^2} &= \frac{d}{dx}(20x^{19}) \\ &= 20 \frac{d}{dx}(x^{19}) \\ &= 20 \cdot 19 \cdot x^{18} \\ &= 380x^{18}\end{aligned}$$

### Question 3:

Find the second order derivative of the function  $x \cos x$

#### Solution:

Consider,  $y = x \cos x$

Then,

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx}(x \cdot \cos x) \\ &= \cos x \cdot \frac{d}{dx}(x) + x \frac{d}{dx}(\cos x) \\ &= \cos x \cdot 1 + x(-\sin x) \\ &= \cos x - x \sin x\end{aligned}$$

Therefore,

$$\begin{aligned}\frac{d^2y}{dx^2} &= \frac{d}{dx}[\cos x - x \sin x] \\ &= \frac{d}{dx}(\cos x) - \frac{d}{dx}(x \sin x) \\ &= -\sin x - \left[ \sin x \cdot \frac{d}{dx}(x) + x \cdot \frac{d}{dx}(\sin x) \right] \\ &= -\sin x - (\sin x + x \cos x) \\ &= -(x \cos x + 2 \sin x)\end{aligned}$$

### Question 4:

Find the second order derivative of the function  $\log x$

#### Solution:

Let  $y = \log x$

Then,

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx}(\log x) \\ &= \frac{1}{x}\end{aligned}$$

Therefore,

$$\begin{aligned}\frac{d^2y}{dx^2} &= \frac{d}{dx}\left(\frac{1}{x}\right) \\ &= \frac{-1}{x^2}\end{aligned}$$



### Question 5:

Find the second order derivative of the function  $x^3 \log x$

#### Solution:

Let  $y = x^3 \log x$

Then,

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx} [x^3 \log x] \\ &= \log x \cdot \frac{d}{dx} (x^3) + x^3 \cdot \frac{d}{dx} (\log x) \\ &= \log x \cdot 3x^2 + x^3 \cdot \frac{1}{x} = \log x \cdot 3x^2 + x^2 \\ &= x^2 (1 + 3 \log x)\end{aligned}$$

Therefore,

$$\begin{aligned}\frac{d^2y}{dx^2} &= \frac{d}{dx} [x^2 (1 + 3 \log x)] \\ &= (1 + 3 \log x) \cdot \frac{d}{dx} (x^2) + x^2 \cdot \frac{d}{dx} (1 + 3 \log x) \\ &= (1 + 3 \log x) \cdot 2x + x^2 \cdot \frac{3}{x} \\ &= 2x + 6 \log x + 3x \\ &= 5x + 6x \log x \\ &= x(5 + 6 \log x)\end{aligned}$$

### Question 6:

Find the second order derivative of the function  $e^x \sin 5x$

#### Solution:

Let  $y = e^x \sin 5x$

Then,

$$\begin{aligned}
 \frac{dy}{dx} &= \frac{d}{dx}(e^x \sin 5x) \\
 &= \sin 5x \times \frac{d}{dx}(e^x) + e^x \frac{d}{dx}(\sin 5x) \\
 &= \sin 5x \cdot e^x + e^x \cdot \cos 5x \cdot \frac{d}{dx}(5x) \\
 &= e^x \sin 5x + e^x \cos 5x \cdot 5 \\
 &= e^x (\sin 5x + 5 \cos 5x)
 \end{aligned}$$

Therefore,

$$\begin{aligned}
 \frac{d^2y}{dx^2} &= \frac{d}{dx}[e^x (\sin 5x + 5 \cos 5x)] \\
 &= (\sin 5x + 5 \cos 5x) \cdot \frac{d}{dx}(e^x) + e^x \cdot \frac{d}{dx}(\sin 5x + 5 \cos 5x) \\
 &= (\sin 5x + 5 \cos 5x) e^x + e^x \left[ \cos 5x \cdot \frac{d}{dx}(5x) + 5(-\sin 5x) \cdot \frac{d}{dx}(5x) \right] \\
 &= e^x (\sin 5x + 5 \cos 5x) + e^x (5 \cos 5x - 25 \sin 5x) \\
 &= e^x (10 \cos 5x - 24 \sin 5x) \\
 &= 2e^x (5 \cos 5x - 12 \sin 5x)
 \end{aligned}$$

### Question 7:

Find the second order derivative of the function  $e^{6x} \cos 3x$

### Solution:

Let  $y = e^{6x} \cos 3x$

Then,

$$\begin{aligned}
 \frac{dy}{dx} &= \frac{d}{dx}(e^{6x} \cos 3x) = \cos 3x \cdot \frac{d}{dx}(e^{6x}) + e^{6x} \cdot \frac{d}{dx}(\cos 3x) \\
 &= \cos 3x \cdot e^{6x} \cdot \frac{d}{dx}(6x) + e^{6x} \cdot (-\sin 3x) \cdot \frac{d}{dx}(3x) \\
 &= 6e^{6x} \cos 3x - 3e^{6x} \sin 3x \quad \dots(1)
 \end{aligned}$$

Therefore,

$$\begin{aligned}
 \frac{d^2 y}{dx^2} &= \frac{d}{dx} (6e^{6x} \cos 3x - 3e^{6x} \sin 3x) = 6 \cdot \frac{d}{dx} (e^{6x} \cos 3x) - 3 \cdot \frac{d}{dx} (e^{6x} \sin 3x) \\
 &= 6 \cdot [6e^{6x} \cos 3x - 3e^{6x} \sin 3x] - 3 \cdot \left[ \sin 3x \cdot \frac{d}{dx} (e^{6x}) + e^{6x} \cdot \frac{d}{dx} (\sin 3x) \right] \quad [\text{using (1)}] \\
 &= 36e^{6x} \cos 3x - 18e^{6x} \sin 3x - 3 \left[ \sin 3x \cdot e^{6x} \cdot 6 + e^{6x} \cdot \cos 3x \cdot 3 \right] \\
 &= 36e^{6x} \cos 3x - 18e^{6x} \sin 3x - 18e^{6x} \sin 3x - 9e^{6x} \cos 3x \\
 &= 27e^{6x} \cos 3x - 36e^{6x} \sin 3x \\
 &= 9e^{6x} (3 \cos 3x - 4 \sin 3x)
 \end{aligned}$$

### Question 8:

Find the second order derivative of the function  $\tan^{-1} x$

#### Solution:

Let  $y = \tan^{-1} x$

Then,

$$\begin{aligned}
 \frac{dy}{dx} &= \frac{d}{dx} (\tan^{-1} x) \\
 &= \frac{1}{1+x^2}
 \end{aligned}$$

Therefore,

$$\begin{aligned}
 \frac{d^2 y}{dx^2} &= \frac{d}{dx} \left( \frac{1}{1+x^2} \right) = \frac{d}{dx} (1+x^2)^{-1} \\
 &= (-1) \cdot (1+x^2)^{-2} \cdot \frac{d}{dx} (1+x^2) = \frac{-1}{(1+x^2)^2} \times 2x \\
 &= \frac{-2x}{(1+x^2)^2}
 \end{aligned}$$

### Question 9:

Find the second order derivative of the function  $\log(\log x)$

#### Solution:

Consider,  $y = \log(\log x)$

Then,

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx} [\log(\log x)] \\ &= \frac{1}{\log x} \cdot \frac{d}{dx} (\log x) \\ &= \frac{1}{\log x} \cdot \frac{1}{x} = (x \log x)^{-1}\end{aligned}$$

Therefore,

$$\begin{aligned}\frac{d^2y}{dx^2} &= \frac{d}{dx} [(x \log x)^{-1}] \\ &= (-1) \cdot (x \log x)^{-2} \cdot \frac{d}{dx} (x \log x) \\ &= \frac{-1}{(x \log x)^2} \cdot \left[ \log x \cdot \frac{d}{dx} (x) + x \cdot \frac{d}{dx} (\log x) \right] \\ &= \frac{-1}{(x \log x)^2} \cdot \left[ \log x \cdot 1 + x \cdot \frac{1}{x} \right] \\ &= \frac{-(1 + \log x)}{(x \log x)^2}\end{aligned}$$

### Question 10:

Find the second order derivative of the function  $\sin(\log x)$

### Solution:

Let  $y = \sin(\log x)$

Then,

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx} [\sin(\log x)] \\ &= \cos(\log x) \cdot \frac{d}{dx} (\log x) \\ &= \frac{\cos(\log x)}{x}\end{aligned}$$

Therefore,

$$\begin{aligned}\frac{d^2y}{dx^2} &= \frac{d}{dx} \left[ \frac{\cos(\log x)}{x} \right] \\ &= \frac{x \cdot \frac{d}{dx} [\cos(\log x)] - \cos(\log x) \cdot \frac{d}{dx} (x)}{x^2}\end{aligned}$$

$$\begin{aligned}\frac{d^2y}{dx^2} &= \frac{x \left[ -\sin(\log x) \cdot \frac{d}{dx}(\log x) \right] - \cos(\log x) \cdot 1}{x^2} \\ &= \frac{-x \sin(\log x) \cdot \frac{1}{x} - \cos(\log x)}{x^2} \\ &= \frac{-[\sin(\log x) + \cos(\log x)]}{x^2}\end{aligned}$$

### Question 11:

If  $y = 5 \cos x - 3 \sin x$ , prove that  $\frac{d^2y}{dx^2} + y = 0$

### Solution:

Given,  $y = 5 \cos x - 3 \sin x$

Then,

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx}(5 \cos x) - \frac{d}{dx}(3 \sin x) \\ &= 5 \frac{d}{dx}(\cos x) - 3 \frac{d}{dx}(\sin x) \\ &= 5(-\sin x) - 3 \cos x \\ &= -(5 \sin x + 3 \cos x)\end{aligned}$$

Therefore,

$$\begin{aligned}\frac{d^2y}{dx^2} &= \frac{d}{dx}[-(5 \sin x + 3 \cos x)] \\ &= -\left[ 5 \cdot \frac{d}{dx}(\sin x) + 3 \cdot \frac{d}{dx}(\cos x) \right] \\ &= -[5 \cos x + 3(-\sin x)] \\ &= -[5 \cos x - 3 \sin x] \\ &= -y\end{aligned}$$

Thus,  $\frac{d^2y}{dx^2} + y = 0$

Hence proved.

### Question 12:

If  $y = \cos^{-1} x$ , find  $\frac{d^2 y}{dx^2}$  in terms of  $y$  alone.

### Solution:

Given,  $y = \cos^{-1} x$

Then,

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx}(\cos^{-1} x) \\ &= \frac{-1}{\sqrt{1-x^2}} \\ &= -(1-x^2)^{-\frac{1}{2}}\end{aligned}$$

Therefore,

$$\begin{aligned}\frac{d^2 y}{dx^2} &= \frac{d}{dx} \left[ -(1-x^2)^{-\frac{1}{2}} \right] \\ &= - \left( -\frac{1}{2} \right) \cdot (1-x^2)^{-\frac{3}{2}} \cdot \frac{d}{dx}(1-x^2) \\ &= \frac{1}{2\sqrt{(1-x^2)^3}} \times (-2x) \\ \frac{d^2 y}{dx^2} &= \frac{-x}{\sqrt{(1-x^2)^3}} \quad \dots(1)\end{aligned}$$

But we need to calculate  $\frac{d^2 y}{dx^2}$  in terms of  $y$

$$\Rightarrow y = \cos^{-1} x$$

$$\Rightarrow x = \cos y$$

Putting  $x = \cos y$  in equation (1), we get

$$\begin{aligned}\frac{d^2y}{dx^2} &= \frac{-\cos y}{\sqrt{(1-\cos^2 y)^3}} \\ &= \frac{-\cos y}{\sqrt{(\sin^2 y)^3}} \\ &= \frac{-\cos y}{\sin^3 y} \\ &= \frac{-\cos y}{\sin y} \times \frac{1}{\sin^2 y} \\ &= -\cot y \cdot \operatorname{cosec}^2 y\end{aligned}$$

### Question 13:

If  $y = 3 \cos(\log x) + 4 \sin(\log x)$ , show that  $x^2 y_2 + x y_1 + y = 0$

### Solution:

Given,  $y = 3 \cos(\log x) + 4 \sin(\log x)$

Then,

$$\begin{aligned}y_1 &= 3 \cdot \frac{d}{dx} [\cos(\log x)] + 4 \cdot \frac{d}{dx} [\sin(\log x)] \\ &= 3 \cdot \left[ -\sin(\log x) \cdot \frac{d}{dx} (\log x) \right] + 4 \cdot \left[ \cos(\log x) \cdot \frac{d}{dx} (\log x) \right] \\ &= \frac{-3 \sin(\log x)}{x} + \frac{4 \cos(\log x)}{x} \\ &= \frac{4 \cos(\log x) - 3 \sin(\log x)}{x}\end{aligned}$$

Therefore,

$$\begin{aligned}
 y_2 &= \frac{d}{dx} \left( \frac{4 \cos(\log x) - 3 \sin(\log x)}{x} \right) \\
 &= \frac{x \cdot \{4 \cos(\log x) - 3 \sin(\log x)\}' - \{4 \cos(\log x) - 3 \sin(\log x)\} \{x\}'}{x^2} \\
 &= \frac{x \cdot [4 \{\cos(\log x)\}' - 3 \{\sin(\log x)\}'] - \{4 \cos(\log x) - 3 \sin(\log x)\} \cdot 1}{x^2} \\
 &= \frac{x \cdot [-4 \sin(\log x) \cdot (\log x)' - 3 \cos(\log x) \cdot (\log x)'] - 4 \cos(\log x) + 3 \sin(\log x)}{x^2} \\
 &= \frac{x \cdot \left[ -4 \sin(\log x) \cdot \frac{1}{x} - 3 \cos(\log x) \cdot \frac{1}{x} \right] - 4 \cos(\log x) + 3 \sin(\log x)}{x^2} \\
 &= \frac{-4 \sin(\log x) - 3 \cos(\log x) - 4 \cos(\log x) + 3 \sin(\log x)}{x^2} \\
 &= \frac{-\sin(\log x) - 7 \cos(\log x)}{x^2}
 \end{aligned}$$

Thus,

$$\begin{aligned}
 x^2 y_2 + x y_1 + y &= \left[ x^2 \left( \frac{-\sin(\log x) - 7 \cos(\log x)}{x^2} \right) + x \left( \frac{4 \cos(\log x) - 3 \sin(\log x)}{x} \right) \right] \\
 &\quad + 3 \cos(\log x) + 4 \sin(\log x) \\
 &= \left[ -\sin(\log x) - 7 \cos(\log x) + 4 \cos(\log x) - 3 \sin(\log x) \right] \\
 &\quad + 3 \cos(\log x) + 4 \sin(\log x) \\
 &= 0
 \end{aligned}$$

Hence proved.

#### Question 14:

If  $y = Ae^{mx} + Be^{nx}$ , show that  $\frac{d^2 y}{dx^2} - (m+n) \frac{dy}{dx} + mny = 0$ .

**Solution:**

Given,  $y = Ae^{mx} + Be^{nx}$

Then,

$$\begin{aligned}
 \frac{dy}{dx} &= A \cdot \frac{d}{dx} (e^{mx}) + B \cdot \frac{d}{dx} (e^{nx}) \\
 &= A e^{mx} \cdot \frac{d}{dx} (mx) + B e^{nx} \cdot \frac{d}{dx} (nx) \\
 &= A m e^{mx} + B n e^{nx}
 \end{aligned}$$



Therefore,

$$\begin{aligned}\frac{d^2y}{dx^2} &= \frac{d}{dx}(Ame^{mx} + Bne^{nx}) \\ &= Am \cdot \frac{d}{dx}(e^{mx}) + Bn \cdot \frac{d}{dx}(e^{nx}) \\ &= Am \cdot e^{mx} \cdot \frac{d}{dx}(mx) + Bn \cdot e^{nx} \cdot \frac{d}{dx}(nx) \\ &= Am^2e^{mx} + Bn^2e^{nx}\end{aligned}$$

Thus,

$$\begin{aligned}\frac{d^2y}{dx^2} - (m+n) \frac{dy}{dx} + mny &= Am^2e^{mx} + Bn^2e^{nx} - (m+n) \cdot (Ame^{mx} + Bne^{nx}) + mn(Ae^{mx} + Be^{nx}) \\ &= Am^2e^{mx} + Bn^2e^{nx} - Am^2e^{mx} - Bmne^{nx} - Amne^{mx} - Bn^2e^{nx} + Amne^{mx} + Bmne^{nx} \\ &= 0\end{aligned}$$

Hence proved.

### Question 15:

If  $y = 500e^{7x} + 600e^{-7x}$ , show that  $\frac{d^2y}{dx^2} = 49y$

**Solution:**

Given,  $y = 500e^{7x} + 600e^{-7x}$

Then,

$$\begin{aligned}\frac{dy}{dx} &= 500 \cdot \frac{d}{dx}(e^{7x}) + 600 \cdot \frac{d}{dx}(e^{-7x}) \\ &= 500 \cdot e^{7x} \cdot \frac{d}{dx}(7x) + 600 \cdot e^{-7x} \cdot \frac{d}{dx}(-7x) \\ &= 3500e^{7x} - 4200e^{-7x}\end{aligned}$$

Therefore,

$$\begin{aligned}\frac{d^2y}{dx^2} &= 3500e^{7x} \cdot \frac{d}{dx}(e^{7x}) - 4200 \cdot \frac{d}{dx}(e^{-7x}) \\ &= 3500 \cdot e^{7x} \cdot \frac{d}{dx}(7x) - 4200 \cdot e^{-7x} \cdot \frac{d}{dx}(-7x) \\ &= 7 \times 3500 \cdot e^{7x} + 7 \times 4200 \cdot e^{-7x} \\ &= 49 \times 500 \cdot e^{7x} + 49 \times 600 \cdot e^{-7x} \\ &= 49(500e^{7x} + 600e^{-7x}) \\ &= 49y\end{aligned}$$

Hence proved.

### Question 16:

If  $e^y(x+1)=1$ , show that  $\frac{d^2y}{dx^2} = \left(\frac{dy}{dx}\right)^2$

#### Solution:

Given,  $e^y(x+1)=1$

$$\Rightarrow e^y(x+1)=1$$

$$\Rightarrow e^y = \frac{1}{x+1}$$

Taking log on both sides, we get

$$y = \log \frac{1}{(x+1)}$$

Differentiating with respect to  $x$ , we get

$$\begin{aligned} \frac{dy}{dx} &= (x+1) \frac{d}{dx} \left( \frac{1}{x+1} \right) \\ &= (x+1) \cdot \frac{-1}{(x+1)^2} \\ &= \frac{-1}{x+1} \end{aligned}$$

Therefore,

$$\begin{aligned} \frac{d^2y}{dx^2} &= \frac{d}{dx} \left( \frac{-1}{x+1} \right) = - \left( \frac{-1}{(x+1)^2} \right) \\ &= \frac{1}{(x+1)^2} = \left( \frac{-1}{x+1} \right)^2 \\ &= \left( \frac{dy}{dx} \right)^2 \end{aligned}$$

Hence proved.

### Question 17:

If  $y = (\tan^{-1} x)^2$ , show that  $(x^2+1)^2 y_2 + 2x(x^2+1)y_1 = 2$

#### Solution:

Given,  $y = (\tan^{-1} x)^2$



Then,

$$\Rightarrow y_1 = 2 \tan^{-1} x \frac{d}{dx} (\tan^{-1} x)$$

$$\Rightarrow y_1 = 2 \tan^{-1} x \cdot \left( \frac{1}{1+x^2} \right)$$

$$\Rightarrow (1+x^2) y_1 = 2 \tan^{-1} x$$

Again, differentiating with respect to  $x$ , we get

$$\Rightarrow (1+x^2) y_2 + 2xy_1 = 2 \left( \frac{1}{1+x^2} \right)$$

$$\Rightarrow (1+x^2)^2 y_2 + 2x(1+x^2) y_1 = 2$$

Hence proved.



MillionStars edu  
Think, Learn & Practice

## EXERCISE 5.8

### Question 1:

Verify Rolle's Theorem for the function  $f(x) = x^2 + 2x - 8, x \in [-4, 2]$

### Solution:

Given,  $f(x) = x^2 + 2x - 8$ , being polynomial function is continuous in  $[-4, 2]$  and also differentiable in  $(-4, 2)$ .

$$\begin{aligned} f(-4) &= (-4)^2 + 2(-4) - 8 \\ &= 16 - 8 - 8 \\ &= 0 \end{aligned}$$

$$\begin{aligned} f(2) &= (2)^2 + 2 \times 2 - 8 \\ &= 4 + 4 - 8 \\ &= 0 \end{aligned}$$

Therefore,  $f(-4) = f(2) = 0$

The value of  $f(x)$  at  $-4$  and  $2$  coincides.

Rolle's Theorem states that there is a point  $c \in (-4, 2)$  such that  $f'(c) = 0$

$$f(x) = x^2 + 2x - 8$$

Therefore,  $f'(x) = 2x + 2$

Hence,

$$f'(c) = 0$$

$$2c + 2 = 0$$

$$c = -1$$

Thus,  $c = -1 \in (-4, 2)$

Hence, Rolle's Theorem is verified.

### Question 2:

Examine if Rolle's Theorem is applicable to any of the following functions. Can you say something about the converse of Rolle's Theorem from these examples?

(i)  $f(x) = [x]$  for  $x \in [5, 9]$

- (ii)  $f(x) = [x]$  for  $x \in [-2, 2]$   
 (iii)  $f(x) = x^2 - 1$  for  $x \in [1, 2]$

### Solution:

By Rolle's Theorem,  $f: [a, b] \rightarrow \mathbf{R}$ ,

If

- (a)  $f$  is continuous on  $[a, b]$   
 (b)  $f$  is continuous on  $(a, b)$   
 (c)  $f(a) = f(b)$

Then, there exists some  $c \in (a, b)$  such that  $f'(c) = 0$

Thus, Rolle's Theorem is not applicable to those functions that do not satisfy any of three conditions of the hypothesis.

- (i)  $f(x) = [x]$  for  $x \in [5, 9]$

Since, the given function  $f(x)$  is not continuous at every integral point.

In general,  $f(x)$  is not continuous at  $x = 5$  and  $x = 9$

Therefore,  $f(x)$  is not continuous in  $[5, 9]$

Also,  $f(5) = [5] = 5$  and  $f(9) = [9] = 9$

Thus,  $f(5) \neq f(9)$

The differentiability of  $f$  in  $(5, 9)$  is checked as follows.

Let  $n$  be an integer such that  $n \in (5, 9)$

The LHD of  $f$  at  $x = n$  is

$$\lim_{h \rightarrow 0^-} \frac{f(n+h) - f(n)}{h} = \lim_{h \rightarrow 0^-} \frac{[n+h] - [n]}{h} = \lim_{h \rightarrow 0^-} \frac{n-1-n}{h} = \lim_{h \rightarrow 0^-} \frac{-1}{h} = \infty$$

The RHD of  $f$  at  $x = n$  is

$$\lim_{h \rightarrow 0^+} \frac{f(n+h) - f(n)}{h} = \lim_{h \rightarrow 0^+} \frac{[n+h] - [n]}{h} = \lim_{h \rightarrow 0^+} \frac{n-n}{h} = \lim_{h \rightarrow 0^+} 0 = 0$$

Since LHD and RHD of  $f$  at  $x = n$  are not equal,  $f$  is not differentiable at  $x = n$

Therefore,  $f$  is not differentiable in  $(5, 9)$ .

It is observed that  $f$  does not satisfy all the conditions of the hypothesis of Rolle's Theorem.

Thus, Rolle's Theorem is not applicable for  $f(x) = [x]$  for  $x \in [5, 9]$ .

(ii)  $f(x) = [x]$  for  $x \in [-2, 2]$

Since, the given function  $f(x)$  is not continuous at every integral point.

In general,  $f(x)$  is not continuous at  $x = -2$  and  $x = 2$

Therefore,  $f(x)$  is not continuous in  $[-2, 2]$

Also,  $f(-2) = [-2] = -2$  and  $f(2) = [2] = 2$

Thus,  $f(-2) \neq f(2)$

The differentiability of  $f$  in  $(-2, 2)$  is checked as follows.

Let  $n$  be an integer such that  $n \in (-2, 2)$

The LHD of  $f$  at  $x = n$  is

$$\lim_{h \rightarrow 0^-} \frac{f(n+h) - f(n)}{h} = \lim_{h \rightarrow 0^-} \frac{[n+h] - [n]}{h} = \lim_{h \rightarrow 0^-} \frac{n-1-n}{h} = \lim_{h \rightarrow 0^-} \frac{-1}{h} = \infty$$

The RHD of  $f$  at  $x = n$  is

$$\lim_{h \rightarrow 0^+} \frac{f(n+h) - f(n)}{h} = \lim_{h \rightarrow 0^+} \frac{[n+h] - [n]}{h} = \lim_{h \rightarrow 0^+} \frac{n-1-n}{h} = \lim_{h \rightarrow 0^+} 0 = 0$$

Since LHD and RHD of  $f$  at  $x = n$  are not equal,  $f$  is not differentiable at  $x = n$

Therefore,  $f$  is not differentiable in  $(-2, 2)$ .

It is observed that  $f$  does not satisfy all the conditions of the hypothesis of Rolle's Theorem.

Thus, Rolle's Theorem is not applicable for  $f(x) = [x]$  for  $x \in [-2, 2]$

(iii)  $f(x) = x^2 - 1$  for  $x \in [1, 2]$

Since,  $f$  being a polynomial function is continuous in  $[1, 2]$  and is differentiable in  $(1, 2)$

Thus,

$$f(1) = (1)^2 - 1 = 0$$

$$f(2) = (2)^2 - 1 = 3$$

Therefore,  $f(1) \neq f(2)$

Since,  $f$  does not satisfy a condition of the hypothesis of Rolle's Theorem.

Hence, Rolle's Theorem is not applicable for  $f(x) = x^2 - 1$  for  $x \in [1, 2]$ .

### Question 3:

If  $f: [-5, 5] \rightarrow \mathbf{R}$  is a differentiable function and if  $f'(x)$  does not vanish anywhere, then prove that  $f(-5) \neq f(5)$ .

### Solution:

Given,  $f: [-5, 5] \rightarrow \mathbf{R}$  is a differentiable function.

Since every differentiable function is a continuous function, we obtain

- (i)  $f$  is continuous on  $[-5, 5]$
- (ii)  $f$  is continuous on  $(-5, 5)$

Thus, by the Mean Value Theorem, there exists  $c \in (-5, 5)$  such that

$$\Rightarrow f'(c) = \frac{f(5) - f(-5)}{5 - (-5)}$$

$$\Rightarrow 10f'(c) = f(5) - f(-5)$$

It is also given that  $f'(x)$  does not vanish anywhere.

Therefore,  $f'(c) \neq 0$

Thus,

$$\Rightarrow 10f'(c) \neq 0$$

$$\Rightarrow f(5) - f(-5) \neq 0$$

$$\Rightarrow f(5) \neq f(-5)$$

Hence proved.

### Question 4:

Verify Mean Value Theorem, if  $f(x) = x^2 - 4x - 3$  in the interval  $[a, b]$ , where  $a = 1$  and  $b = 4$ .

### Solution:

Given,  $f(x) = x^2 - 4x - 3$

$f$ , being a polynomial function, is continuous in  $[1, 4]$  and is differentiable in  $(1, 4)$ , whose derivative is  $2x - 4$ .

Thus,

$$f(1) = 1^2 - 4 \times 1 - 3 = -6$$

$$f(4) = 4^2 - 4 \times 4 - 3 = -3$$

Therefore,

$$\begin{aligned} \frac{f(b) - f(a)}{b - a} &= \frac{f(4) - f(1)}{4 - 1} \\ &= \frac{-3 - (-6)}{3} \\ &= \frac{3}{3} \\ &= 1 \end{aligned}$$

Mean Value Theorem states that there is a point  $c \in (1, 4)$  such that  $f'(c) = 1$

Hence,

$$\Rightarrow f'(c) = 1$$

$$\Rightarrow 2c - 4 = 1$$

$$\Rightarrow c = \frac{5}{2} \quad \left[ \text{where } c = \frac{5}{2} \in (1, 4) \right]$$

Thus, mean value theorem is verified for the given function.

### Question 5:

Verify Mean Value Theorem, if  $f(x) = x^3 - 5x^2 - 3x$  in the interval  $[a, b]$  where  $a = 1$  and  $b = 3$ .  
Find all  $c \in (1, 3)$  for which  $f'(c) = 0$ .

### Solution:

Given,  $f$  is  $f(x) = x^3 - 5x^2 - 3x$

$f$ , being a polynomial function, is continuous in  $[1, 3]$  and is differentiable in  $(1, 3)$ , whose derivative is  $3x^2 - 10x - 3$

Thus,

$$f(1) = 1^3 - 5 \times 1^2 - 3 \times 1 = -7$$

$$f(3) = 3^3 - 5 \times 3^2 - 3 \times 3 = -27$$

Therefore,



$$\begin{aligned}\frac{f(b)-f(a)}{b-a} &= \frac{f(3)-f(1)}{3-1} \\ &= \frac{-27-(-7)}{3-1} \\ &= -10\end{aligned}$$

Mean Value Theorem states that there exists a point  $c \in (1,3)$  such that  $f'(c) = -10$   
Hence,

$$\begin{aligned}\Rightarrow f'(c) &= -10 \\ \Rightarrow 3c^2 - 10c - 3 &= -10 \\ \Rightarrow 3c^2 - 10c + 7 &= 0 \\ \Rightarrow 3c^2 - 3c - 7c + 7 &= 0 \\ \Rightarrow 3c(c-1) - 7(c-1) &= 0 \\ \Rightarrow (c-1)(3c-7) &= 0 \\ \Rightarrow c = 1, \frac{7}{3} &\quad \left[ \text{where } c = \frac{7}{3} \in (1,3) \right]\end{aligned}$$

Thus, Mean Value Theorem is verified for the given function and  $c = \frac{7}{3} \in (1,3)$  is the only point for which  $f'(c) = 0$ .

### Question 6:

Examine the applicability of Mean Value Theorem for all three functions given

- (i)  $f(x) = [x]$  for  $x \in [5,9]$
- (ii)  $f(x) = [x]$  for  $x \in [-2,2]$
- (iii)  $f(x) = x^2 - 1$  for  $x \in [1,2]$

### Solution:

Mean Value Theorem states that for a function  $f: [a,b] \rightarrow \mathbf{R}$ , if

- (a)  $f$  is continuous on  $[a,b]$
- (b)  $f$  is continuous on  $(a,b)$

Then there exists some  $c \in (a,b)$  such that  $f'(c) = \frac{f(b)-f(a)}{b-a}$

Thus, Mean Value Theorem is not applicable to those functions that do not satisfy any of three conditions of the hypothesis.

- (i)  $f(x) = [x]$  for  $x \in [5,9]$

Since, the given function  $f(x)$  is not continuous at every integral point.

In general,  $f(x)$  is not continuous at  $x=5$  and  $x=9$

Therefore,  $f(x)$  is not continuous in  $[5,9]$

The differentiability of  $f$  in  $(5,9)$  is checked as follows.

Let  $n$  be an integer such that  $n \in (5,9)$

The LHD of  $f$  at  $x=n$  is

$$\lim_{h \rightarrow 0^-} \frac{f(n+h) - f(n)}{h} = \lim_{h \rightarrow 0^-} \frac{[n+h] - [n]}{h} = \lim_{h \rightarrow 0^-} \frac{n-1-n}{h} = \lim_{h \rightarrow 0^-} \frac{-1}{h} = \infty$$

The RHD of  $f$  at  $x=n$  is

$$\lim_{h \rightarrow 0^+} \frac{f(n+h) - f(n)}{h} = \lim_{h \rightarrow 0^+} \frac{[n+h] - [n]}{h} = \lim_{h \rightarrow 0^+} \frac{n-n}{h} = \lim_{h \rightarrow 0^+} 0 = 0$$

Since LHD and RHD of  $f$  at  $x=n$  are not equal,  $f$  is not differentiable at  $x=n$

Therefore,  $f$  is not differentiable in  $(5,9)$ .

It is observed that  $f$  does not satisfy all the conditions of the hypothesis of Mean Value Theorem.

Thus, Mean Value Theorem is not applicable for  $f(x)=[x]$  for  $x \in [5,9]$

(ii)  $f(x)=[x]$  for  $x \in [-2,2]$

Since, the given function  $f(x)$  is not continuous at every integral point.

In general,  $f(x)$  is not continuous at  $x=-2$  and  $x=2$

Therefore,  $f(x)$  is not continuous in  $[-2,2]$

The differentiability of  $f$  in  $(-2,2)$  is checked as follows.

Let  $n$  be an integer such that  $n \in (-2,2)$

The LHD of  $f$  at  $x=n$  is

$$\lim_{h \rightarrow 0^-} \frac{f(n+h) - f(n)}{h} = \lim_{h \rightarrow 0^-} \frac{[n+h] - [n]}{h} = \lim_{h \rightarrow 0^-} \frac{n-1-n}{h} = \lim_{h \rightarrow 0^-} \frac{-1}{h} = \infty$$

The RHD of  $f$  at  $x = n$  is

$$\lim_{h \rightarrow 0^+} \frac{f(n+h) - f(n)}{h} = \lim_{x \rightarrow 0^+} \frac{[n+h] - [n]}{h} = \lim_{h \rightarrow 0^+} \frac{n-n}{h} = \lim_{h \rightarrow 0^+} 0 = 0$$

Since LHD and RHD of  $f$  at  $x = n$  are not equal,  $f$  is not differentiable at  $x = n$

Therefore,  $f$  is not differentiable in  $(-2, 2)$ .

It is observed that  $f$  does not satisfy all the conditions of the hypothesis of Mean Value Theorem.

Thus, Mean Value Theorem is not applicable for  $f(x) = [x]$  for  $x \in [-2, 2]$ .

(iii)  $f(x) = x^2 - 1$  for  $x \in [1, 2]$

Since,  $f$  being a polynomial function is continuous in  $[1, 2]$  and is differentiable in  $(1, 2)$

It is observed that  $f$  satisfies all the conditions of the hypothesis of Mean Value Theorem.

Hence, Mean Value Theorem is applicable for  $f(x) = x^2 - 1$  for  $x \in [1, 2]$ .

It can be proved as follows.

We have,  $f(x) = x^2 - 1$

Then,

$$f(1) = (1)^2 - 1 = 0,$$

$$f(2) = (2)^2 - 1 = 3$$

Therefore,

$$\frac{f(b) - f(a)}{b - a} = \frac{f(2) - f(1)}{2 - 1} = \frac{3 - 0}{1} = 3$$

Hence,  $f'(x) = 2x$

Thus,

$$\Rightarrow f'(c) = 3$$

$$\Rightarrow 2c = 3$$

$$\Rightarrow c = \frac{3}{2}$$

$$\Rightarrow c = 1.5 \quad [\text{where } 1.5 \in [1, 2]]$$

## MISCELLANEOUS EXERCISE

### Question 1:

Differentiate with respect to  $x$  the function  $(3x^2 - 9x + 5)^9$ .

### Solution:

Let  $y = (3x^2 - 9x + 5)^9$

Using chain rule, we get

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx} (3x^2 - 9x + 5)^9 \\ &= 9(3x^2 - 9x + 5)^8 \cdot \frac{d}{dx} (3x^2 - 9x + 5) \\ &= 9(3x^2 - 9x + 5)^8 \cdot (6x - 9) \\ &= 9(3x^2 - 9x + 5)^8 \cdot 3(2x - 3) \\ &= 27(3x^2 - 9x + 5)^8 (2x - 3)\end{aligned}$$

### Question 2:

Differentiate with respect to  $x$  the function  $\sin^3 x + \cos^6 x$ .

### Solution:

Let  $y = \sin^3 x + \cos^6 x$

Using chain rule, we get

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx} (\sin^3 x) + \frac{d}{dx} (\cos^6 x) \\ &= 3\sin^2 x \cdot \frac{d}{dx} (\sin x) + 6\cos^5 x \cdot \frac{d}{dx} (\cos x) \\ &= 3\sin^2 x \cdot \cos x + 6\cos^5 x \cdot (-\sin x) \\ &= 3\sin x \cos x (\sin x - 2\cos^4 x)\end{aligned}$$

### Question 3:

Differentiate with respect to  $x$  the function  $(5x)^{3\cos 2x}$ .

**Solution:**

Let  $y = (5x)^{3 \cos 2x}$

Taking logarithm on both the sides, we obtain

$$\log y = 3 \cos 2x \log 5x$$

Differentiating both sides with respect to  $x$ , we get

$$\begin{aligned} \frac{1}{y} \frac{dy}{dx} &= 3 \left[ \log 5x \cdot \frac{d}{dx} (\cos 2x) + \cos 2x \cdot \frac{d}{dx} (\log 5x) \right] \\ \frac{dy}{dx} &= 3y \left[ \log 5x \cdot (-\sin 2x) \cdot \frac{d}{dx} (2x) + \cos 2x \cdot \frac{1}{5x} \cdot \frac{d}{dx} (5x) \right] \\ &= 3y \left[ -2 \sin 2x \cdot \log 5x + \frac{\cos 2x}{x} \right] \\ &= y \left[ \frac{3 \cos 2x}{x} - 6 \sin 2x \log 5x \right] \\ &= (5x)^{3 \cos 2x} \left[ \frac{3 \cos 2x}{x} - 6 \sin 2x \log 5x \right] \end{aligned}$$

**Question 4:**

Differentiate with respect to  $x$  the function  $\sin^{-1}(x\sqrt{x})$ ,  $0 \leq x \leq 1$ .

**Solution:**

Let  $y = \sin^{-1}(x\sqrt{x})$

Using chain rule, we get

$$\begin{aligned}
 \frac{dy}{dx} &= \frac{d}{dx} \sin^{-1}(x\sqrt{x}) \\
 &= \frac{1}{\sqrt{1-(x\sqrt{x})^2}} \times \frac{d}{dx}(x\sqrt{x}) \\
 &= \frac{1}{\sqrt{1-x^3}} \cdot \frac{d}{dx}\left(x^{\frac{3}{2}}\right) \\
 &= \frac{1}{\sqrt{1-x^3}} \cdot \frac{3}{2} x^{\frac{1}{2}} \\
 &= \frac{3\sqrt{x}}{2\sqrt{1-x^3}} \\
 &= \frac{3}{2} \sqrt{\frac{x}{1-x^3}}
 \end{aligned}$$

### Question 5:

Differentiate with respect to  $x$  the function  $\frac{\cos^{-1} \frac{x}{2}}{\sqrt{2x+7}}, -2 < x < 2$ .

### Solution:

Let  $y = \frac{\cos^{-1} \frac{x}{2}}{\sqrt{2x+7}}$

Using quotient rule, we get

$$\begin{aligned}
 \frac{dy}{dx} &= \frac{\sqrt{2x+7} \cdot \frac{d}{dx} \left( \cos^{-1} \frac{x}{2} \right) - \left( \cos^{-1} \frac{x}{2} \right) \cdot \frac{d}{dx} (\sqrt{2x+7})}{(\sqrt{2x+7})^2} \\
 &= \frac{\sqrt{2x+7} \left[ \frac{-1}{\sqrt{1 - \left( \frac{x}{2} \right)^2}} \cdot \frac{d}{dx} \left( \frac{x}{2} \right) \right] - \left( \cos^{-1} \frac{x}{2} \right) \cdot \frac{1}{2\sqrt{2x+7}} \cdot \frac{d}{dx} (2x+7)}{2x+7} \\
 &= \frac{\sqrt{2x+7} \cdot \frac{-1}{\sqrt{4-x^2}} - \left( \cos^{-1} \frac{x}{2} \right) \cdot \frac{2}{2\sqrt{2x+7}}}{2x+7} \\
 &= \frac{-\sqrt{2x+7}}{(\sqrt{4-x^2}) \cdot (2x+7)} - \frac{\cos^{-1} \frac{x}{2}}{(\sqrt{2x+7}) (2x+7)} \\
 &= - \left[ \frac{1}{\sqrt{4-x^2} \sqrt{2x+7}} + \frac{\cos^{-1} \frac{x}{2}}{(2x+7)^{\frac{3}{2}}} \right]
 \end{aligned}$$

### Question 6:

Differentiate with respect to  $x$  the function  $\cot^{-1} \left[ \frac{\sqrt{1+\sin x} + \sqrt{1-\sin x}}{\sqrt{1+\sin x} - \sqrt{1-\sin x}} \right], 0 < x < \frac{\pi}{2}$ .

### Solution:

Let  $y = \cot^{-1} \left[ \frac{\sqrt{1+\sin x} + \sqrt{1-\sin x}}{\sqrt{1+\sin x} - \sqrt{1-\sin x}} \right] \quad \dots(1)$

Then,

$$\begin{aligned}
 \frac{\sqrt{1+\sin x} + \sqrt{1-\sin x}}{\sqrt{1+\sin x} - \sqrt{1-\sin x}} &= \frac{(\sqrt{1+\sin x} + \sqrt{1-\sin x})^2}{(\sqrt{1+\sin x} - \sqrt{1-\sin x})(\sqrt{1+\sin x} + \sqrt{1-\sin x})} \\
 &= \frac{(1+\sin x) + (1-\sin x) + 2\sqrt{(1+\sin x)(1-\sin x)}}{(1+\sin x) - (1-\sin x)} \\
 &= \frac{2 + 2\sqrt{1-\sin^2 x}}{2\sin x} = \frac{1 + \cos x}{\sin x} \\
 &= \frac{1 + 2\cos^2 \frac{x}{2} - 1}{2\sin \frac{x}{2} \cos \frac{x}{2}} = \frac{2\cos^2 \frac{x}{2}}{2\sin \frac{x}{2} \cos \frac{x}{2}} \\
 &= \cot \frac{x}{2}
 \end{aligned}$$

Therefore, equation (1) becomes,

$$y = \cot^{-1} \left( \cot \frac{x}{2} \right)$$

$$\Rightarrow y = \frac{x}{2}$$

Thus,

$$\begin{aligned}
 \Rightarrow \frac{dy}{dx} &= \frac{1}{2} \frac{d}{dx} (x) \\
 &= \frac{1}{2}
 \end{aligned}$$

### Question 7:

Differentiate with respect to  $x$  the function  $(\log x)^{\log x}$ ,  $x > 1$ .

### Solution:

$$\text{Let } y = (\log x)^{\log x}$$

Taking logarithm on both the sides, we obtain

$$\log y = \log x \cdot \log(\log x)$$



Differentiating both sides with respect to  $x$ , we obtain

$$\begin{aligned}\Rightarrow \frac{1}{y} \frac{dy}{dx} &= \frac{d}{dx} [\log x \cdot \log(\log x)] \\ \Rightarrow \frac{1}{y} \frac{dy}{dx} &= \log(\log x) \cdot \frac{d}{dx}(\log x) + \log x \cdot \frac{d}{dx} [\log(\log x)] \\ \Rightarrow \frac{dy}{dx} &= y \left[ \log(\log x) \cdot \frac{1}{x} + \log x \cdot \frac{1}{\log x} \cdot \frac{d}{dx}(\log x) \right] \\ \Rightarrow \frac{dy}{dx} &= y \left[ \frac{1}{x} \cdot \log(\log x) + \frac{1}{x} \right] \\ \Rightarrow \frac{dy}{dx} &= (\log x)^{\log x} \left[ \frac{1}{x} + \frac{\log(\log x)}{x} \right]\end{aligned}$$

### Question 8:

Differentiate with respect to  $x$  the function  $\cos(a \cos x + b \sin x)$ , for some constant  $a$  and  $b$ .

#### Solution:

Let  $y = \cos(a \cos x + b \sin x)$

Using chain rule, we get

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx} \cos(a \cos x + b \sin x) \\ &= -\sin(a \cos x + b \sin x) \cdot \frac{d}{dx}(a \cos x + b \sin x) \\ &= -\sin(a \cos x + b \sin x) \cdot [a(-\sin x) + b \cos x] \\ &= (a \sin x - b \cos x) \cdot \sin(a \cos x + b \sin x)\end{aligned}$$

### Question 9:

Differentiate with respect to  $x$  the function  $(\sin x - \cos x)^{(\sin x - \cos x)}$ ,  $\frac{\pi}{4} < x < \frac{3\pi}{4}$

#### Solution:

Let  $y = (\sin x - \cos x)^{(\sin x - \cos x)}$

Taking log on both the sides, we obtain

$$\begin{aligned}\log y &= \log \left[ (\sin x - \cos x)^{(\sin x - \cos x)} \right] \\ &= (\sin x - \cos x) \log (\sin x - \cos x)\end{aligned}$$

Differentiating both sides with respect to  $x$ , we obtain

$$\begin{aligned}\frac{1}{y} \frac{dy}{dx} &= \frac{d}{dx} \left[ (\sin x - \cos x) \log (\sin x - \cos x) \right] \\ \Rightarrow \frac{1}{y} \frac{dy}{dx} &= \log (\sin x - \cos x) \cdot \frac{d}{dx} (\sin x - \cos x) + (\sin x - \cos x) \cdot \frac{d}{dx} \log (\sin x - \cos x) \\ \Rightarrow \frac{1}{y} \frac{dy}{dx} &= \log (\sin x - \cos x) \cdot (\cos x + \sin x) + (\sin x - \cos x) \cdot \frac{1}{(\sin x - \cos x)} \cdot \frac{d}{dx} (\sin x - \cos x) \\ \Rightarrow \frac{dy}{dx} &= (\sin x - \cos x)^{(\sin x - \cos x)} \left[ (\cos x + \sin x) \cdot \log (\sin x - \cos x) + (\cos x + \sin x) \right] \\ \Rightarrow \frac{dy}{dx} &= (\sin x - \cos x)^{(\sin x - \cos x)} (\cos x + \sin x) [1 + \log (\sin x - \cos x)]\end{aligned}$$

#### Question 10:

Differentiate with respect to  $x$  the function  $x^x + x^a + a^x + a^a$ , for some fixed  $a > 0$  and  $x > 0$ .

#### Solution:

$$\text{Let } y = x^x + x^a + a^x + a^a$$

$$\text{Also, let } x^x = u, x^a = v, a^x = w \text{ and } a^a = s$$

Therefore,

$$\begin{aligned}\Rightarrow y &= u + v + w + s \\ \Rightarrow \frac{dy}{dx} &= \frac{du}{dx} + \frac{dv}{dx} + \frac{dw}{dx} + \frac{ds}{dx} \quad \dots(1)\end{aligned}$$

$$\text{Now, } u = x^x$$

Taking logarithm on both the sides, we obtain

$$\begin{aligned}\Rightarrow \log u &= \log x^x \\ \Rightarrow \log u &= x \log x\end{aligned}$$

Differentiating both sides with respect to  $x$ , we obtain

$$\begin{aligned}\frac{1}{u} \frac{du}{dx} &= \log x \cdot \frac{d}{dx}(x) + x \cdot \frac{d}{dx}(\log x) \\ \frac{du}{dx} &= u \left[ \log x \cdot 1 + x \cdot \frac{1}{x} \right] \\ &= x^x [\log x + 1] = x^x (1 + \log x) \quad \dots(2)\end{aligned}$$

Now,  $v = x^a$

Hence,

$$\begin{aligned}\frac{dv}{dx} &= \frac{d}{dx}(x^a) \\ &= ax^{a-1} \quad \dots(3)\end{aligned}$$

Now,  $w = a^x$

Taking logarithm on both the sides, we obtain

$$\begin{aligned}\Rightarrow \log w &= \log a^x \\ \Rightarrow \log w &= x \log a\end{aligned}$$

Differentiating both sides with respect to  $x$ , we obtain

$$\begin{aligned}\frac{1}{w} \frac{dw}{dx} &= \log a \cdot \frac{d}{dx}(x) \\ \frac{dw}{dx} &= w \log a \\ &= a^x \log a \quad \dots(4)\end{aligned}$$

Now,  $s = a^a$

Since  $a$  is constant,  $a^a$  is also a constant.

Hence,

$$\frac{ds}{dx} = 0 \quad \dots(5)$$

From (1), (2), (3), (4) and (5), we obtain

$$\begin{aligned}\frac{dy}{dx} &= x^x (1 + \log x) + ax^{a-1} + a^x \log a + 0 \\ &= x^x (1 + \log x) + ax^{a-1} + a^x \log a\end{aligned}$$

### Question 11:

Differentiate with respect to  $x$  the function  $x^{x^2-3} + (x-3)^{x^2}$ , for  $x > 3$ .

### Solution:

Let  $y = x^{x^2-3} + (x-3)^{x^2}$

Also, let  $u = x^{x^2-3}$  and  $v = (x-3)^{x^2}$

Therefore,

$$y = u + v$$

$$\Rightarrow \frac{dy}{dx} = \frac{du}{dx} + \frac{dv}{dx} \quad \dots(1)$$

Now,  $u = x^{x^2-3}$

Taking logarithm on both the sides, we obtain

$$\begin{aligned} \log u &= \log(x^{x^2-3}) \\ &= (x^2-3)\log x \end{aligned}$$

Differentiating both sides with respect to  $x$ , we obtain

$$\frac{1}{u} \frac{du}{dx} = \log x \cdot \frac{d}{dx}(x^2-3) + (x^2-3) \cdot \frac{d}{dx}(\log x)$$

$$\Rightarrow \frac{1}{u} \frac{du}{dx} = \log x \cdot 2x + (x^2-3) \cdot \frac{1}{x}$$

$$\Rightarrow \frac{du}{dx} = x^{x^2-3} \left[ \frac{x^2-3}{x} + 2x \log x \right] \quad \dots(2)$$

Now,  $v = (x-3)^{x^2}$

Taking logarithm on both the sides, we obtain

$$\begin{aligned} \log v &= \log(x-3)^{x^2} \\ &= x^2 \log(x-3) \end{aligned}$$

Differentiating both sides with respect to  $x$ , we obtain

$$\frac{1}{v} \frac{dv}{dx} = \log(x-3) \cdot \frac{d}{dx}(x^2) + (x^2) \cdot \frac{d}{dx}[\log(x-3)]$$

$$\Rightarrow \frac{1}{v} \frac{dv}{dx} = \log(x-3) \cdot 2x + x^2 \cdot \frac{1}{x-3} \cdot \frac{d}{dx}(x-3)$$

$$\Rightarrow \frac{dv}{dx} = v \left[ 2x \log(x-3) + \frac{x^2}{x-3} \cdot 1 \right]$$

$$\Rightarrow \frac{dv}{dx} = (x-3)^{x^2} \left[ \frac{x^2}{x-3} + 2x \log(x-3) \right] \quad \dots(3)$$

From (1), (2), and (3), we obtain

$$\frac{dy}{dx} = x^{x^2-3} \left[ \frac{x^2-3}{x} + 2x \log x \right] + (x-3)^{x^2} \left[ \frac{x^2}{x-3} + 2x \log(x-3) \right]$$

### Question 12:

Find  $\frac{dy}{dx}$ , if  $y = 12(1 - \cos t)$ ,  $x = 10(t - \sin t)$ ,  $-\frac{\pi}{2} < t < \frac{\pi}{2}$

### Solution:

The given function is  $y = 12(1 - \cos t)$ ,  $x = 10(t - \sin t)$

Hence,

$$\frac{dx}{dt} = \frac{d}{dt} [10(t - \sin t)]$$

$$= 10 \cdot \frac{d}{dt} (t - \sin t)$$

$$= 10(1 - \cos t)$$

$$\frac{dy}{dt} = \frac{d}{dt} [12(1 - \cos t)]$$

$$= 12 \cdot \frac{d}{dt} (1 - \cos t)$$

$$= 12 \cdot [0 - (-\sin t)]$$

$$= 12 \sin t$$

Therefore,

$$\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{12 \sin t}{10(1 - \cos t)}$$

$$= \frac{12 \cdot 2 \sin \frac{t}{2} \cdot \cos \frac{t}{2}}{10 \cdot 2 \sin^2 \frac{t}{2}}$$

$$= \frac{6}{5} \cot \frac{t}{2}$$

### Question 13:

Find  $\frac{dy}{dx}$ , if  $y = \sin^{-1} x + \sin^{-1} \sqrt{1-x^2}$ ,  $-1 \leq x \leq 1$ .

### Solution:

The given function is  $y = \sin^{-1} x + \sin^{-1} \sqrt{1-x^2}$

Hence,

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx} \left[ \sin^{-1} x + \sin^{-1} \sqrt{1-x^2} \right] \\ &= \frac{d}{dx} (\sin^{-1} x) + \frac{d}{dx} (\sin^{-1} \sqrt{1-x^2}) \\ \frac{dy}{dx} &= \frac{1}{\sqrt{1-x^2}} + \frac{1}{\sqrt{1-(\sqrt{1-x^2})^2}} \cdot \frac{d}{dx} (\sqrt{1-x^2}) \\ &= \frac{1}{\sqrt{1-x^2}} + \frac{1}{x} \cdot \frac{1}{2\sqrt{1-x^2}} \cdot \frac{d}{dx} (1-x^2) \\ &= \frac{1}{\sqrt{1-x^2}} + \frac{1}{2x\sqrt{1-x^2}} (-2x) \\ &= \frac{1}{\sqrt{1-x^2}} - \frac{1}{\sqrt{1-x^2}} \\ &= 0\end{aligned}$$

#### Question 14:

If  $x\sqrt{1+y} + y\sqrt{1+x} = 0$  for  $-1 < x < 1$ , prove that  $\frac{dy}{dx} = -\frac{1}{(1+x)^2}$ .

**Solution:**

The given function is  $x\sqrt{1+y} + y\sqrt{1+x} = 0$   
 $\Rightarrow x\sqrt{1+y} = -y\sqrt{1+x}$

Squaring both sides, we obtain

$$\begin{aligned}x^2(1+y) &= y^2(1+x) \\ \Rightarrow x^2 + x^2y &= y^2 + xy^2 \\ \Rightarrow x^2 - y^2 &= xy^2 - x^2y \\ \Rightarrow x^2 - y^2 &= xy(y-x) \\ \Rightarrow (x+y)(x-y) &= xy(y-x) \\ \Rightarrow x+y &= -xy \\ \Rightarrow (1+x)y &= -x \\ \Rightarrow y &= \frac{-x}{(1+x)}\end{aligned}$$

Differentiating both sides with respect to  $x$ , we obtain

$$\begin{aligned}\frac{dy}{dx} &= - \left[ \frac{(1+x) \frac{d}{dx}(x) - (x) \cdot \frac{d}{dx}(1+x)}{(1+x)^2} \right] \\ &= - \frac{(1+x) - x}{(1+x)^2} \\ &= - \frac{1}{(1+x)^2}\end{aligned}$$

Hence proved.

### Question 15:

$$\frac{\left[1 + \left(\frac{dy}{dx}\right)^2\right]^{\frac{3}{2}}}{\frac{d^2y}{dx^2}}$$

If  $(x-a)^2 + (y-b)^2 = c^2$  for  $c > 0$ , prove that  $\frac{\left[1 + \left(\frac{dy}{dx}\right)^2\right]^{\frac{3}{2}}}{\frac{d^2y}{dx^2}}$  is a constant independent of  $a$  and  $b$ .

### Solution:

The given function is  $(x-a)^2 + (y-b)^2 = c^2$

Differentiating both sides with respect to  $x$ , we obtain

$$\begin{aligned}\frac{d}{dx}[(x-a)^2] + \frac{d}{dx}[(y-b)^2] &= \frac{d}{dx}(c^2) \\ \Rightarrow 2(x-a) \cdot \frac{d}{dx}(x-a) + 2(y-b) \cdot \frac{d}{dx}(y-b) &= 0 \\ \Rightarrow 2(x-a) \cdot 1 + 2(y-b) \cdot \frac{dy}{dx} &= 0 \\ \Rightarrow \frac{dy}{dx} &= \frac{-(x-a)}{y-b} \quad \dots(1)\end{aligned}$$

Therefore,

$$\begin{aligned}
 \frac{d^2y}{dx^2} &= \frac{d}{dx} \left[ \frac{-(x-a)}{y-b} \right] \\
 &= - \left[ \frac{(y-b) \cdot \frac{d}{dx}(x-a) - (x-a) \cdot \frac{d}{dx}(y-b)}{(y-b)^2} \right] \\
 &= - \left[ \frac{(y-b) - (x-a) \cdot \frac{dy}{dx}}{(y-b)^2} \right] \\
 &= - \left[ \frac{(y-b) - (x-a) \cdot \left\{ \frac{-(x-a)}{y-b} \right\}}{(y-b)^2} \right] \quad [\text{Using (1)}] \\
 &= - \left[ \frac{(y-b)^2 + (x-a)^2}{(y-b)^3} \right]
 \end{aligned}$$

Hence,

$$\begin{aligned}
 \frac{\left[ 1 + \left( \frac{dy}{dx} \right)^2 \right]^{\frac{3}{2}}}{\frac{d^2y}{dx^2}} &= \frac{\left[ 1 + \frac{(x-a)^2}{(y-b)^2} \right]^{\frac{3}{2}}}{- \left[ \frac{(y-b)^2 + (x-a)^2}{(y-b)^3} \right]} \\
 &= \frac{\left[ \frac{(y-b)^2 + (x-a)^2}{(y-b)^2} \right]^{\frac{3}{2}}}{- \left[ \frac{(y-b)^2 + (x-a)^2}{(y-b)^3} \right]} \\
 &= \frac{\left[ \frac{c^2}{(y-b)^2} \right]^{\frac{3}{2}}}{- \frac{c^2}{(y-b)^3}} \\
 &= \frac{(y-b)^3}{c^2} \\
 &= -c
 \end{aligned}$$



$-c$  is a constant and is independent of  $a$  and  $b$ .

Hence proved.

### Question 16:

If  $\cos y = x \cos(a + y)$  with  $\cos a \neq \pm 1$ , prove that  $\frac{dy}{dx} = \frac{\cos^2(a + y)}{\sin a}$ .

#### Solution:

The given function is  $\cos y = x \cos(a + y)$

Therefore,

$$\begin{aligned} \Rightarrow \frac{d}{dx}[\cos y] &= \frac{d}{dx}[x \cos(a + y)] \\ \Rightarrow -\sin y \frac{dy}{dx} &= \cos(a + y) \cdot \frac{d}{dx}(x) + x \cdot \frac{d}{dx}[\cos(a + y)] \\ \Rightarrow -\sin y \frac{dy}{dx} &= \cos(a + y) + x \cdot [-\sin(a + y)] \frac{dy}{dx} \\ \Rightarrow [x \sin(a + y) - \sin y] \frac{dy}{dx} &= \cos(a + y) \quad \dots (1) \end{aligned}$$

Since,  $\cos y = x \cos(a + y) \Rightarrow x = \frac{\cos y}{\cos(a + y)}$

Then, equation (1) becomes,

$$\begin{aligned} \left[ \frac{\cos y}{\cos(a + y)} \cdot \sin(a + y) - \sin y \right] \frac{dy}{dx} &= \cos(a + y) \\ \Rightarrow [\cos y \cdot \sin(a + y) - \sin y \cdot \cos(a + y)] \cdot \frac{dy}{dx} &= \cos^2(a + y) \\ \Rightarrow \sin(a + y - y) \frac{dy}{dx} &= \cos^2(a + y) \\ \Rightarrow \frac{dy}{dx} &= \frac{\cos^2(a + y)}{\sin a} \end{aligned}$$

Hence proved.

### Question 17:

If  $x = a(\cos t + t \sin t)$  and  $y = a(\sin t - t \cos t)$ , find  $\frac{d^2 y}{dx^2}$ .

### Solution:

The given function is  $x = a(\cos t + t \sin t)$  and  $y = a(\sin t - t \cos t)$   
Therefore,

$$\begin{aligned}\frac{dx}{dt} &= a \cdot \frac{d}{dt}(\cos t + t \sin t) \\ &= a \left[ -\sin t + \sin t + t \cdot \frac{d}{dt}(t) + t \cdot \frac{d}{dt}(\sin t) \right] \\ &= a [-\sin t + \sin t + t \cos t] \\ &= at \cos t\end{aligned}$$

$$\begin{aligned}\frac{dy}{dt} &= a \cdot \frac{d}{dt}(\sin t - t \cos t) \\ &= a \left[ \cos t - \left\{ \cos t \cdot \frac{d}{dt}(t) + t \cdot \frac{d}{dt}(\cos t) \right\} \right] \\ &= a [\cos t - \{\cos t - t \sin t\}] \\ &= at \sin t\end{aligned}$$

$$\frac{dy}{dx} = \frac{\left(\frac{dy}{dt}\right)}{\left(\frac{dx}{dt}\right)} = \frac{at \sin t}{at \cos t} = \tan t$$

$$\begin{aligned}\frac{d^2y}{dx^2} &= \frac{d}{dx} \left( \frac{dy}{dx} \right) = \frac{d}{dx}(\tan t) = \sec^2 t \cdot \frac{dt}{dx} \\ &= \sec^2 t \cdot \frac{1}{at \cos t} \quad \left[ \frac{dx}{dt} = at \cos t \Rightarrow \frac{dt}{dx} = \frac{1}{at \cos t} \right] \\ &= \frac{\sec^3 t}{at}, 0 < t < \frac{\pi}{2}\end{aligned}$$

### Question 18:

If  $f(x) = |x|^3$ , show that  $f''(x)$  exists for all real  $x$ , and find it.

### Solution:

It is known that  $|x| = \begin{cases} x, & \text{if } x \geq 0 \\ -x, & \text{if } x < 0 \end{cases}$

Therefore, when  $x \geq 0$ ,  $f(x) = |x|^3 = x^3$

In this case,  $f'(x) = 3x^2$  and hence,  $f''(x) = 6x$

When  $x < 0$ ,  $f(x) = |x|^3 = (-x)^3 = -x^3$

In this case,  $f'(x) = -3x^2$  and hence,  $f''(x) = -6x$

Thus, for  $f(x) = |x|^3$ ,  $f''(x)$  exists for all real  $x$  and is given by,

$$f''(x) = \begin{cases} 6x, & \text{if } x \geq 0 \\ -6x, & \text{if } x < 0 \end{cases}$$

### Question 19:

Using mathematical induction prove that  $\frac{d}{dx}(x^n) = nx^{n-1}$  for all positive integers  $n$ .

#### Solution:

To prove:  $P(n): \frac{d}{dx}(x^n) = nx^{n-1}$  for all positive integers  $n$ .

For  $n = 1$ ,

$$P(1): \frac{d}{dx}(x) = 1 = 1 \cdot x^{1-1}$$

Therefore,  $P(n)$  is true for  $n = 1$ .

Let  $P(k)$  is true for some positive integer  $k$ .

$$\text{That is, } P(k): \frac{d}{dx}(x^k) = kx^{k-1}$$

It has to be proved that  $P(k+1)$  is also true.

Consider

$$\begin{aligned} \frac{d}{dx}(x^{k+1}) &= \frac{d}{dx}(x \cdot x^k) \\ &= x^k \cdot \frac{d}{dx}(x) + x \cdot \frac{d}{dx}(x^k) \quad [\text{By applying product rule}] \\ &= x^k \cdot 1 + x \cdot k \cdot x^{k-1} \end{aligned}$$

$$\begin{aligned} \frac{d}{dx}(x^{k+1}) &= x^k + kx^k \\ &= (k+1) \cdot x^k \\ &= (k+1) \cdot x^{(k+1)-1} \end{aligned}$$

Thus,  $P(k+1)$  is true whenever  $P(k)$  is true.

Therefore, by the principle of mathematical induction, the statement  $P(n)$  is true for every positive integer  $n$ .

Hence, proved.

### Question 20:

Using the fact that  $\sin(A+B) = \sin A \cos B + \cos A \sin B$  and the differentiation, obtain the sum formula for cosines.

#### Solution:

Given,  $\sin(A+B) = \sin A \cos B + \cos A \sin B$

Differentiating both sides with respect to  $x$ , we obtain


$$\begin{aligned} \frac{d}{dx} [\sin(A+B)] &= \frac{d}{dx} (\sin A \cos B) + \frac{d}{dx} (\cos A \sin B) \\ \Rightarrow \cos(A+B) \cdot \frac{d}{dx} (A+B) &= \cos B \cdot \frac{d}{dx} (\sin A) + \sin A \cdot \frac{d}{dx} (\cos B) + \sin B \cdot \frac{d}{dx} (\cos A) + \cos A \cdot \frac{d}{dx} (\sin B) \\ \Rightarrow \cos(A+B) \cdot \frac{d}{dx} (A+B) &= \cos B \cdot \cos A \frac{dA}{dx} + \sin A (-\sin B) \frac{dB}{dx} + \sin B (-\sin A) \frac{dA}{dx} + \cos A \cos B \frac{dB}{dx} \\ \Rightarrow \cos(A+B) \cdot \left[ \frac{dA}{dx} + \frac{dB}{dx} \right] &= (\cos A \cos B - \sin A \sin B) \cdot \left[ \frac{dA}{dx} + \frac{dB}{dx} \right] \\ \Rightarrow \cos(A+B) &= \cos A \cos B - \sin A \sin B \end{aligned}$$

### Question 21:

Does there exist a function which is continuous everywhere but not differentiable at exactly two points? Justify your answer?

#### Solution:

Consider,  $y = \begin{cases} |x| & -\infty < x \leq 1 \\ 2-x & 1 \leq x \leq \infty \end{cases}$



It can be seen from the above graph that the given function is continuous everywhere but not differentiable at exactly two points which are 0 and 1.

### Question 22:

If  $y = \begin{vmatrix} f(x) & g(x) & h(x) \\ l & m & n \\ a & b & c \end{vmatrix}$ , prove that  $\frac{dy}{dx} = \begin{vmatrix} f'(x) & g'(x) & h'(x) \\ l & m & n \\ a & b & c \end{vmatrix}$

### Solution:

Given,  $y = \begin{vmatrix} f(x) & g(x) & h(x) \\ l & m & n \\ a & b & c \end{vmatrix}$

$$\Rightarrow y = (mc - nb)f(x) - (lc - na)g(x) + (lb - ma)h(x)$$

Then,

$$\begin{aligned} \frac{dy}{dx} &= \frac{d}{dx}[(mc - nb)f(x)] - \frac{d}{dx}[(lc - na)g(x)] + \frac{d}{dx}[(lb - ma)h(x)] \\ &= (mc - nb)f'(x) - (lc - na)g'(x) + (lb - ma)h'(x) \\ &= \begin{vmatrix} f'(x) & g'(x) & h'(x) \\ l & m & n \\ a & b & c \end{vmatrix} \end{aligned}$$

Thus,  $\frac{dy}{dx} = \begin{vmatrix} f'(x) & g'(x) & h'(x) \\ l & m & n \\ a & b & c \end{vmatrix}$  proved.

### Question 23:

If  $y = e^{a \cos^{-1} x}$ ,  $-1 \leq x \leq 1$ , show that  $(1-x^2) \frac{d^2 y}{dx^2} - x \frac{dy}{dx} - a^2 y = 0$

### Solution:

The given function is  $y = e^{a \cos^{-1} x}$

Taking logarithm on both the sides, we obtain

$$\Rightarrow \log y = a \cos^{-1} x \log e$$

$$\Rightarrow \log y = a \cos^{-1} x$$

Differentiating both sides with respect to  $x$ , we obtain

$$\Rightarrow \frac{1}{y} \frac{dy}{dx} = a \cdot \frac{-1}{\sqrt{1-x^2}}$$

$$\Rightarrow \frac{dy}{dx} = \frac{-ay}{\sqrt{1-x^2}}$$

By squaring both the sides, we obtain

$$\Rightarrow \left( \frac{dy}{dx} \right)^2 = \frac{a^2 y^2}{1-x^2}$$

$$\Rightarrow (1-x^2) \left( \frac{dy}{dx} \right)^2 = a^2 y^2$$

Again, differentiating both sides with respect to  $x$ , we obtain

$$\Rightarrow \left( \frac{dy}{dx} \right)^2 \frac{d}{dx} (1-x^2) + (1-x^2) \times \frac{d}{dx} \left[ \left( \frac{dy}{dx} \right)^2 \right] = a^2 \frac{d}{dx} (y^2)$$

$$\Rightarrow \left( \frac{dy}{dx} \right)^2 (-2x) + (1-x^2) \times 2 \frac{dy}{dx} \cdot \frac{d^2 y}{dx^2} = a^2 \cdot 2y \cdot \frac{dy}{dx}$$

$$\Rightarrow -x \frac{dy}{dx} + (1-x^2) \frac{d^2 y}{dx^2} = a^2 \cdot y \quad \left[ \frac{dy}{dx} \neq 0 \right]$$

$$\Rightarrow (1-x^2) \frac{d^2 y}{dx^2} - x \frac{dy}{dx} - a^2 y = 0$$

Hence proved.