

COLLAGEN, A KEY COMPONENT FOR HEALTHY AND YOUTHFUL SKIN

COLLAGEN IS ONE OF THE MOST ABUNDANT PROTEINS IN YOUR BODY, MAKING UP AROUND 75% OF YOUR SKIN³

Among the various collagen types, type 1 and type 3 are the most prevalent in the skin:⁴

- Collagen type 1 is the main structural component of the skin⁵
- Collagen type 3 plays a role in the processes of healing and repair⁵

WITH AGE, YOUR COLLAGEN LEVELS RAPIDLY DEPLETE^{6,7}

Throughout adult life, about 1% of collagen is lost per year⁷

BY AGE 35



Up to **15**% of collagen loss⁷

35 BY AGE **45**



Up to **25**% of collagen loss⁷

BY AGE **55**




Up to **35**% of collagen loss⁷

COLLAGEN HELPS IN MAINTAINING
YOUR SKIN'S SHAPE AND FIRMNESS^{4,6}

SCULPTRA INCREASES LEVELS
OF COLLAGEN

Sculptra is an injectable collagen biostimulator made from poly-L-lactic acid (PLLA)*, a fully absorbed polymer which works deep into the skin to improve its inner structure over a series of treatments.^{1,8-10}


PLLA MICROPARTICLES WORK DEEP WITHIN THE SKIN TO STIMULATE THE REPLACEMENT OF LOST COLLAGEN^{1,8-10}



PLLA microparticles are injected subcutaneously or into the deep dermis



PLLA microparticles stimulate the replacement of lost collagen



Over time, collagen production rebuilds the skin's structure

BY STIMULATING THE PRODUCTION OF COLLAGEN TYPE 1, SCULPTRA CAN GRADUALLY RESTORE SKIN FIRMNESS, QUALITY AND VOLUME^{1,8-10}

The most common side effects after initial treatment include injection site swelling, tenderness, redness, pain, bruising, bleeding, itching and lumps.⁸

*PLLA has been used in medical products for more than 30 years and has been tested for safety.11

SCULPTRA OFFERS YOU PROVEN EFFICACY AND HIGH PATIENT SATISFACTION FOR MORE THAN 2 YEARS

INCREASE IN SKIN ELASTICITY AND IMPROVED OVERALL SKIN QUALITY AND RADIANCE*12





Skin elasticity



Skin radiance



Skin smoothness

ASSESSMENT OF WRINKLES

PATIENTS SHOWING LONG-LASTING IMPROVEMENT *13



1.5 years after last treatment



PATIENT SATISFACTION

PATIENTS SATISFIED WITH THEIR TREATMENT RESULTS 2 YEARS AFTER THEIR LAST TREATMENT¹⁴



SCULPTRA HELPS YOU OBTAIN A MORE YOUTHFUL-LOOKING SKIN



Sculptra has made a massive difference to me. It's given me confidence and taken years off my face.

CARA. 42

Refore

8 months after treatment

Actual patient. Individual results may vary. Treated with 3 vials of Sculptra, 3 sessions.

I am absolutely delighted with the increased volume and improved skin tone and feel I look much younger and healthier!

STEWART, 54



Before

7 months after treatment

Actual patient. Individual results may vary.

Treated with 5 vials of Sculptra 3 sessions.

© Dr Linda Eve. Pictures are reproduced with permission.

^{*}Global skin quality improvement included radiance, smoothness, and pigment uniformity as well as a decrease of erythema index and pore size.

†Randomised, controlled, double-blind, multicentre study. Patients received 3 treatments every 4 weeks with either PLLA (treatment group) or saline (control group) injections. †Efficacy of Sculptra for the correction of nasolabial fold wrinkles based on Investigator's Global Evaluations.

TEST THE FIRMNESS OF YOUR SKIN

AS YOU AGE, YOUR SKIN WILL DEVELOP WRINKLES, SAG AND LOSE ITS ABILITY TO SNAP BACK AFTER STRETCHING

One simple way to assess if your skin has lost its elasticity and firmness is by pinching your skin.

- Does your skin feel looser?
- Does it take longer for the skin to bounce back?

ARE YOU LOOKING FOR A SUBTLE TREATMENT, ABLE TO OFFER GRADUAL AND LONG-LASTING BENEFITS?

WITH SCULPTRA. ACHIEVE MORE YOUTHFUL-LOOKING SKIN











TO LEARN MORE ABOUT SCULPTRA TALK TO YOUR AESTHETIC PRACTITIONER

*Actual patient. Individual results may vary. Treated with 5 vials of Sculptra 3 sessions.

SCULPTRA®

IS KEY TO FIRMER AND REJUVENATED SKIN

Stimulates the natural replacement of lost collagen^{1,8-10}

Improves skin firmness and elasticity 10,12

Proven efficacy for more than 2 years 13,15

Increase in skin elasticity and improved overall skin quality and radiance¹²

High patient satisfaction for more than 2 years¹⁴

An established safety profile with over 50 publications to date, in more than 10,000 patients¹³⁻¹⁶



REFERENCES:

- 1. Goldberg D, et al. Dermatol Surg. 2013;39(6):915-922.
- 2. Hexsel D, et al. Dermatol Surg. 2020;46(8):1122-1124.
- 3. Shoulders MD, Raines RT. Annu Rev Biochem. 2009;78:929-958.
- 4. Aziz J, et al. Skin Pharmacol Physiol. 2016;29(4):190-203.
- 5. Wu J, et al. J Biol Chem. 2010;285(24):18537-18544.
- 6. Farage MA, et al. Adv Wound Care (New Rochelle). 2013;2(1):5-10.
- 7. Shuster S, et al. Br J Dermatol. 1975;93:639-643.
- 8. Sculptra. EU IFU.
- 9. Lacombe V. Facial Plast Surg. 2009:25(2):95-99.
- 10. Mest DR, Humble G. Dermatol Surg. 2006;32(11):1336-1345.
- 11. Lowe NJ. J Eur Acad Dermatol Venereol. 2006:20(1):2-6.
- 12. Bohnert K, et al. Plast Reconstr Surg. 2019;45(5):718-724.
- 13. Brandt FS, et al. Aesthet Surg J. 2011;31(5):521-528.
- 14. Brown SA, et al. Plast Reconstr Surg. 2011;127(4):1684-1692.
- 15. Narins RS, et al. J Am Acad Dermatol. 2010;62(3):448-462.
- 16. Data on File (MA-47133).

ABBREVIATIONS:

PLLA = Poly-L-lactic acid.

Adverse events should be reported. For the UK, Reporting forms and information can be found at www.mhra.gov.uk/yellowcard or search for Yellow Card in the Google Play or Apple App Store.

For Ireland, Suspected adverse events can be reported via HPRA Pharmacovigilance, Website: www.hpra.ie; Adverse events should also be reported to Galderma (UK) Ltd, Email: medinfo.uk@galderma.com

Tel: **+44 (0) 300 3035674**