



Case:
History Central vein Occlusion

Multiple intravitreal injections


Dense lens changes


Potential vision 20/200 ?





































Figure 1. Final visual acuity after suprachoroidal hemorrhage associated with pars plana vitrectomy. Each point represents an individual. The x- and y-axes are visual acuity (LogMAR).

Surgical Management of Suprachoroidal Hemorrhage

Although uncommon, blood in the suprachoroidal space can warrant surgery.

By AMAN CHANDRA, MRCSEd, FRCOphth, PhD • TOM WILLIAMSON, MD, FRCS, FRCOphth


Retinal Physician, July 1, 2014



[Ophthalmology](#). 2016 Jun;123(6):1252-6. doi: 10.1016/j.ophtha.2016.02.014. Epub 2016 Mar 18.

Previous Intravitreal Therapy Is Associated with Increased Risk of Posterior Capsule Rupture during Cataract Surgery.

Lee AY1, Day AC2, Egan C3, Bailey C4, Johnston RL5, Tsaloumas MD6, Denniston AK7, Tufail A8; United Kingdom Age-related Macular Degeneration and Diabetic Retinopathy Electronic Medical Records Users Group.



[Ophthalmology](#). 2016 Jun;123(6):1252-6. doi: 10.1016/j.ophtha.2016.02.014. Epub 2016 Mar 18.

Previous Intravitreal Therapy Is Associated with Increased Risk of Posterior Capsule Rupture during Cataract Surgery.

Lee AY¹, Day AC², Egan C³, Bailey C⁴, Johnston RL⁵, Tsaloumas MD⁶, Denniston AK⁷, Tufail A⁸; United Kingdom Age-related Macular Degeneration and Diabetic Retinopathy Electronic Medical Records Users Group.

Nebraska Laser Eye Associates

www.nebraskaeye.com