Office Use Only					
APPL					
RAD					
CK					



Orthopedic Foundation for Animals
2300 E Nifong Blvd, Columbia, MO 65201
Phone (573) 442-0418 | Fax (573) 875-5073
Email ofa@offa.org | www.ofa.org
A Not-for-Profit Organization

Application for Basic Cardiac Database

Registered name:			AKC registration number: Other registry name:			ry name:	
KMA-RILEY ROO		T-			Other regist	ry #:	
Breed: GREAT DANE		Sex:	Date of birth (MM/DD/YY): 11/12/2018				
Microchip/tattoo:			Registration number of sire		Registration number of dam: WS50410809		
985112010515751			WP8314630		VVS5		
Owner name: Co-Owner name: DR. WENDY WHITLOCK MR. MAR		JACKSON	PETCARDIA			Date of evaluation (MM/DD/YY): 09/02/2021	
Mailing address:	IVIIX. IVIAIXI	CJACKSON	Mailing address:	~		09/02/2021	
maining address.			774 MALET	A LN			
City:	State:	Zip/postal code:	City:		State:	Zip/postal code:	
COLORADO SPRINGS	CO	80923	CASTLE RO	OCK	CO	80103	
Phone:	E-mail:		Phone:		E-mail:		
			(720) 724-8	013	INFO	@PETCARDIA.COM	
Veterinary Exam Results Clinical findings based on cardiac aus	cultation is require	ed. (see page 2)					
		AUSCULTATI	ON (REQUIRED)				
	Normal	Abnormal		ımia 🗆			
Murmur G	rade: I 🗆			V□	VI 🗆		
PMI:	Left □	Right	☐ Base ☐ /	Арех 🗆			
Timing:	Systolic	Diastolic	☐ Contin	uous 🗆			
Extra Soun	ds: Click 🗌	Gallop	☐ Split S1☐ Sp	olit S2 🗆			
•	ar examination—hea ular examination—h	rt disease is not evide leart disease cannot	ent be diagnosed nor exclu ase; indicate suspected			ding.	
☐ I certify that the standards for cardiad I DID verify microchip/tattoo on this	dog 🗔 I	DID NOT verify micr	ochip/tattoo on this do	og	amination.		
Veterinarian Signature Che	ck one box: 📮 P	ractitioner, 📮 Sp	pecialist, 🖵 Cardio	ologist		Date	
Fees Animals Over 12 Months Litter of 3 or more submit	ted together	\$30.00	Minimum of 5 in	dividuals		/co-owned by same person. \$7.50 dation for Animals.	
Card number	c	ardholder name		Exp date M	IM/YY	CVV	

Methods of Examination

Clinical Examination

- 1. The clinical cardiac examination should be conducted in a systematic manner. The arterial and venous pulses, mucous membranes, and precordium should be evaluated. Heart rate should be obtained. The clinical examination should be performed by an individual with advanced training in cardiac diagnosis. Board certication by the American College of Veterinary Internal Medicine, Specialty of Cardiology is considered by the American Veterinary Medical Association as the benchmark of clinical prociency for veterinarians in clinical cardiology, and examination by a Diplomate of this specialty board is recommended. However, any licensed veterinarian may be able to perform this examination by auscultation.
- 2. Cardiac auscultation should be performed in a quiet, distraction-free environment. The animal should be standing and restrained, but sedative drugs should be avoided. Panting must be controlled, and if necessary, the dog should be given time to rest and acclimate to the environment. The clinician should be able to identify the cardiac valve areas for auscultation. The examiner should gradually move the stethoscope across all valve areas and also should auscultate over the subaortic area, ascending aorta, pulmonary artery, and the left craniodorsal cardiac base. Following examination of the left precordium, the right precordium should be examined.
 - The mitral valve area is located over and immediately dorsal to the palpable left apical impulse and is identified by palpation with the tips of the ingers. The stethoscope is then placed over themitral area and the heart sounds identified.
 - The aortic valve area is dorsal and 1 or 2 intercostal spaces cranial to the left apical impulse. The second heart sound will become most intense when the stethoscope is centered over the aortic valve area. Murmurs originating from or radiating to the subaortic area of auscultation are evident immediately caudoventral to the aortic valve area. Murmurs originating from or radiating into the ascending aorta will be evident craniodorsal to the aortic valve and may also project to the right cranial thorax and to the carotid arteries in the neck.
 - The pulmonic valve area is ventral and one intercostal space cranial to the aortic valve area. Murmurs originating from or radiating into the main pulmonary artery will be evident dorsal to the pulmonic valve over the left hemithorax.
 - The tricuspid valve area is a relatively large area located on the right hemithorax, opposite and slightly cranial to the mitral valve area.
 - The clinician should also auscultate along the ventral right precordium (right sternal border) and over the right craniodorsal cardiac border.
 - Any cardiac murmurs or abnormal sounds should be noted.
 Murmurs should be described as indicated below.

- 3. Description of cardiac murmurs—A full description of the cardiac murmur should be made and recorded in the medical record.
 - Murmurs should be designated as systolic, diastolic, or continuous.
 - The point of maximal murmur intensity should be indicated as described above. When a precordial thrill is palpable, the murmur will generally be most intense over this vibration.
 - Murmurs that are only detected intermittently or are variable should be so indicated.
 - The radiation of the murmur should be indicated.
 - Grading of heart murmurs is as follows:

Grade 1—a very soft murmur only detected after very careful auscultation

Grade 2—a soft murmur that is readily evident

Grade 3—a moderately intense murmur not associated with a palpable precordial thrill (vibration)

Grade 4—a loud murmur; a palpable precordial thrill is not present or is intermittent

Grade 5—a loud cardiac murmur associated with a palpable precordial thrill and not audible when the stethoscope is lifted from the thoracic wall

Grade 6—a loud cardiac murmur associated with a palpable precordial thrill and audible even when the stethoscope is lifted from the thoracic wall

Other descriptive terms may be indicated at the discretion
of the examiner; these include such timing descriptors as:
proto(early)-systolic, ejection or crescendo-decrescendo,
holo-systolic or pan-systolic, decrescendo, and tele(late)systolic and descriptions of subjective characteristics such
as: musical, vibratory, harsh, and machinery.

4. Effects of heart rate, heart rhythm, and exercise.

- Some heart murmurs become evident or louder with changes in autonomic activity, heart rate, or cardiac cycle length. Such changes may be induced by exercise or other stresses. The importance of evaluating heart murmurs after exercise is currently unresolved. It appears that some dogs with congenital subaortic stenosis or with dynamic out ow tract obstruction may have murmurs that only become evident with increased sympathetic activity or after prolonged cardiac lling periods during marked sinus arrhythmia. It also should be noted that some normal, innocent heart murmurs may increase in intensity after exercise. Furthermore, panting artifact may be a problem after exercise.
- It is most likely that examining dogs after exercise will result in increased sensitivity to diagnosis of soft murmurs but probably decreased speci city as well. Auscultation of the heart following exercise is at the discretion of the examining veterinarian.
- At this time the OFA does not require a post exercise examination in the assessment of heart murmurs in dogs; however, this practice may be modi ed should de nitive information become available.