



ProFume[®] Gas Fumigant (Sulfuryl Fluoride) Technical Bulletin



ProFume® Gas Fumigant (Sulfuryl Fluoride) Technical Bulletin

This technical bulletin for ProFume® gas fumigant (sulfuryl fluoride) is provided for reference purposes only and is not a substitute for or an addition to labeling and Safety Data Sheets. *Always read and follow label directions.* The information and any recommendations in this bulletin ("Information") are presented in good faith; however, Douglas Products makes no warranty as to the completeness or accuracy of the information. This information is supplied upon the condition that persons receiving it will make their own determinations as to its suitability for their purposes prior to use and consult with their advisors to ensure compliance with all federal, national/country, state and local regulations. In no event will Douglas Products be responsible for damages of any nature whatsoever resulting from the use of or reliance upon this information.

NO REPRESENTATIONS OR WARRANTIES, EITHER EXPRESSED OR IMPLIED, OF MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE OR OF ANY OTHER NATURE ARE MADE HEREUNDER WITH RESPECT TO THE INFORMATION OR THE PRODUCTS TO WHICH THE INFORMATION REFERS.

Douglas Products reserves all rights to, and in, the contents of this document. This bulletin has been prepared specifically for ProFume and no other fumigant. No part of this document may be copied, in part or in whole, or reproduced for another purpose without the prior written permission of Douglas Products.

Contact your local Douglas Products or distributor representative for additional information on ProFume.

Overview

ProFume® gas fumigant (sulfuryl fluoride) is a broad-spectrum postharvest fumigant developed in collaboration with leading stored-product researchers, fumigators and food industries worldwide. Fumigation is usually the preferred method of postharvest pest eradication because ProFume, with its excellent penetration properties, can control pests anywhere they occur within the commodity and the storage, transportation or food processing structure under treatment. ProFume, because of its unique mode of action, can also be used as a viable rotation tool to manage and help prevent resistance issues with other fumigants and insecticides. Food tolerances have been established for ProFume which enable its use on a wide variety of commodities for global trade and in mills, food processing facilities, warehouses, storage containers, chambers and transportation vehicles.

Noteworthy features

ProFume is:

- A broad-spectrum fumigant effective on all life stages of insects, other arthropods and rodent pests
- Flexible for use in long- or short-exposure fumigations
- Nonflammable, odorless, colorless gas that rapidly vaporizes and distributes quickly
- Noncorrosive gas for use in sensitive areas having equipment and electronic devices; very low reactivity as a gas and does not react with materials to form unpleasant odors or flavors
- Rapidly penetrates porous materials, rapidly aerates from materials and commodities, and has low sorption on fumigated materials
- An inorganic gas that does not leave surface residues on inert surfaces, such as stainless steel, glass and ceramic, after fumigation
- Not an ozone depleter and does not interact or cause local ozone formation
- A new mode of action which can be utilized for resistance management strategies

The Sulfuryl Fluoride Story

History: The Dow Chemical Company developed sulfuryl fluoride as a structural fumigant and marketed it as Vikane® gas fumigant since 1961. Vikane has been successfully used to fumigate more than 3 million structures, including homes, museums, cathedrals, historical landmarks, rare-book libraries, and scientific and medical research laboratories to eradicate termites, wood-boring beetles, cockroaches, rodents, bed bugs and other structure-infesting pests.

Postharvest insect pests that infest grains and dried fruits and tree nuts in mills, warehouses and food storage facilities cause substantial economic and quality losses. Postharvest losses due to insects in the United States alone were estimated at \$5 billion per year (Pimental 1991). With the adoption of the Montreal Protocol, the search for replacements for methyl bromide began in the early 1990s. Beginning in 1995, Dow AgroSciences formed partnerships with leading stored product researchers, fumigators and food industries around the world to evaluate and develop sulfuryl fluoride as a postharvest fumigant.



Pest Control Studies: Researchers at six world-recognized stored-product research laboratories in the United States (USDA-ARS and Dried Fruit and Tree Nut Association [DFA] in Fresno, California) and Europe (Food and Environmental Research Agency in the UK, Julius Kuehn Institute in Germany, the University of Milan in Italy and Laboratoire National des Denrées Stockées in France) defined the dosages required to control all the life stages of target stored-product insect pests under a range of fumigation conditions (Baltaci et al. 2008, 2009; Baraket et al. 2009; Bell 2006; Bell et al. 1999, 2003; Bell and Savvidou 1999; Ducom et al. 2003; Schneider and Hartsell 1999; Zettler et al. 1999; Zettler and Gill 1999). These studies, conducted in laboratory and field trials, confirmed the effectiveness of ProFume® gas fumigant on all life stages, including diapausing stages and eggs, of a wide range of post-harvest insect pests, including the important pest species of Coleoptera and Lepidoptera.

Researchers also evaluated the efficacy of commercial applications of ProFume® gas fumigant through trapping of in-situ insects and/or control of confined bioassay insects placed in flour mills and food processing plants (Drinkall et al. 2003, Hartzler et al. 2010, Reichmuth et al. 2003, Small 2007, Subramanyam 2006, Suss and Savoldelli 2008, Thoms et al. 2008, Tsai et al. 2006) and in commodities such as dates (Williams and Thoms 2008, Williams 2009) and cocoa beans (Bookout and Milyo 2006). Fumigations were prepared and conducted by commercial applicators. Researchers concluded ProFume is an effective fumigant for disinfestation of structures and commodities.

Food Quality: Extensive studies have been conducted to verify fumigation with ProFume does not affect the quality of commodities when compared to the same nonfumigated commodities. These trials were not required for registration but have been important for acceptance of ProFume by the food processing industry.

Grain kernels of three wheat varieties (durum, hard red winter and soft red winter wheat) tested in residue trials were also tested in a battery of quality trials conducted by the Department of Grain Science and Industry, Kansas State University, using state-of-the-art equipment and protocols approved by the American Association of Cereal Chemists (AACC 2000). For all quality trials, grain was fumigated with a dosage of ProFume that exceeded the current maximum label dosage by 33%. Fumigation with ProFume did not affect wheat kernel quality, based on test weight; thousand-kernel weight; and moisture, protein, lipid, dietary fiber and thiamine content. Milling quality of fumigated grain, as determined by falling number, was not affected. Properties of dough and baked goods from flour milled from fumigated and nonfumigated red wheat were equivalent. This determination was based on dough peak mixing time, stability (ability of flour to adjust to overmixing and undermixing), and absorption rate of water measured using a farinograph, dough viscosity measured using a viscoanalyzer, and dough tenacity, extensibility, strength and volume as measured using an alveograph. Baking tests showed bread volume, cookie height, and width and spread of dough were not affected by grain fumigation. Properties of dough and spaghetti of semolina flour milled from fumigated and nonfumigated durum wheat were equivalent. This was based on visual and handling characteristics of raw dough and spaghetti, color of raw and dried spaghetti, cooking time and cooking yield of spaghetti, and tensile strength of cooked spaghetti.

Taste testing trials for high-value commodities (dried fruit, tree nuts and cocoa) were conducted. Single and repeated fumigations were conducted at a dosage of ProFume that exceeded the current maximum label dosage by 33% for dried fruits and tree nuts. Walnuts were also vacuum fumigated. Tests were conducted by experts trained in sensory evaluation using two methods of taste testing. No significant differences in taste quality of fumigated and nonfumigated dried fruit and tree nuts were observed in these trials.

For cocoa sensory trials, cocoa beans from Sulawesi and Ivory Coast were fumigated at three dosages. Fumigated and nonfumigated cocoa beans were roasted, cracked, winnowed and milled into liquor for testing. Taste testing was conducted by the nine member companies of the Chocolate Manufacturers Association (CMA, now part of the National Confectioner's Association), using their in-house experts and protocols. The consensus of this testing is that there are no sensory issues with ProFume, and the CMA members approved ProFume as a suitable fumigant.

Food Residues: Research by Meikle (1964) identified the two types of residue found in food commodities that were fumigated with sulfuryl fluoride: the parent sulfuryl fluoride and the degradation products fluoride anion (F-) and sulfate (SO₄²⁻). Sulfate as a terminal residue of sulfuryl fluoride is not of any toxicological concern due to its natural abundance in living systems. Research by Meikle and Stewart (1962) and Scheffrahn et al. (1989) investigated these residues in a variety of food commodities. Their research demonstrated that sulfuryl fluoride residues are transient in food commodities and rapidly decrease to very low (ppb)

or nondetectable levels. Fluoride and sulfate can form permanent residues in commodities containing proteinaceous material, particularly when fat (oil) is present. In oil alone there were essentially no detectable residues of fluoride or sulfate.


Extensive residue trials fumigating a wide variety of commodities with the maximum label dosage rates of ProFume® gas fumigant have been conducted following Good Laboratory Practice (GLP). Representative types for each commodity were selected for the residue trials. For example, three of six classes of wheat milled in the United States (durum, hard red winter and soft red winter wheat) were tested in the GLP residue trials. These classes represent common wheat varieties milled and have a wide range in protein content (available amino functional groups for reaction) that could affect F anion residue formation. In addition to the whole commodity, any processed fractions that would be fumigated were tested. Therefore, processed fractions of cereal grains, such as flour, germ and meal for wheat, were evaluated in the GLP residue trials.

In addition to laboratory residue trials, GLP residue trials were conducted in flour mills and commercial food processing facilities to determine F anion residues in fumigated commodities following processing. Residue data from all trials were used to develop risk assessments for adults and children based on percentage of commodity treated annually with ProFume, contribution of commodity to diet and other factors.

Residue Tolerances: The United States Environmental Protection Agency (EPA) used the residue trials described above to develop residue tolerances for the parent compound, sulfuryl fluoride and F anion in commodities fumigated with ProFume. These residue tolerances are published in the Federal Register: 40 CFR 180.575 for sulfuryl fluoride and 40 CFR 180.145 for fluorine compounds. Other countries/regions where ProFume gas fumigant is registered for application, including Australia, Canada, Mexico and the European Union, have also used the above residue trials to establish residue tolerances for ProFume.

For international trade, maximum residue tolerances (MRLs) for sulfuryl fluoride are established in CODEX for tree nuts, dried fruits, and cereal grains and their processed fractions (Codex Alimentarius/FAO, see: <http://www.fao.org/fao-who-codexalimentarius/standards/pestres/en/>). Not all countries use CODEX, such as Japan and the EU, and instead have their own listing of MRLs for sulfuryl fluoride. Since ProFume was first registered in the United States in 2004, a wide variety of commodities fumigated with ProFume have been traded globally. A global MRL database is available through Bryant Christie Inc. (www.globalmrl.com). Global MRL databases may be available through other service providers and regulatory authorities.

Commodities: In many countries where ProFume is registered, it is labeled for fumigation of a variety of raw agricultural commodities. Depending upon the country, these commodities may include cereal grains (whole kernels and processed fractions) such as wheat (soft red, hard red, durum and white), rice, corn (field and popcorn), barley and oats; dried fruits such as raisins, prunes, figs, apples, apricots and dates; tree nuts such as walnuts, almonds, hazelnuts and pecans; and dried herbs, ginger, cocoa and coffee beans, cottonseed, dried legumes, ham, cheese and peanuts. Seeds of commodities can also be fumigated with ProFume. A complete listing of commodities can be found on the labeling for the country in which ProFume is registered.



Lack of Reactivity with Inert Materials:

ProFume does not react with commodity packaging or building components and contents, including computers, electronics and manufacturing equipment. Testing of computers conducted at 122°F (50°C) and up to tenfold the maximum permissible label dosage of ProFume (15,000 oz-h/1000 ft³ [= g-h/m³]) showed no adverse effects on computers, even when a computer was left on during fumigation (Bell et al. 2003). Additional testing was conducted with copper tubing exposed to threefold the maximum label dosage at temperatures up to 104°F (40°C) and relative humidity up to 70%. The copper tubing showed


no discoloration immediately after fumigation and for the two-month observation period following fumigation (Bell et al. 2003).

The chemical properties of sulfuryl fluoride do not result in fluoride deposition on inert surfaces. Research conducted by DFA showed no detectable increase in fluoride residues on stainless steel, glass and ceramic surfaces following fumigation with ProFume® gas fumigant at the maximum label dosage at 95°F (35°C) for 24 hours (Nead-Nylander and Thoms 2013).

Stewardship: Douglas Products has a rigorous stewardship policy for all its sulfuryl fluoride products. Fumigators using ProFume are required to participate in training programs, adhere to a written stewardship policy and allow Douglas Products or a distributor representative for ProFume to observe their initial ProFume fumigation.

Registration: The first registration received for ProFume was in Switzerland in 2003. ProFume was first registered in the United States in 2004. As of 2017, ProFume is registered in 18 countries, including Australia, Canada, Mexico and many countries in Europe. Douglas Products continues to work on registering ProFume for use in additional countries.

Targeted Pests: The pests for which ProFume is used depends upon the use patterns and provisions in the labeling. In the United States, ProFume is used to control all life stages (egg, larva, pupa and adult) of insects, such as Indian meal moth (*Plodia interpunctella*), Mediterranean flour moth (*Ephestia kuehniella*), codling moth (*Cydia pomonella*), navel orange worm (*Amyelois transitella*), flour beetles (*Tribolium* spp.), sawtoothed grain beetle (*Oryzaephilus suranamensis*), warehouse beetle (*Trogoderma variable*), granary weevil (*Sitophilus granarius*), rice weevil (*Sitophilus oryzae*), lesser grain borer (*Rhizopertha dominica*), hide beetle (*Dermestes maculatus*), drugstore beetle (*Stegobium paniceum*), rusty grain beetle (*Cryptolestes ferrugineus*), almond moth (*Ephestes cautella*), cigarette beetle (*Lasioderma serricorne*), cowpea weevil (*Callosobruchus maculatus*), bean weevil (*Ancanthoscelides obtectus*), cocoa moth (*Ephestia elutella*), cockroach species, and spiders, other invertebrate pests, and rats and mice.



Indian meal moth Lesser grain borer Warehouse beetle

Insect photos courtesy of the Canadian Grain Commission.

Dosage: Fumigation dosage is measured as a product of both concentration (C) and exposure time (T) per unit area (C x T). Because of variables in structures, the environment, pest species to be controlled and other fumigation factors, each fumigation is different, and therefore the amount of ProFume needed per unit area is variable. The computer-based Fumiguide® program calculates dosages based on input of a wide range of fumigation variables. The data used to produce the Fumiguide are the result of 10 years of research by six stored-product research laboratories in the United States and Europe (see "Pest Control Studies" on page 4) and nearly 1,200 bioassays of the key stored product insects evaluated during 51 commercial fumigations.

Precision Fumigation™: Precision Fumigation tools and techniques have been researched, developed and promoted for ProFume. Precision Fumigation can be defined as optimizing fumigant use to maximize efficiency and minimize risk. Precision Fumigation is achieved by integrating all the factors affecting control, such as pest biology, temperature, exposure time and improved sealing techniques into the fumigation management plan.

Mode of Action: Once sulfuryl fluoride enters an insect or other arthropod through the spiracles in postembryonic life stages, or diffusion through openings in the egg shell, the compound is broken down to the insecticidally active fluoride anion. The fluoride anion disrupts the glycolysis and fatty acid cycles, depriving the insect of necessary cellular energy. Insecticidal activity results from fluoride inhibition of enzyme systems utilizing magnesium within the glycolysis cycle (Meikle et al. 1963). After inhibition of the glycolysis and fatty acid cycles, insects attempt to utilize protein and amino acids to maintain a viable energy level. However, these alternative energy-producing processes are insufficient to maintain a proper metabolic rate for survival.

Resistance Management: Laboratory trials (Jagadeesan et al. 2015, Jagadeesan and Nayak 2017, Nayak et al. 2016) and field trials in bulk grain storages (Nayak et al. 2016, Opit et al. 2016) have verified that label dosage rates of ProFume® gas fumigant provide effective control of all life stages of key phosphine-resistant insect pests, including the lesser grain borer (*R. dominica*), red flour beetle (*Tribolium castaneum*), rice weevil, (*S. oryzae*) and rusty grain beetle (*C. ferrugineus*). These trials verified no cross-resistance between phosphine and sulfuryl fluoride.

Modeling studies conducted for Indian meal moth (*P. interpunctella*) and red flour beetle (*T. castaneum*) indicated that there is a very low probability for resistance development because of high rates of insect immigration, low selection pressure, no known cross-resistance and overlapping generations (Prabhakaran et al. 2001).

Due to its efficacy for controlling phosphine-resistant insect pests, ProFume is used as a rotational treatment to prevent/delay development of insecticide resistance or as a tool to control phosphine-resistant insect populations.

Description of Chemistry

Active Substance Name		Physical Properties	
IUPAC Name	sulfuryl fluoride	Molecular Weight	102.1 AMU
CAS Name	sulfuryl fluoride	Odor	Odorless
CAS Registry Number	2699-79-8	Specific Gravity	4.18 mg/ml at 25° C and 760 mm Hg
Common Name	sulfuryl fluoride, sulphuryl fluoride	Melting Point	-136.7° C (-93.7° F) at 760 mm Hg
Structure and Empirical Formula	SO ₂ F ₂	Boiling Point	-55.2° C (-67° F) at 760 mm Hg
		Vapor Pressure	12,087 mm Hg at 20° C
		Flammability	Nonflammable
		Octanol Water Partition Coefficient (log KOW)	0.14 at 20° C
		Solubility in Water	750 mg/kg at 25° C
		Solvent Solubility	1-Octanol 14 g/l Heptane 22 g/l Ethyl Acetate 59 g/l 1,2 dichloroethane 25 g/l Methanol 33 g/l Acetone 71 g/l

Thermal Degradation: In the presence of extremely high temperatures exceeding 752°F (400°C), such as an open flame or glowing heat element, sulfuryl fluoride will degrade to form hydrogen fluoride (HF). HF dissolves in water to form hydrofluoric acid, which can etch glass, metal and ceramic surfaces near the heat source. Thus, prior to fumigation, all open flames and glowing heat filaments are turned off or disconnected within the fumigated space.

Mammalian Toxicity

Introduction: Sulfuryl fluoride is an odorless, colorless gas and at low concentrations, nonirritating to mucous membranes and gives no warning of its presence. ProFume® gas fumigant is toxic and must be handled carefully in regard to the potential hazards it presents. Therefore, ProFume is labeled for use only by certified fumigators who are trained in proper fumigation techniques as part of the Douglas Products stewardship program.

The toxicology of sulfuryl fluoride has been extensively studied; a review of this research is summarized by Eisenbrandt and Hotchkiss (2010). Governmental fluoride standards have been set after extensive review of toxicological, medical and epidemiological data that included consideration of women and children. Reviews were completed by the World Health Organization (WHO) in 1984; the U.S. Public Health Service, Department of Health and Human Services in 1991 and 2015; and the National Research Council in 1993 and 2006.

Metabolism: Inhalation is the primary route of exposure to ProFume. Metabolism studies in laboratory animals indicate that sulfuryl fluoride is rapidly hydrolyzed in respiratory tissues to fluorosulfate, with release of fluoride, followed by further hydrolysis to sulfate and release of additional fluoride. This rapid metabolism results in dose-dependent increases in fluoride in the blood, tissues and urine following inhalation exposure to sulfuryl fluoride. On the other hand, parent sulfuryl fluoride is not present in blood, tissues or urine. Sulfate is a normal constituent of the body, is generally regarded as safe and thus is unlikely to contribute to the systemic toxicity of sulfuryl fluoride. Fluorosulfate and fluoride are rapidly eliminated from the plasma and tissues and excreted in the urine. Fluoride is cleared rapidly from plasma by the kidney as well as through uptake into bone. A comprehensive review of the pharmacokinetics of fluoride is provided by Whitford (1996).

Acute Toxicity: Oral ingestion of sulfuryl fluoride is unlikely because of its physical properties. Sulfuryl fluoride is not toxic dermally; the gas is not absorbed through the skin in acutely toxic amounts. Rats exposed dermally for four hours to concentrations of 9599 ppm did not show evidence of toxicity. The main route of exposure is through inhalation. Like other fumigants, sulfuryl fluoride can cause adverse effects after acute inhalation exposure, depending on the exposure concentration and duration. A summary of acute toxicology data is provided below:

Study Type	Animal	Sex	Results
Acute Oral	Rat and Guinea Pig	n/a	LD ₅₀ 100 mg/kg
Acute Dermal	Rat, F-344	Male and Female	4-Hr LC ₅₀ 1122 ppm males 4-Hr LC ₅₀ 991 ppm females
Acute Inhalation	Rat, F-344	Male and Female	1-Hr LC ₅₀ 3730 ppm males 4-Hr LC ₅₀ 3021 ppm females
Acute Inhalation	Mice, B6C3F1	Male and Female	4-Hr LC ₅₀ 400-600 ppm
Acute Inhalation	Mice, CD1	Male and Female	4-Hr LC ₅₀ 400-600 ppm

Subchronic Toxicity Studies: A summary of subchronic toxicology data is provided below:

Study Type	Animal	Sex	NOEL
Dietary Study	Rat	Male and Female	19 ppm
13 wk Inhalation	Rat, F-344	Male and Female	30 ppm
13 wk Inhalation	Rabbit, NZW	Male and Female	30 ppm
13 wk Inhalation	Mice, CD-1	Male and Female	30 ppm
13 wk Inhalation	Dog, Beagles	Male and Female	100 ppm

Chronic Toxicity and Repeated Exposure Toxicity Studies: Sulfuryl fluoride has been studied extensively in a variety of toxicological studies in laboratory animals, including repeated exposure studies. These studies varied from two weeks to two years in duration and evaluated sulfuryl fluoride concentrations above those permitted for human exposure (Eisenbrandt and Hotchkiss 2010). The toxicity of sulfuryl fluoride is species, dose and time dependent. Repeated inhalation exposure (six hours/day, five or seven days/week) of laboratory animals to sulfuryl fluoride results in inflammation of the upper and lower respiratory tissues (portal of entry irritant). Repeated exposure to sulfuryl fluoride at higher concentrations results in target organ effects in the kidney, thyroid and brain. Inhalation exposure to sulfuryl fluoride does not result in developmental effects and does not have effects on reproduction. Lifetime studies in which rats and mice were exposed to sulfuryl fluoride to assess if the chemical has potential to cause cancer also were negative. Therefore, sulfuryl fluoride is not teratogenic or carcinogenic.

Remarkably little difference in effects was observed among the species of laboratory animals in the sulfuryl fluoride repeated exposure toxicological studies (Eisenbrandt and Hotchkiss 2010). The no-observable-effect levels (NOELs) for studies that were two to 13 weeks in duration (six hours/day, five days/week exposures) were in the range of 30 to 100 ppm sulfuryl fluoride. The NOELs for specialized neurotoxicity studies were the same or higher than the NOELs for general toxicity. Importantly, an acute neurotoxicity study demonstrated that rats exposed for six hours a day for two days to 100 ppm or 300 ppm did not have signs of neurotoxicity.

Ecotoxicology and Environmental Fate

Ecotoxicology

Because ProFume® gas fumigant is a gas and applied in closed spaces, the likelihood of exposure to nontarget terrestrial and aquatic wildlife species is low. Ecotoxicology studies required for labeling and classification purposes have been conducted, and summaries of data are below:

Study Type	Species and Strain	Value
Acute Toxicity	Rainbow Trout	96h LC ₅₀ 0.89 mg/L
Acute Toxicity	<i>Daphnia Magna</i>	48h EC ₅₀ 0.62 mg/L
Algal Growth Inhibition	<i>Selenastrum Capricornutum</i>	72hEC ₅₀ 0.58 mg/L

Environmental Fate

Dow AgroSciences was awarded the Stratospheric Ozone Protection Award by the EPA in 2002 for the development of ProFume, and both the United Nations Montreal Protocol Innovators Award and the EPA's "Best of the Best" Ozone Protection Award in 2007. These awards were given to recognize the innovation required to develop ProFume for global use to protect the stratospheric ozone layer.

Fate in Air: When ProFume gas fumigant is aerated from a structure or commodity, it rapidly dissipates into the atmosphere because of its high vapor pressure. The relatively small amounts of ProFume released are calculated to have virtually no impact on the global atmosphere and environment. Sulfuryl fluoride is fully oxidized, and thus is not expected to interact or contribute to local ozone formation such as smog because of its low reactivity in the atmosphere. Sulfuryl fluoride is broken down mainly through hydrolysis to release fluoride and sulfate. The relative contribution of sulfuryl fluoride to acid rain is infinitesimally small compared to the massive amount of sulfur released into the atmosphere from industry. Sulfuryl fluoride contains no chlorine or bromine and thus cannot react to deplete stratospheric ozone by the known mechanisms (Bailey 1992).

Sulfuryl fluoride is not listed as a greenhouse gas under the Kyoto Protocol, which is the international treaty regulating greenhouse gases, and more recent climate agreements such as the Paris Agreement. Compared to greenhouse gases, such as carbon dioxide, methane and nitrous oxide, listed in the Kyoto Protocol and other climate agreements, the amount of sulfuryl fluoride emitted into the atmosphere is extremely small. Based on published estimates of the global warming potential of sulfuryl fluoride (Papadimitriou et al. 2008, Carpenter et al. 2014, Rigby et al. 2014), the current and projected annual emissions of sulfuryl fluoride would be equivalent to only about 0.05% of the total anthropogenic global carbon dioxide emission. During their evaluation of sulfuryl fluoride in 2008 and 2009, European authorities concluded that emissions of sulfuryl fluoride represented a negligible share of global greenhouse gas emissions. As part of its stewardship program, Douglas Products supports efforts for ongoing monitoring of trace atmospheric levels of sulfuryl fluoride through the Advanced Global Atmospheric Gases Experiment (AGAGE) project. Monitoring results from AGAGE continues to confirm the extremely low atmospheric levels and negligible contributions of sulfuryl fluoride as a potential greenhouse gas.

Fate in Soil: Predictive models using the physical properties of sulfuryl fluoride estimate that less than 0.1% of sulfuryl fluoride will be found in the soil at equilibrium. This is due to the high vapor pressure, which results in rapid dissipation into the atmosphere.

Fate in Water: Sulfuryl fluoride quickly hydrolyzes in water to form fluorosulfate and fluoride. Degradation rates increase with increases in aqueous pH. The half-life for sulfuryl fluoride in water with pH of 5.9, 8.1 and 9.2 is three days, 18 minutes and 1.8 minutes, respectively.

Worker Exposure

- Like all fumigants, sulfuryl fluoride can cause adverse effects after acute exposure, depending on the exposure concentration and duration. Therefore, ProFume® gas fumigant is labeled for use only by trained fumigators certified/licensed by their state or national regulatory authority to fumigate.
- Exposure guidelines: Permissible Exposure Limits (PELs) for sulfuryl fluoride for workers are based on a Time Weighted Average (TWA), are established by national regulatory authorities and vary globally from 1 to 5 ppm. Examples are given below:
 - *United States:* Both the American Conference of Governmental Industrial Hygienists (ACGIH) Threshold Limit Value (TLV) and Occupational Safety and Health Administration (OSHA) PEL are 5 ppm for workers. The EPA PEL for fumigation workers, specifically, is 1 ppm.
 - *Canada:* The Pest Management Regulatory Agency (PMRA) established a 1 ppm PEL for workers.
 - *Australia:* The Australian Pesticide and Veterinary Medicines Authority (AVPMA) established a 3 ppm PEL for workers.
 - *Europe:* The European Food Safety Authority (EFSA) established a 3 ppm PEL for workers, which is followed in labeling for European countries where ProFume is registered for application, with the exception of France (1 ppm for biocidal use and 2 ppm for phytosanitary use).
- A positive-pressure self-contained breathing apparatus (SCBA) is necessary when entering areas being fumigated where the concentration is unknown or is greater than the PEL (1 to 3 ppm, depending upon country) as measured by a detection device with sufficient sensitivity. Country regulations may specify that a fumigator wear an SCBA when conducting specific fumigation tasks, such as during application of ProFume.
- Pharmacokinetic modeling studies of potential human exposure scenarios at 1 ppm sulfuryl fluoride show that blood levels of fluoride from sulfuryl fluoride will be minimal (Poet et al. 2012). The results show that peak plasma fluoride derived from sulfuryl fluoride will be lower in adults and children re-entering a recently fumigated structure than would be expected from drinking water. The predicted fluoride levels in plasma also are lower in workers exposed during a typical work year of fumigating structures than from fluoride exposures from water or several dietary sources of fluoride.
- Eye protection requires wearing goggles or full-face shield during fumigant introduction. No special skin protection is required. Skin contact with the liquid may cause freeze damage where the liquid is confined to the skin.

Formulations:

ProFume is packaged in steel cylinders as a liquid under pressure, containing 99.8% sulfuryl fluoride (125 pounds/57 kg per cylinder) with no other pesticides, solvents or additives.

References Cited

- American Association of Cereal Chemists (AACC). 2000. Approved methods of the AACC, 10th ed. AACC, St. Paul, MN.
- Bailey, R. 1992. Sulfuryl fluoride: Fate in the atmosphere. Dow Chemical Company. DECO-ES Report 2511. Midland, Michigan.
- Baltaci, D., D. Klementz B. Gerowitt, M. J. Drinkall, and Ch. Reichmuth. 2008. Sulfuryl fluoride against all life stages of Rust red grain beetle (*Cryptolestes ferrugineus*) and Merchant grain beetle (*Oryzaephilus mercator*). In: Obenauf, G. L. (ed.), Proceedings of the 2008 Annual International Research Conference on Methyl Bromide Alternatives and Emissions Reductions, Nov. 11-14, 2008, Orlando, FL.
- Baltaci, D., D. Klementz, B. Gerowitt, M. J. Drinkall, and Ch. Reichmuth. 2009. Lethal effects of sulfuryl fluoride on eggs of different ages and other life stages of the warehouse moth *Ephestia elutella* (Hubner). *J. Stored Prod. Res.* 45: 19-23.
- Barakat, D. A., D. Klementz, and Ch. Reichmuth. 2009. Response of eggs of *Corcyra cephalonica* towards sulfuryl fluoride. In: Obenauf, G. L. (ed.), Proceedings of the 2009 Annual International Research Conference on Methyl Bromide Alternatives and Emissions Reductions, Nov. 9-12, 2009, San Diego, CA.
- Bell, C. H. 2006. Factors affecting the efficacy of sulphuryl fluoride as a fumigant. In: Lorini, I., B. Bacaltchuk, H. Beckel, D. Deckers, E. Sundfeld, J. P. dos Santos, J. D. Biagi, J. C. Celaro, L. R. D'A. Faroni, L. de O. F. Bortolini, M. R. Sartori, M. C. Elias, R. N. C. Guedes, R. G. da Fonseca, V. M. Scussel (eds.), Proceedings of the 9th International Working Conference on Stored Product Protection, October 15-18, 2006, Campinas, São Paulo, Brazil. Brazilian Post-Harvest Association-ABRAPOS, Passo Fundo, RS, Brazil, 2006, pp. 519-526.
- Bell, C. H. and N. Savvidou. 1999. The toxicity of Vikane (sulfuryl fluoride) to age groups of eggs of the Mediterranean flour moth (*Ephestia kuehniella*). *J. Econ. Entomol.* 35: 233-247.
- Bell, C. H., N. Savvidou, and T. J. Wontner-Smith. 1999. The toxicity of sulfuryl fluoride (Vikane®) to eggs of insect pests of flour mills. In: Jin, Z., Q. Liang, Y. Liang, X. Tan, and L. Guan, (eds.), Proceedings of the 7th International Working Conference on Stored-Product Protection, 14-19 October 1998, Beijing, China. Sichuan Publishing House of Science and Technology, Chengdu, China, 1999, pp. 345-355.
- Bell, C. H., T. J. Wontner-Smith and N. Savvidou. 2003. Some properties of sulfuryl fluoride in relation to its use as a fumigant in the cereals industry. In: Credland, P. F., D. M. Armitage, C. H. Bell, P. M. Cogan, E. Highley (eds.), Proceedings of the 8th International Working Conference on Stored Product Protection, July 22-26, 2002, York, UK. CAB International, Wallingford, United Kingdom, pp. 910-915.
- Bookout, A. and G. Milyo. 2006. Commercial use of ProFume® on stored cocoa beans. In: Obenauf, G. L. (ed.), Proceedings of the 2008 Annual International Research Conference on Methyl Bromide Alternatives and Emissions Reductions, Nov. 6-9, 2006, Orlando, FL.
- Carpenter, L. J., S. Reimann, J. B. Burkholder, C. Clerbaux, B. D. Hall, R. Hossaini, J. C. Laube, and S. A. Yvon-Lewis. 2014. Ozone-Depleting Substances (ODSs) and Other Gases of Interest to the Montreal Protocol, Chapter 1 in Scientific Assessment of Ozone Depletion: 2014, Global Ozone Research and Monitoring Project – Report No. 55, World Meteorological Organization, Geneva, Switzerland.
- Drinkall, M. J., V. Zaffagnini, L. Süß, and D. P. Locatelli. 2003. Efficacy of sulfuryl fluoride on stored product insects in a semolina mill trial in Italy. In: Credland, P.F., D. M. Armitage, C. H. Bell, P. M. Cogan, E. Highley (eds.), Proceedings of the 8th International Working Conference on Stored Product Protection, July 22-26, 2002, York, UK. CAB International, Wallingford, United Kingdom, pp. 884-887.
- Ducom, P., S. Dupuis, V. Stefanini, and A. A. Guichard. 2003. Sulfuryl fluoride as a new fumigant for the disinfestation of flour mills in France. In: Credland, P.F., D. M. Armitage, C. H. Bell, P. M. Cogan, E. Highley (eds.), Proceedings of the 8th International Working Conference on Stored Product Protection, July 22-26, 2002, York, UK. CAB International, Wallingford, United Kingdom, pp. 900-903.
- Eisenbrandt, D. L. and J. A. Hotchkiss. 2010. Sulfuryl fluoride. In: Hayes Handbook of Pesticide Toxicology, Vol. 2, R. Krieger, ed. Elsevier Inc., pp. 2245-2258.
- Hartzer, M., B. Subramanyam, M. Brijwani, W. Chayaprasert, and D. E. Maier. 2010. Methyl bromide, sulfuryl fluoride, and heat: Effectiveness against red flour beetle. In: Obenauf, G. L. (ed.), Proceedings of the 2010 Annual International Research Conference on Methyl Bromide Alternatives and Emissions Reductions, Nov. 2-5, 2010, Orlando, FL.
- Jagadeesan, R. and Nayak, M. K. 2017. Phosphine resistance does not confer cross-resistance to sulfuryl fluoride in four major stored grain insect pests. *Pest Manag. Sci.* 73: 1391-1401.
- Jagadeesan, R., M. K. Nayak, H. Pavic, K. Chndra, and P. J. Collins. 2015. Susceptibility to sulfuryl fluoride and lack of cross-resistance to phosphine in developmental stages of the red flour beetle, *Tribolium castaneum* (Coleoptera: Tenebrionidae). *Pest Manag. Sci.* 71: 1379-1386.
- Meikle, R. W. 1964. The fate of sulfuryl fluoride in wheat flour. *J. Ag. Food Chem.* 12: 464-467.
- Meikle, R. W. and D. J. Stewart. 1962. Structural fumigants, the residue potential of sulfuryl fluoride, methyl bromide, and methanesulfonyl fluoride in structural fumigations. *J. Ag. Food Chem.* 10: 393-379.
- Meikle, R. W., D. J. Stewart, and G. A. Globus. 1963. Drywood termite metabolism of Vikane fumigant as shown by labeled pool technique. *J. Ag. Food Chem.* 11: 226-230.
- Nayak, M. K., R. Jagadeesan, R. R. Kaur, G. J. Daglish, R. Reid, H. Pavic, L. W. Smith, and P. J. Collins. 2016. Use of Sulfuryl fluoride in the Management of Strongly Phosphine-Resistant Insect Pest Populations in Bulk Grain Storages in Australia. *J. Grain Storage Research*: 100-107.
- Nead-Nylander, Barb, and E. M. Thoms. 2013. Update on ProFume® gas fumigant as a global, post-harvest fumigant. In: Obenauf, G. L. (ed.), Proceedings of the 2013 Annual International Research Conference on Methyl Bromide Alternatives and Emissions Reductions, Nov. 4-7, 2013, San Diego, CA.
- Opit, G. P., E. M. Thoms, T. W. Phillips, and M. E. Payton. 2016. Effectiveness of sulfuryl fluoride fumigation for control of phosphine-resistant grain insects infesting stored wheat. *J. Econ. Entomol.* 109: 930-941.
- Papadimitriou, V. C., R. W. Portmann, D. W. Fahey, J. Mühle, R. F. Weiss, and J. B. Burkholder. 2008. Experimental and Theoretical Study of the Atmospheric Chemistry and Global Warming Potential of SO₂F₂. *J. of Physical Chem.* 112:12657-12666.
- Pimentel, D. 1991. Ecology and Management of Food-Industry Pests (J.R. Gorham, ed.) Association of Official Analytical Chemists, Arlington, p. 5.
- Poet, T. S., P. M. Hinderliter, D. L. Eisenbrandt, G. M. Whitford, and M. J. Bartels. 2012. PBPK Modeling of species- and age-specific sulfuryl fluoride dose metrics. Society of Toxicology, San Francisco, CA, March 2012; The Toxicologist 126: #1871.
- Prabhakaran, S., B. M. Schneider, R. E. Williams, and S. L. Ray. 2001. Computer modeling for simulating stored products insect pest population dynamics and insecticide resistance potential in food processing plants, pp. 523 – 529. In: Donahaye, E. J., Navarro, S. and Leesch J. G. (eds.), Proceedings of the International Conference of Controlled Atmosphere and Fumigation in Stored Products, Fresno, CA, Oct. 29-Nov. 3, 2000, Executive Printing Services, Clovis, CA.
- Reichmuth, Ch., W. Rassmann, G. Binker, G. Fröba, and M. J. Drinkall. 2003. Disinfestation of rust-red flour beetle (*Tribolium castaneum*), saw-toothed grain beetle (*Oryzaephilus surinamensis*), yellow meal worm (*Tenebrio molitor*), Mediterranean flour moth (*Ephestia kuehniella*) and Indian meal moth (*Plodia interpunctella*) with sulfuryl fluoride in flour mills. In: Credland, P.F., D. M. Armitage, C. H. Bell, P. M. Cogan, E. Highley (eds.), Proceedings of the 8th International Working Conference on Stored Product Protection, July 22-26, 2002, York, UK. CAB International, Wallingford, United Kingdom, pp. 736-738.



®Trademark of Douglas Products.
ProFume is a federally Restricted Use Pesticide. Always read and follow label directions.
These materials have been created specifically for ProFume and no other fumigant.
These materials may not be copied, whole or in part, or reproduced without the
permission of Douglas Products.
©2018 Douglas Products. U01-129-032 (01/18) DP

