






## REVIEW

Agronomy, Soils, and Environmental Quality

# A global meta-analysis of cover crop response on soil carbon storage within a corn production system

Deepak R. Joshi<sup>1</sup>  | Heidi L. Sieverding<sup>2</sup> | Hui Xu<sup>3</sup> | Hoyoung Kwon<sup>3</sup> |  
 Michael Wang<sup>3</sup>  | Sharon A Clay<sup>1</sup>  | Jane M. Johnson<sup>4</sup> | Resham Thapa<sup>5</sup>  |  
 Shaina Westhoff<sup>1</sup> | David E. Clay<sup>1</sup> 

<sup>1</sup>Department of Agronomy, Horticulture & Plant Science, South Dakota State University, Brookings, South Dakota, USA

<sup>2</sup>Department of Civil and Environmental Engineering, South Dakota School of Mines & Technology, Rapid City, South Dakota, USA

<sup>3</sup>Energy Systems and Infrastructure Analysis Division, Argonne National Laboratory, Lemont, Illinois, USA

<sup>4</sup>North Central Soil Conservation Research Laboratory, USDA Agriculture Research Service, Morris, Minnesota, USA

<sup>5</sup>Department of Agriculture and Environmental Science, Tennessee State University, Nashville, Tennessee, USA

## Correspondence

David E. Clay, South Dakota State University, Brookings, SD, USA.  
 Email: david.clay@sdstate.edu

Assigned to Associate Editor Michael Kaiser

## Funding information

SD Corn Utilization Council; South Dakota State University; USDA-NRCS; U.S. Department of Energy, Grant/Award Number: DE-AC02-06CH11357; U.S. Department of Agriculture

## Abstract

By influencing soil organic carbon (SOC), cover crops play a key role in shaping soil health and hence the system's long-term sustainability. However, the magnitude by which cover crops impacts SOC depends on multiple factors, including soil type, climate, crop rotation, tillage type, cover crop growth, and years under management. To elucidate how these multiple factors influence the relative impact of cover crops on SOC, we conducted a meta-analysis on the impacts of cover crops within rotations that included corn (*Zea mays* L.) on SOC accumulation. Information on climatic conditions, soil characteristics, management, and cover crop performance was extracted, resulting in 198 paired comparisons from 61 peer-reviewed studies. Over the course of each study, cover crops on average increased SOC by 7.3% (95% CI, 4.9%–9.6%). Furthermore, the impact of cover crop-induced increases in percent change SOC was evaluated across soil textures, cover crop types, crop rotations, biomass amounts, cover crop durations, tillage practices, and climatic zones. Our results suggest that current cover crop-based corn production systems are sequestering 5.5 million Mg of SOC per year in the United States and have the potential to sequester 175 million Mg SOC per year globally. These findings can be used to improve carbon footprint calculations and develop science-based policy recommendations. Taken altogether, cover cropping is a promising strategy to sequester atmospheric C and hence make corn production systems more resilient to changing climates.

**Abbreviations:**  $\bar{X}_{CC}$ , mean SOC stock for the cover crop treatment;  $\bar{X}_{NCC}$ , mean SOC stock for no cover crop treatment; CO<sub>2e</sub>, carbon dioxide equivalence; GHG, greenhouse gas; ln(*R*), natural log of response ratios; *N*<sub>CC</sub>, number of replications for the cover crop treatment; *N*<sub>NCC</sub>, number of replications for no cover crop treatment; SD, standard deviation; SOC, soil organic carbon; SOM, soil organic matter.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial](https://creativecommons.org/licenses/by-nc/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2023 The Authors. *Agronomy Journal* published by Wiley Periodicals LLC on behalf of American Society of Agronomy.

## 1 | INTRODUCTION

In 2020, 1162 million metric tons of corn (*Zea mays* L.) grain was produced globally on 202 million hectares of land (FAO-STAT, 2022). Corn grain is used to produce many products including human food, animal feed, energy products, plastics, cosmetics, diapers, and baby powder (Erenstein et al., 2021; Grote et al., 2021). However, because growing corn can contribute to greenhouse gas (GHG) emissions, it is important to minimize corn's carbon footprint (Chaplot & Smith, 2022; Lee et al., 2021). It has been hypothesized that the carbon footprint can be reduced by growing cover crops within the corn production system (Joshi et al., 2022).

Cover crops are plants that typically are not intended to be harvested and are used to reduce erosion by covering the soil between two cash crops. Farmers have many management options when growing cover crops including what, when, and where to plant (Reese et al., 2014). Interactions among management, climate, and soil conditions dictate cover crop performance, which ultimately impacts cash crop yields and the carbon dioxide equivalence ( $CO_{2e}$ ) (Abdalla et al., 2019; Jian et al., 2020; Joshi et al., 2022; McClelland et al., 2021; Poeplau & Don, 2015). For example, in arid and semiarid climates, water used by the cover crop can reduce cash crop yields, whereas in temperate environments cover crops can improve soil and plant health (Reese et al., 2014).

The  $CO_2$  equivalent ( $CO_{2e}$ ) is used to reduce the complexity of GHG emissions from multiple gases into a single value (Joshi et al., 2022). In crop production, the dominant GHG considered in  $CO_{2e}$  calculations are  $N_2O$ ,  $CO_2$ , and  $CH_4$ . This paper considers only one of those gases,  $CO_2$ , because it is directly related to SOC. Carbon dioxide emissions can be assessed by several approaches including direct emission measurement, the prediction of emissions with models, the measurement of temporal changes in SOC, or by some combination of all three (Joshi et al., 2022). This meta-analysis is based on the reported temporal changes in SOC stocks in field experiments that contained cover crop and no cover crop treatments.

Cover crops have been reported to have a mixed effect on the amount of carbon sequestered in the soil (Blanco-Canqui, 2022). In a review of US studies, Blanco-Canqui (2022) found that only 29% of the total 77 paired comparisons reported higher SOC stocks in cover crops as compared to no cover crop treatments. The positive effect of cover crops on SOC was attributed to soils with low initial SOC, higher cover crop biomass production, and the use of cover crops for many years. In 71% of the total comparisons included in his study, cover crops had no effect on SOC stocks. Blanco-Canqui (2022) attributed this lack of impact to many factors including tillage, cover crop species, fertilization, irrigation, initial SOC, soil texture, and climate. Blanco-Canqui (2022) did not conduct a quantitative synthesis because most studies do not provide the

### Core Ideas

- In the 61 studies that contained cover crops, the average SOC increase over each project was 7.3%.
- The amount of SOC stored was influenced by crop rotation, cover crop type and biomass, soil texture, and climate.
- Average cover crops increased SOC storage in the surface 30 cm was 0.88 Mg SOC (ha × per year).
- Globally, cover crop-based corn production systems have the potential to sequester 175 million Mg SOC per year.

required information (variances). Fortunately, techniques are available to overcome this barrier (Adams et al., 1997; Basche & DeLong, 2017; Hedges et al., 1999).

Previous meta-analytic studies determined cover crop effects on SOC including a wide range of crop rotations (Abdalla et al., 2019; Jian et al., 2020; McClelland et al., 2021; Poeplau & Don, 2015). However, our current study is unique in that it focuses on corn cropping systems and it considers studies within and outside the United States. The objective of this paper was to determine the impact of cover crops on SOC accumulation within rotations that included corn crops.

## 2 | MATERIALS AND METHODS

### 2.1 | Literature search and data extraction

This meta-analysis was conducted by searching for digital online peer-reviewed articles that were published prior to May 2022 (Figure 1). In a search of the Web of Science and Google Scholar, relevant articles were collected followed by data extracted, an assessment of data quality, and statistical analysis and interpretation. The keywords used in the Web of Science and Google Scholar search were soil organic matter, soil organic carbon, soil carbon, soil C, corn, maize, *Zea mays*, cover crop, green manure, rye, oat, vetch, and catch crop. This search resulted in 3856 published articles, which were published prior to May 2022. In addition to the publication date, the articles had to meet the following inclusion criteria: (1) corn had to be included in the rotation; (2) changes in soil organic carbon (SOC) had to be reported; (3) the study had to contain cover crop and no cover crop treatments; (4) the cover crop was not harvested, but was terminated or incorporated, and (5) the replicated field experiment had to be completed for at least 2 years. Because many studies were missing critical information that could not be obtained elsewhere, only 61 were selected for data extraction (Supporting Information).




FIGURE 1 Workflow diagram for peer-reviewed papers selection during meta-analysis.

During data extraction, information on crop rotation, tillage type, cover crop type and biomass produced, method and timing of cover crop termination, fertilizer application, crop yield, location (latitude and longitude), annual temperature and precipitation, soil organic carbon (SOC) and depth of sampling, soil pH, texture, bulk density (bd), when the study was initiated and completed, number of replications, and irrigation were extracted. Whenever total carbon in soil was reported as soil organic matter (SOM), it was converted to soil organic carbon by assuming that organic matter contained 58% carbon (Xu et al., 2019).

Bulk density was used in the model building, as well as to convert gravimetric values to volumetric amounts using the following equations as reported by Xu et al. (2019):

$$\text{SOC (Mg ha}^{-1}\text{)} = \text{SOC (\%)} \times \text{soil increment (cm)} \times \text{bd (g cm}^{-3}\text{)} \quad (1)$$

$$\text{SOC (Mg ha}^{-1}\text{)} = \text{SOC (g kg}^{-1}\text{)} \times \text{soil increment (cm)} \times \text{bd (g cm}^{-3}\text{)} \times 0.1 \quad (2)$$

Questions and gaps in the databases were filled by contacting the authors, extracting soil information from the Web Soil Survey for US studies and ISRIC SoilGrids for non-US studies. Missing climate information was obtained from NOAA (2022). Where possible, soil information was standardized to three soil depths (0–15, 0–30, and 0–60 cm). In the case of

multiple soil depth measurements at the same site, SOC for the whole soil profile, that is, from surface to deepest layer, was used in this meta-analysis. This was done to avoid pseudoreplication that may result when data from multiple soil depths are included in the analysis.

Whenever initial SOC amounts were not provided, it was assumed that the initial SOC stocks were identical for the cover crop and no cover treatments. If there was a difference in the initial SOC stocks between treatments, we either added or subtracted the difference in the final SOC stocks as explained by Xu et al. (2019). Moreover, the SOC stocks were standardized to the 0–15, 0–30, and 0–60 cm depths. For example, if a particular study reported data differently than our standardized soil depth increments, such as 0–10, 10–20, and 20–30 cm, then bulk density along with the depth increments were used, as explained in Equation (1) and (2), to first calculate SOC stock for the respective depth increments, which were afterward added together to determine the SOC stock for the whole soil profile. Moreover, if the depth increment reported in a study did not align with our standard depth categories, we followed the approach of Xu et al. (2019) and adjusted reported values based on if the site was tilled. For tilled sites, the vertical distribution of SOC stocks was assumed to be homogeneous for the first 30 cm. Therefore, for all tilled 0–30 cm depth increments, SOC stock was determined from the surface SOC stock value reported in the study. For no-tillage sites, a conversion factor of 1.35 was used to convert SOC stock provided for the 0–20 cm depth to the 0–30 cm depth (Puget and Lal, 2005; Xu et al., 2019; Yang & Wander, 1999).

## 2.2 | Statistical analysis

The statistical analysis was separated into multiple categories that included exploratory data analysis, cumulative meta-analysis and sensitivity/publication bias analysis, and model building. In exploratory data analysis, the distribution of the study site location was graphically presented using ArcMap. In addition, the type and amount of information collected for each category was determined using frequency plots.

### 2.2.1 | Cumulative meta-analysis: Overall cover crop effects

The cumulative meta-analysis determined the overall effect of the treatments (i.e., effect sizes) on the measured parameters. In this analysis, the effect size was determined as the natural log of the response ratio using the equation,

$$\ln(R) = \ln(\bar{X}_{CC}/\bar{X}_{NCC}) = \ln(\bar{X}_{CC}) - \ln(\bar{X}_{NCC}) \quad (3)$$

where  $\ln(R)$  is the natural log of response ratios,  $\bar{X}_{CC}$  is the mean SOC stock or corn yield values for the cover crop treatment, and  $\bar{X}_{NCC}$  is the mean SOC stock or corn yield values for the no cover crop treatment (Hedges et al., 1999). In a meta-analysis, individual effect sizes are usually weighted by the inverse of sample variances to increase the weights of studies with lower variances (Philibert et al., 2012). However, because many studies do not report sample variances, standard deviation (SD), standard errors (SE), or the coefficient of variability (CV), weighting factors ( $w_i$ ) for the  $i$ th observation (i.e., individual effect sizes) were determined based on Adams et al. (1997). This approach is based on the number of replications used in each study ( $N$ ) and it was determined with the following equation:

$$w_i = (N_{CC} * N_{NCC}) / (N_{CC} + N_{NCC}) \quad (4)$$

where  $N_{CC}$  and  $N_{NCC}$  were the number of replications for the cover crop and no cover crop treatments, respectively. Nevertheless, meta-analysis employing sample variances (SD, SE, or CV) as weighting factors is regarded as a more rigorous methodology; thus, additional research is necessary to compare the reported and estimated approaches.

To account for various sources of dependencies between effect sizes within and across studies, we created a multi-level mixed-effects meta-analytic model utilizing the “nlme” package in R (Pinheiro et al., 2017; R Core Team, 2017; Van den Noortgate et al., 2013). In this model, effect sizes were assigned as a fixed effect, study/site/common controls were nested as random effects, and  $w_i$  as weighting factors (Thapa, Mirsky, et al., 2018, Thapa, Poffenberger, et al., 2018). In the

end, we estimated the robust standard error for the mean effect size by utilizing the “clubSandwich” package in R, which is cluster-based technique for robust variance estimation (Pustejovsky & Tipton, 2022; Thapa, Mirsky, et al., 2018). Using robust standard errors, the 95% confidence intervals (CIs) were calculated for the weighted mean effect sizes, that is, the natural log means [ $\ln(R)$ ]. The mean effect sizes and their 95% CIs were back-transformed to percent change in the response using the equation

$$\% \text{ change in response} = \left( e^{\overline{\ln(R)}} - 1 \right) \times 100\% \quad (5)$$

where  $\overline{\ln(R)}$  is the mean effect sizes. If the 95% CIs did not contain zero, the overall cover crop effect on the response variable was considered significantly different from the controls ( $p < 0.05$ ). The C sequestration rate, that is, the rate of change in SOC ( $\text{Mg SOC} [\text{ha} \times \text{year}]^{-1}$ ) was determined with the equation

$$\text{SOC}_{\text{rate}} = (\text{SOC}_{\text{cc},T1} - \text{SOC}_{\text{cc},T0}) / T \quad (6)$$

where  $\text{SOC}_{\text{cc},T1}$  and  $\text{SOC}_{\text{cc},T0}$  refer to the final (T1) and initial (T0) SOC amounts for the cover crop treatment after  $T$  years.

### 2.2.2 | Moderator analysis: Effect of soil, climate, and management on the overall effects

Moderator or subgroup analysis was conducted to determine if the overall cover crop effects on percent change in SOC were affected by potential co-variables. Co-variables considered in this study include soil texture, climatic zone, crop rotation, tillage type, cover crop biomass, and years under cover crop management. To perform this analysis, separate means and their 95% CIs for each of the moderators were determined by assigning each one as a sole co-variate in the original multi-level meta-analytic mixed-effects model described above. The mean cover crop effect for each moderator was considered significant if their 95% CIs did not contain zero ( $p < 0.05$ ) and the treatment was considered different if the 95% CIs did not overlap. Moderator analysis on cover crop SOC responses was conducted by grouping the metadata into the following categories:

1. Cover crop performance by the amount of cover crop biomass produced ( $\leq 3$ ,  $3-7$  and  $\geq 7$  Mg biomass  $\text{ha}^{-1}$ ),
2. The tillage type (cultivated [CT] and no-tillage [NT]),
3. Crop rotation types (corn-corn, corn-soybean, and corn-other). Here corn-other includes corn rotation with any other crops such as rice (*Oryza sativa*), sunflower

(*Helianthus annuus* L.), groundnut (*Arachis hypogaea*), tomato (*Solanum lycopersicum*), etc.

4. The cover crop type (legume, non-legume, and mixed). The most common legume cover crops were Hairy vetch (*Vicia villosa*), lupin (*Lupinus polyphyllus*), Mucuna (*Mucuna pruriens*), Sesbania (*Sesbania sesban*), and mungbean (*Vigna radiata*). The most common non-legumes were cereal rye (*Secale cereale*), canola (*Brassica napus*), radish (*Raphanus sativus*), and oat (*Avena sativa*). Cover crop mixtures contained two or more species. For example, cereal rye + hairy vetch, winter lentil (*Lens culinaris*) + wheatgrass (*Triticum*), and oat + hairy vetch.
5. Years under cover crop management. It was categorized into three different categories: <5 years, 5–10 years, and >10 years.
6. The soil textures at the study site. These textures were categorized into fine (clay, silty clay loam, clay loam, and sandy clay), medium (silt loam and loam), and coarse (sandy loam and sandy).
7. The Köppen climate zone of the study sites were tropical, temperate, and cold categories. The tropical region included: Af (tropical rainforest climate), Aw (tropical wet and dry climate), BSh (hot semi-arid climate), BSk (cold semi-arid climate), and BWh (hot desert climate) Köppen climate zones. The temperate region included Cfa (humid subtropical climate), Csa (hot summer Mediterranean climate), Cfb (temperate oceanic climate), Csb (warm summer Mediterranean climate), Cwa (monsoon subtropical climate), and Cwb (subtropical highland climate) and lastly the cold climatic region included Dfa (hot summer humid continental climate), Dfb (warm summer humid continental climate) and Dwa (monsoon-influenced hot summer humid continental climate) Köppen climate zones. The mean annual temperature for the cold, temperate, and tropical climates were 10, 16, and 22°C, respectively, and the mean annual precipitation for the cold, temperate, and tropical climates were 883, 1116, and 1184 mm, respectively.

### 2.2.3 | Publication bias and sensitivity analysis

The publication bias is classically conducted using a funnel plot analysis, but many studies did not provide sample variance information required to create meaningful funnel plots. Therefore, a histogram was constructed to check the distribution of all individual effect sizes in the dataset to test for evidence of publication bias (Basche & DeLonge, 2017; Gurevitch et al., 2001; Thapa, Mirsky, et al., 2018). We also conducted a *Jackknife* sensitivity analysis to determine the sensitivity to any given study and hence, the robustness of the analysis (Philibert et al., 2012;

Thapa, Mirsky, et al., 2018). During *Jackknife* analysis, each study was assigned a unique study ID and data from one of the studies was excluded from the database in each calculation.

### 2.2.4 | Stepwise multiple linear regression

Stepwise regression involves recursively adding and removing predictors in the predictive model to find a subset of variables that provides the best precision and accuracy (Iduseri & Osemwenkhae, 2015). A combination of forward and backward regression models was constructed by using bulk density, clay percent, sand percent, silt percent, annual temperature, annual rainfall, initial SOC stock, N fertilizer application rate, and cover crop biomass to predict the cover crop effects on SOC stocks, that is, the natural log of the response ratios (Equation 3). However, only studies that provided both cover crop biomass and initial SOC stock were included in the regression modeling. The “stepAIC” function, from the “MASS” package in R studio was used that had a combination of both forward and backward regression during model building.

## 3 | RESULTS AND DISCUSSION

### 3.1 | Database description

A total of 61 articles met the criteria for inclusion in the meta-analysis. These studies resulted in 198 pairwise comparisons from 67 sites located on 5 different continents (Figure 2). North America (62.8%) had the highest number of sites followed by Asia (11.4%), Africa (10%), South America (8.5%), and Europe (7.14%). Sixteen different countries included in the analysis were the United States, Brazil, China, India, Argentina, Bangladesh, Benin, Italy, Kenya, Mexico, Pakistan, Poland, South Africa, Spain, Sweden, and Ethiopia (Figure 2). In the United States, most of the studies were in the Cfa, Dfa, and Dfb Köppen climate zones. These climate zones are partially aligned with what is referred to as the Corn Belt. Among all studies, nine were published between 2021 and 2022, 38 were published between 2011 and 2020, 11 were published between 2001 and 2010, and three were published between 1990 and 2000. (Figure 3a).

The number of years under cover crop management varied among studies with 41, 14, 9, and 6 studies having durations of between 2 and 5, 6 and 10, 11 and 15, and 16 and 20 years, respectively (Figure 3b). The most common soil texture was medium (44.3%) followed by coarse (30%) (Figure 3c). The studies were conducted in tropical, temperate, and cold climate zones. Of these, most studies were conducted in the temperate (38.5%) and cold (45.7%) zones (Figure 3d).




FIGURE 2 Location of all study sites (green dots) in the world map.





FIGURE 3 Different categories and their distribution of studies based on (a) publication year, (b) duration of study, (c) soil texture, and (d) Köppen climate zone.

### 3.2 | Soil organic carbon change due to cover crops

Of 198 observations assessed in our analysis, cover crops had a negative effect on SOC in 25% of the comparisons and a positive effect on SOC in 75%. When weighted across all observations, cover crops when compared with the no-cover crop treatment increased the percent change in SOC by 7.3% (95% CI, 4.9%–9.6%; Figure 4). These findings were consistent with other meta-analysis studies and confirmed the positive impact of cover crops on SOC sequestration (Abdalla et al., 2019; Jian et al., 2020; McClelland et al., 2021; Poelplau & Don, 2015). The strong positive impact on the percent

change in SOC is attributed to greater C inputs via root exudates from living cover crops during growth as well as the stabilization of shoot and root biomass C from decaying cover crops following its termination (Crystal-Ornelas et al., 2021; Janzen et al., 2022).

The net effect of cover crops on SOC is impacted by what, where, how much, and how long the cover crop is grown (Figure 4). In the reported studies, increasing the amount of cover crop biomass increased SOC storage. For example, cover crops in the moderate (3–7 Mg ha<sup>-1</sup> per year<sup>-1</sup>) and high (>7 Mg ha<sup>-1</sup> per year<sup>-1</sup>) biomass categories increased the percent SOC gains by 10.9% (95% CI, 6.1%–16.1%) and 12.4% (95% CI, 5.6%–19.6%), respectively. Whereas a low




**FIGURE 4** Percent change in soil organic carbon (SOC) due to overall cover crop, cover crop biomass, cover crop types, years under cover crops, soil depth, soil texture, tillage, crop rotation, and climate. Total number of pairwise comparisons are shown in the parenthesis above the means. The percent change in SOC change was considered significant when the 95% confidence intervals (CIs) (shown as bars) did not overlap with zero. The vertical dashed line passing through zero represents no effect of cover crops.

cover crop biomass  $<3 \text{ Mg ha}^{-1}$  per year $^{-1}$  the cover crop did not increase percent SOC gains (95% CI,  $-0.5\%$ – $7.6\%$ ). These results suggest that to increase SOC, cover crop biomass may need to be  $>3 \text{ Mg ha}^{-1}$  per year. A systemic review conducted by Blanco-Canqui (2022) also reported that cover crop biomass less than  $2 \text{ Mg ha}^{-1}$  per year may have no effect on SOC stocks and that SOC gains increase with greater biomass production. McClelland et al. (2021) also found that SOC stocks increased substantially with greater cover crop biomass production ( $>7 \text{ Mg ha}^{-1}$  per year).

Our meta-regression analysis further corroborates these previous studies by showing a positive relationship between

individual mean effect sizes of cover crops on SOC and above-ground cover crop biomass ( $p < 0.01$ ; Figure 5a). This observed effect suggests that degradation follows first-order kinetics (Joshi et al., 2020; Sainju & Singh, 1997). Although none of the studies assessed provided information on cover crop root biomass, a higher above-ground shoot biomass is typically associated with a higher below-ground root biomass. From a SOC perspective, cover crop root biomass may be more critical than shoot biomass as studies have pointed out that crop root contribution to SOC can be very high (Balesdent & Balabane, 1996; Gale et al., 2000; Wilhelm et al., 2004; Wilts et al., 2004). To address the relative contribution



**FIGURE 5** Meta-regression analysis showing the relationship between the ln response ratio (soil organic carbon [SOC] stock in cover crop/SOC stock in no-cover crop) and cover crop biomass production (a), and years under cover crop management (b). Different color points represent different cover crop types. The solid line represents the linear model between predicted ( $y$ ) and measured ( $x$ ) values. The horizontal dashed line passing through zero represents the line between SOC gains (>0) and losses (<0).

of shoots and roots future studies should quantify both cover crop shoot and root biomass to better elucidate SOC responses to cover crop performance.

All cover crop types increased SOC when compared to no cover crops (Figure 4). For instance, legume cover crops increased SOC by 8.7% (95% CI, 5.5%–12.0%), whereas the SOC gains due to multi-species cover crop mixtures and non-legumes were 6.6% (95% CI, 2.6%–10.9%) and 6.7% (95% CI, 4.1%–9.4%) respectively. The % change in SOC did not differ among cover crop types. This finding is consistent with previous studies that also observe a lack of SOC response to cover crop species (Abdalla et al., 2019; Poeplau & Don, 2015).


The length of time that cover crops were utilized had a mixed impact on SOC gains, which were 7.7% (95% CI, 4.3%–11.14%), 6.0% (95% CI, 2.1%–10.1%), and 8.4% (95% CI, 2.8%–14.2%) for systems that had contained cover crops for <5 years, 5–10 years, and >10 years, respectively (Figure 4). Our lack of difference between SOC gains and cover crop duration is attributed to the large variation in the data set (Figure 5b). These results were aligned with the analysis that showed that the length of time that cover crops were adopted did not impact SOC gains ( $p = 0.926$ , Figure 5b). McClelland et al. (2021) had similar results. However, few others have reported that SOC gains increase with time. For example, Das et al. (2022), reported that short-term studies had a smaller impact on SOC gains than mid-term and long-term studies. Blanco-Canqui (2022) reported higher SOC gains were observed when cover crops had been adopted for at least 5 years, Crystal-Ornelas et al. (2021) observed that

the soil health benefits from adopting cover crops accrue over time, and Poeplau and Don (2015) reported that cover crops increased SOC at the rate of 0.32 Mg ha<sup>-1</sup> per year. A comparison across studies indicates that cover crop-induced SOC gains most likely increase with time.

In the 0–15 cm depth, cover crops increased the % increase in SOC by 7.7% (95% CI, 3.5%–12.0%; Figure 5). When averaged over 0–30 cm depth, the percent SOC increase was 7.8% (95% CI, 4.3%–11.3%). These results suggest that cover crops increased the percent change in SOC stocks on the surface by 30 cm. Across soil textures, cover crops also had a positive impact on SOC, with increases of 6.5% (95% CI, 2.1%–10.6%), 7.6% (95% CI, 4.1%–11.2%), and 7.9% (95% CI, 3.3%–12.7%) in fine, medium, and coarse-textured soils, respectively (Figure 4). A meta-analysis conducted by Bai et al. (2019) also reported a greater increase in SOC in coarse and medium-textured soils than in fine-textured soils. The large percent SOC change in the coarse-textured soils may be attributed to the soils having relatively low initial SOC levels (Augustin & Cihacek, 2016; Vieira et al., 2009).

In the corn followed by corn rotation, cover crops increased the percent SOC gain by 7.9% (95% CI, 4.5%–11.5%), whereas in the corn followed by soybean rotation, the cover crops increase was 4.6% (95% CI, 0.9%–8.4%; Figure 4). This apparent crop rotation effect could be caused by the surface corn residue covering the soil which slowed mineralization and/or the addition of low C/N ratio soybean residue that stimulated mineralization. Clearly, additional research is needed to better define differences between these rotations.





**FIGURE 6** Percent change in soil organic carbon (SOC) due to cover crops in conventional (CT) and no-tillage (NT) systems at 0–15, 0–30, and 0–60 cm depths. Total number of pairwise comparisons are presented in parenthesis. Error bars are 95% confidence intervals (CIs) and percent SOC change were considered significant only when 95% CIs did not overlap with zero. The vertical dashed line passing through zero represents no effect of cover crops.

Averaged across all depths, cover crops increased SOC by 7.5% (95% CI, 4.0%–11.2%) and 7.1% (95% CI, 4.2%–10.0%) in conventional (CT) and no-tillage (NT) systems, respectively (Figure 4). These results suggest that tillage did not influence the potential for cover crops to increase SOC. The lack of differences was surprising because no-tillage generally has lower mineralization rates than conventionally tilled soils. However, when the data was analyzed by soil depths, an apparent tillage effect was observed (Figure 6). For example, in the 0–15 cm soil depth, the 95% confidence interval in cover crops percent SOC gain ranged from 8.6% to 33.7%, whereas in the NT the confidence interval ranged from 0.3% to 10.5%. This apparent difference was attributed to the low number of comparisons ( $n = 7$ ) in the tilled treatment.

Cover crops increased SOC in all climate zones (Figure 4). The SOC percent increases by cover crops in cold, temperate, and tropical climates were 6.1% (95% CI, 2.6%–9.8%), 9.2% (95% CI, 2.7%–16.1%), 7.6% (95% CI, 4.1%–11.2%), respectively. The large confident intervals for the climate zones indicate that the climate zone did not affect C storage. The apparent lack of difference may be attributed to more rapid cover crop growth and more rapid cover crop decomposition in the tropical than the cold and temperate zones (McClelland et al., 2021; Snapp et al., 2005; Thapa, Poffenbarger et al., 2018). Taken together, climate can influence SOC storage by affecting both the production and decomposition of the cover crop biomass.

### 3.3 | Soil carbon sequestration potential of cover crops


The SOC rate of change for the 0–15 and 0–30 cm depths were 0.44 (95% CI, 0.15–0.72), and 0.88 (95% CI, 0.56–1.2) Mg SOC ( $\text{ha} \times \text{year}^{-1}$ ), respectively. By subtracting these two depths from each other, SOC sequestration rate in the 15–30 cm depth was calculated. This calculation suggests that almost the same amount of carbon was stored in the 0–15 cm depth (0.44 Mg SOC ( $\text{ha} \times \text{year}^{-1}$ )) and 15–30 cm (0.44 Mg SOC ( $\text{ha} \times \text{year}^{-1}$ )) depth. The lack of differences between the 0–15 and 15–30 cm depths may be attributed to higher SOC mineralization rates in the 0–15 than the 15–30 cm soil depth (Clay et al., 2015). The annual amount of carbon sequestered was comparable to others. Poeplau and Don (2015) reported that for the 0–22 cm soil depth, SOC was sequestered at a rate of 0.32 Mg SOC ( $\text{ha} \times \text{year}^{-1}$ ), whereas Blanco-Canqui (2022) reported that SOC sequestration rates ranged from 0.2 to 0.9 Mg SOC ( $\text{ha} \times \text{year}^{-1}$ ) for the 0–30 cm soil depth.

Based on the amount of US cover crop cultivated land seeded to corn (~6.2 million ha; Cruthfield, 2016), ~5.5 (95% CI, 3.1–7.4) million Mg of SOC per year are being sequestered annually. If all US corn fields (36.4 million ha) used cover crops, 32.0 million Mg SOC per year might be sequestered annually, which would result in a  $\text{CO}_2\text{e}$  value of 107 million metric tons. When extended over the globe, the 200 million ha of soil seeded to corn has the potential to sequester ~175 (95% CI, 112–240) million Mg SOC per year. Sieverding et al. (2020) used data provided by Pelton (2019) to determine that the  $\text{CO}_2\text{e}$  for corn grown in the United States was ~3309 kg  $\text{CO}_2\text{e} \text{ ha}^{-1}$ . Liu et al. (2020) reported that the  $\text{CO}_2\text{e}$  could be reduced by 4453 kg  $\text{CO}_2\text{e} \text{ ha}^{-1}$  for corn with cover crops that are grown in Nebraska.

### 3.4 | Modeling cover crop-induced changes in SOC stocks

To predict the change in SOC stocks caused by cover crops, 10 factors were utilized as variables in a stepwise multiple linear regression model, with  $\ln(R)$  as the response variable. The  $\ln(R)$  value ranged from negative to zero, which indicates a reduction to no change in SOC due to cover crop cultivation, whereas positive  $\ln(R)$  indicates an increase in SOC under cover crops. Out of 10 different variables, cover crop-induced changes in SOC stocks were best predicted by bulk density, clay percent, cover crop biomass temperature, and initial SOC. The resulting equation was,

$$\ln(R) = -0.44 + 0.26 \times \text{bulk density} - 0.0025 \times \text{Clay} \\ + 0.004 \times \text{cover crop biomass} + 0.02 \times \text{temperature} \\ - 0.001 \times \text{Initial SOC}, R^2 = 0.61, n = 48,$$



**FIGURE 7** Percent change in corn yield across different cropping systems due to different cover crop types compared with no cover crop. Total number of pairwise comparisons are presented in parenthesis. Error bars are 95% confidence intervals (CIs), and percent yield change was considered significant only when 95% CIs did not overlap with zero.

$$p - \text{value} < 0.001 \quad (6)$$

where clay was in percent, cover crop biomass was in  $\text{Mg ha}^{-1}$  per year, and annual temperature was in  $^{\circ}\text{C}$ . This result suggests that increases in SOC with cover crops decrease with increasing SOC and clay content and increase with cover crop biomass production. This interpretation is consistent with Blanco-Canqui (2022) and Poeplau and Don (2015). Another potential factor affecting  $\ln(R)$  may be years under cover crop management. However, dataset used in the model ( $n = 48$ ) was dominated with studies that had cover crop management for 5 years or less, hence it was not included as model input. The exclusion of years of cover crop from the model suggests that more long-term studies are needed.


### 3.5 | Cover crop impact on corn yield

Contrasting results have been recorded globally about the impact of cover crops on the yield of the main crop. For example, cover crops can reduce corn yields (Eckert, 1991; Olson et al., 2014; Ruis et al., 2017), not influence yields (Bich et al., 2014), and increase crop yields (Astier et al., 2006; Calegari et al., 2008; Fronning et al., 2008; Reese et al., 2014). Mixed findings for cover crops impact on corn yields may be attributed to cover crops and main crops competing for water, nutrients, and light (Munawar et al., 1990) and improving nutrient and water use efficiency (Thapa, Poffenbarger, et al., 2018; Thorup-Kristensen et al., 2003). Despite the multiple agroecosystem benefits from cover crops including SOC sequestration, uncertainties in the yield impacts of cover crops most likely have slowed farmer adoption of cover crops (Singer et al., 2007).

In our meta-analysis, only 27 studies provided corn yield information resulting in a total of 93 pair-wise comparisons. Due to the relatively small sample size that had high variability, care must be used in interpreting this information. When the data were pooled across all cropping systems, legume cover crops increased corn yields by 34.9% (95% CI, 11.6%–63.3%; Figure 7). This large yield increase was not expected and may be attributed to the small number of comparisons included in the analysis, low yields in the studies that provided yield data, and/or that the cash crop did not have an adequate amount of N. Legume cover crops can fix atmospheric N which may be made available to the cash crop following mineralization (Daryanto et al., 2018; Marcillo & Miguez, 2017; Thapa, Tully et al., 2022a, Thapa, Cabrera, et al., 2022; Thorup-Kristensen et al., 2003). Planting cover crops not only adds N to the crop that follows, but also provides other ecosystem services that have a positive effect on subsequent crop yields. For instance, cover crops can improve soil properties, conserve soil moisture, and suppress weeds (Joshi et al., 2022; Thorup-Kristensen et al., 2003). By preserving soil moisture, cover crop residues help main crops use water more effectively, resulting in high N uptake even at low soil inorganic N conditions (Frye et al., 1988).

## 4 | PUBLICATION BIAS AND SENSITIVITY ANALYSIS

The distribution of individual effect sizes indicating cover crop effects on SOC was presented in Figure 8b. The histogram showed that the observations were close to normal distribution, and that the meta-analysis was not subject to publication bias. Moreover, *Jackknife* sensitivity analysis showed that no one study appeared to have a disproportionate impact



**FIGURE 8** The (a) distribution of natural log of response ratios and (b) normally distributed histogram showing no evidence for publication bias for cover crops effects on soil organic carbon (SOC) stocks across the 198 observations. Sensitivity analysis (c) was conducted using the *Jackknife* technique. The overall percent change in SOC stocks is shown by the solid red line, and the lower and higher 95% confidence intervals are also provided as dashed red lines. The removal of any single study had no effect on the results.

on the results and that the cover crops' overall effect size estimates for SOC obtained in this meta-analysis was robust (Figure 8c).

## 5 | LIMITATIONS AND FUTURE STUDY CONSIDERATIONS

Many of the papers failed to report important information used in the meta-analysis. For example, initial SOC, changes in SOC over the entire rooting zone, cover crop biomass produced, cash crop yields, pH, bulk densities, soil texture, and nutrient concentrations were often not reported. Moreover, many studies did not report the measured sample variance values such as standard deviation, standard errors, or the coefficient of variability. Therefore, providing such information in future studies will be helpful for a more robust meta-analysis. Also not reporting important information, such as yields of the cash crop, or the amount of cover crop biomass reduces the ability to evaluate important interactions. Others have noted similar data gaps with which data are reported (Abdalla et al., 2019; Jian et al., 2020; Poelplau & Don, 2015). To improve global predictive models,

these database gaps need to be minimized. Overall, our study found that cover crops have a positive effect on SOC sequestration. However, we did not consider different greenhouse gas ( $\text{NO}_2$  and  $\text{CH}_4$ ) emission during the growth and decomposition to determine the full carbon footprint of cover crop. Therefore, assessing SOC stocks along with greenhouse gases in response to cover crops should be considered in the future to increase our understanding of the corn cover crop system.

## 6 | CONCLUSION

In this meta-analysis, data from 61 publications were used to determine the effects of cover crops on SOC stocks and corn yields within a corn cropping system. A database was created by extracting information provided by individual studies along with soils, climate, management, and cover crop information. Overall, integrating cover crops in corn production systems increased SOC stocks by 7.3% (95% CI, 4.9%–9.6%). The SOC stock responses for cover crops varied by location depending on soil (texture and initial SOC), climate, management (cover crop types, tillage, and crop

rotations), and cover crop biomass production. The SOC gains from cover crop adoption are more likely to be observed in low-C and coarse-textured soils. Similarly, cover crop biomass production had a strong positive effect on the magnitude of SOC accumulation. Cover crop performance depends on soil nutrient status, climate, management (cover crop types, cover crop planting, and termination dates/methods), and growing window depending on main crop rotations.

Warmer growing conditions and more rainfall in temperate and tropical climatic zones support higher cover crop biomass production and return rates. However, higher temperatures also help to accelerate cover crop decomposition. The net SOC gains were similar across climate zones. Based on these results, we can conclude that soil, climate, and management influence both cover crop performance and its ability for SOC sequestration.

At an average SOC sequestration rate of 0.88 Mg ha<sup>-1</sup> per year at 0–30 cm, current corn fields with cover crops are potentially sequestering 5.5 million Mg of SOC-C per year in the United States and 160 million Mg SOC per year globally. If all US corn fields used cover crops, 32.1 million Mg SOC per year could be sequestered annually in the United States, which would result in a CO<sub>2e</sub> value of 107 million metric tons.

These findings imply that cover crop-induced increases in SOC can improve soil health and soils' ability to adapt to changing climates while also mitigating it. In the long term, improved SOC under cover crops will lead to better soil functions and productivity. Findings from this study can be used to identify areas that may have the greatest potential to sequester carbon and also shape management decisions for optimizing SOC storage under cover crops. Taken altogether, we conclude that growing cover crops on croplands rather than leaving them in a fallow phase can be one of the strategies to sequester atmospheric C and mitigate greenhouse gas emissions, thereby potentially enhancing the sustainability of corn production systems worldwide.

## AUTHOR CONTRIBUTIONS

**Deepak R. Joshi:** Conceptualization; data curation; formal analysis; investigation; methodology; resources; software; validation; visualization; writing—original draft; writing—review and editing. **Heidi L. Sieverding:** Conceptualization; data curation; methodology; resources; writing—review and editing. **Hui Xu:** Conceptualization; data curation; methodology; software; validation; writing—review and editing. **Hoyoung Kwon:** Conceptualization; methodology; resources; validation; writing—review and editing. **Michael Wang:** Conceptualization; data curation; methodology; resources; validation; writing—review and editing. **Sharon A Clay:** Conceptualization; resources; supervision; validation; writing—review and editing. **Jane M. Johnson:** Resources; supervision; validation; writing—review editing. **Resham Thapa:** Formal analysis; writing—review and

editing. **Shaina Westhoff:** Conceptualization; methodology; writing—review and editing. **David E. Clay:** Conceptualization; funding acquisition; project administration; resources; supervision; validation; writing—review and editing.

## ACKNOWLEDGMENTS

Funding was provided by SD Corn Utilization Council, USDA-NRCS, South Dakota State University Experiment Station, and US-DOE. Argonne National Laboratory's contribution to this research was supported by the Bioenergy Technologies Office of the Office of Energy Efficiency and Renewable Energy of the US Department of Energy, under contract number DE-AC02-06CH11357. Trade names or commercial products mentioned in this article are solely for the purpose of providing specific information and do not infer either recommendation or endorsement by the US Department of Agriculture. The findings and conclusions in this publication are those of the authors and should not be construed to represent any official USDA or US Government determination or policy. USDA is an equal opportunity provider and employer.

## CONFLICT OF INTEREST STATEMENT

Authors declare no conflicts of interest.

## ORCID

Deepak R. Joshi  <https://orcid.org/0000-0002-5142-284X>  
 Michael Wang  <https://orcid.org/0000-0002-8315-0201>  
 Sharon A Clay  <https://orcid.org/0000-0003-4166-6995>  
 Resham Thapa  <https://orcid.org/0000-0002-0059-764X>  
 David E. Clay  <https://orcid.org/0000-0002-5031-9744>

## REFERENCES

- Abdalla, M., Hastings, A., Cheng, K., Yue, Q., Chadwick, D., Espenberg, M., Truu, J., Rees, R. M., & Smith, P. (2019). A critical review of the impacts of cover crops on nitrogen leaching, net greenhouse gas balance and crop productivity. *Global Change Biology*, 25, 2530–2543. <https://doi.org/10.1111/gcb.14644>
- Adams, D. C., Gurevitch, J., & Rosenberg, M. S. (1997). Resampling tests for meta-analysis of ecological data. *Ecology*, 78, 1277–1283. [https://doi.org/10.1890/0012-9658\(1997\)078\[1277:RTFMAO\]2.0.CO;2](https://doi.org/10.1890/0012-9658(1997)078[1277:RTFMAO]2.0.CO;2)
- Astier, M., Maass, J. M., Etchevers-Barra, J. D., Peña, J. J., & González, F. D. L. (2006). Short-term green manure and tillage management effects on maize yield and soil quality in an Andisol. *Soil and Tillage Research*, 88, 153–159. <https://doi.org/10.1016/j.still.2005.05.003>
- Augustin, C., & Cihacek, L. J. (2016). Relationships between soil carbon and soil texture in the Northern Great Plains. *Soil Science*, 181, 386–392. <https://doi.org/10.1097/SS.0000000000000173>
- Bai, X., Huang, Y., Ren, W., Coyne, M., Jacinthe, P.-A., Tao, B., Hui, D., Yang, J., & Matocha, C. (2019). Responses of soil carbon sequestration to climate-smart agriculture practices: A meta-analysis. *Global Change Biology*, 25, 2591–2606. <https://doi.org/10.1111/gcb.14658>
- Basche, A., & Delongue, M. (2017). The impact of continuous living cover on soil hydrologic properties: A meta-analysis. *Soil Science Society of*

- America Journal*, 81, 1179–1190. <https://doi.org/10.2136/sssaj2017.03.0077>
- Bich, A. D., Reese, C. L., Kennedy, A. C., Clay, D. E., & Clay, S. A. (2014). Corn yield is not reduced by mid-season establishment of cover crops in northern Great Plains environments. *Crop Management*, 13(1), 1–8. <https://doi.org/10.2134/CM-2014-0009-RS>
- Blanco-Canqui, H. (2022). Cover crops and carbon sequestration: Lessons from US studies. *Soil Science Society of America Journal*, 86, 501–519. <https://doi.org/10.1002/saj2.20378>
- Balesdent, J., & Balabane, M. (1996). Major contribution of roots to soil carbon storage inferred from maize cultivated soils. *Soil Biology & Biochemistry*, 28(9), 1261–1263.
- Calegari, A., Hargrove, W. L., Rheinheimer, D. D. S., Ralisch, R., Tessier, D., Tourdonnet, S., & Fatima Guimarães, M. (2008). Impact of long-term no-tillage and cropping system management on soil organic carbon in an Oxisol: A model for sustainability. *Agronomy Journal*, 100, 1013–1019. <https://doi.org/10.2134/agronj2007.0121>
- Chaplot, V., & Smith, P. (2022). Cropping leads to loss of soil organic matter: How can we prevent it? *Pedosphere*, 33, 9–10. <https://doi.org/10.1016/j.pedsph.2022.06.002>
- Clay, D. E., Reicks, G., Carlson, C. G., Moriles-Miller, J., Stone, J. J., & Clay, S. A. (2015). Tillage and corn residue harvesting impact surface and subsurface carbon sequestration. *Journal of Environmental Quality*, 44, 803–809. <https://doi.org/10.2134/jeq2014.07.0322>
- Crutchfield, J. (2016). USDA Foreign Agricultural Service overview for operational monitoring of current crop conditions and production forecasts, AGU Fall Meeting Abstracts. pp. GC53A–1260.
- Crystal-Ornelas, R., Thapa, R., & Tully, K. L. (2021). Soil organic carbon is affected by organic amendments, conservation tillage, and cover cropping in organic farming systems: A meta-analysis. *Agriculture, Ecosystems, and Environment*, 312, 107356. <https://doi.org/10.1016/j.agee.2021.107356>
- Daryanto, S., Fu, B., Wang, L., Jacinthe, P.-A., & Zhao, W. (2018). Quantitative synthesis on the ecosystem services of cover crops. *Earth-Science Reviews*, 185, 357–373. <https://doi.org/10.1016/j.earscirev.2018.06.013>
- Das, S., Chatterjee, S., & Rajbanshi, J. (2022). Responses of soil organic carbon to conservation practices including climate-smart agriculture in tropical and subtropical regions: A meta-analysis. *Science of the Total Environment*, 805, 0048–9697. <https://doi.org/10.1016/j.scitotenv.2021.150428>
- Eckert, D. J. (1991). Chemical attributes of soils subjected to no-till cropping with rye cover crops. *Soil Science Society of America Journal*, 55, 405–409. <https://doi.org/10.2136/sssaj1991.03615995005500020019x>
- Erenstein, O., Chamberlin, J., & Sonder, K. (2021). Estimating the global number and distribution of maize and wheat farms. *Global Food Security*, 30, 100558. <https://doi.org/10.1016/j.gfs.2021.100558>
- FAOSTAT. (2022). Database. FAO. <http://www.fao.org/faostat/en/#data>
- Fronning, B. E., Thelen, K. D., & Min, D.-H. (2008). Use of manure, compost, and cover crops to supplant crop residue carbon in corn stover removed cropping systems. *Agronomy Journal*, 100, 1703–1710. <https://doi.org/10.2134/agronj2008.0052>
- Frye, W. W., Blevins, R. L., Smith, M. S., Corak, S. J., & Varco, J. J. (1988). Role of annual legumes cover crops in efficient use of water and nitrogen. In W. L. Hargrove (Ed.), *Cropping strategies for efficient use of water and nitrogen*, (pp. 129–154). ASA.
- Gale, W. J., Cambardella, C. A., & Bailey, T. B. (2000). Root-derived carbon and the formation and stabilization of aggregates. *Soil Science Society of America Journal*, 64, 201–207. <https://doi.org/10.2136/sssaj2000.641201x>
- Grote, U., Fasse, A., Nguyen, T. T., & Erenstein, O. (2021). Food security and the dynamics of wheat and maize value chains in Africa and Asia. *Frontiers in Sustainable Food Systems*, 4, Article 617009. <https://doi.org/10.3389/fsufs.2020.617009>
- Gurevitch, J., Curtis, P. S., & Jones, M. H. (2001). Meta-analysis in ecology. *Advances in Ecological Research*, 199, 247. [https://doi.org/10.1016/S0065-2504\(01\)32013-5](https://doi.org/10.1016/S0065-2504(01)32013-5)
- Hedges, L. V., Gurevitch, J., & Curtis, P. S. (1999). The meta-analysis of response ratios in experimental ecology. *Ecology*, 80, 1150–1156. [https://doi.org/10.1890/0012-9658\(1999\)080%5b1150:TMAORR%5d2.0.CO;2](https://doi.org/10.1890/0012-9658(1999)080%5b1150:TMAORR%5d2.0.CO;2)
- Iduseri, A., & Osemwenkhae, J. E. (2015). An efficient variable selection method for predictive discriminant analysis. *Annals of Data Science*, 2, 489–504. <https://doi.org/10.1007/s40745-015-0061-9>
- Janzen, H. H., Van Groenigen, K. J., Powlson, D. S., Schwinghamer, T., & Van Groenigen, J. W. (2022). Photosynthetic limits on carbon sequestration in croplands. *Geoderma*, 416, 115810. <https://doi.org/10.1016/j.geoderma.2022.115810>
- Jian, J., Du, X., Reiter, M. S., & Stewart, R. D. (2020). A meta-analysis of global cropland soil carbon changes due to cover cropping. *Soil Biology and Biochemistry*, 143, 107735. <https://doi.org/10.1016/j.soilbio.2020.107735>
- Joshi, D. R., Clay, D. E., Clay, S. A., & Smart, A. J. (2020). Seasonal losses of surface litter in northern great plains mixed-grass prairies. *Rangeland Ecology & Management*, 73, 259–264. <https://doi.org/10.1016/j.rama.2019.11.003>
- Joshi, D. R., Clay, D. E., Clay, S. A., Moriles-Miller, J., Daigh, A. L. M., Reicks, G., & Westhoff, S. (2022). Quantification and machine learning based N<sub>2</sub>O–N and CO<sub>2</sub>–C emissions predictions from a decomposing rye cover crop. *Agronomy Journal*, 1–15. <https://doi.org/10.1002/agj2.21185>
- Lee, U., Kwon, H., Wu, M., & Wang, M. (2021). Retrospective analysis of the US corn ethanol industry for 2005–2019: Implications for greenhouse gas emission reductions. *Biofuels, Bioproducts and Biorefining*, 15, 1318–1331. <https://doi.org/10.1002/bbb.2225>
- Liu, X., Kwon, H., Northrup, D., & Wang, M. (2020). Shifting agricultural practices to produce sustainable, low carbon intensity feedstocks for biofuel production. *Environmental Research Letters*, 15, 8084014. <https://doi.org/10.1088/1748-9326/ab794e>
- Marcillo, G. S., & Miguez, F. E. (2017). Corn yield response to winter cover crops: An updated meta-analysis. *Journal of Soil and Water Conservation*, 72, 226. <https://doi.org/10.2489/jswc.72.3.226>
- McClelland, S. C., Paustian, K., & Schipanski, M. E. (2021). Management of cover crops in temperate climates influences soil organic carbon stocks: A meta-analysis. *Ecological Applications*, 31, e02278. <https://doi.org/10.1002/eap.2278>
- Munawar, A., Blevins, R. L., Frye, W. W., & Saul, M. R. (1990). Tillage and cover crop management for soil water conservation. *Agronomy Journal*, 82, 773–777. <https://doi.org/10.2134/agronj1990.00021962008200040024x>
- NOAA, National Centers for Environmental Information. (2022). *Climate data online: Dataset discovery*. <https://www.ncdc.noaa.gov/cdo-web/datasets>
- Olson, K., Ebelhar, S. A., & Lang, J. M. (2014). Long-term effects of cover crops on crop yields, soil organic carbon stocks and

- sequestration. *Open Journal of Soil Science*, 4, 284–292. <https://doi.org/10.4236/ojss.2014.48030>
- Pelton, R. (2019). Spatial greenhouse gas emissions from US county corn production. *International Journal of Life Cycle Assessment*, 24, 13–25. <https://doi.org/10.1007/s11367-018-1506-0>
- Philibert, A., Loyce, C., & Makowski, D. (2012). Assessment of the quality of meta-analysis in agronomy. *Agriculture, Ecosystems & Environment*, 148, 72–82.
- Pinheiro, J., Bates, D., DebRoy, S., & Sarkar, D. (2017). *nlme: Linear and nonlinear mixed effects models. R package version 3.1-131*. R Foundation for Statistical Computing.
- Poepflau, C., & Don, A. (2015). Carbon sequestration in agricultural soils via cultivation of cover crops: A meta-analysis. *Agriculture, Ecosystems & Environment*, 200, 33–41.
- Puget, P., & Lal, R. (2005). Soil organic carbon and nitrogen in a Mollisol in central Ohio as affected by tillage and land use. *Soil and Tillage Research*, 80, 201–213. <https://doi.org/10.1016/j.still.2004.03.018>
- Pustejovsky, J. E., & Tipton, E. (2022). Meta-analysis with robust variance estimation: Expanding the range of working models. *Prevention Science*, 23(3), 425–438. <https://doi.org/10.1007/s11121-021-01246-3>
- R Core Team. (2017). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing. <http://www.R-project.org>
- Reese, C. L., Clay, D. E., Clay, S. A., Bich, A. D., Kennedy, A. C., Hansen, S. A., & Moriles, J. (2014). Winter cover crops impact on corn production in semiarid regions. *Agronomy Journal*, 106, 1479–1488. <https://doi.org/10.2134/agronj13.0540>
- Ruis, S. J., Blanco-Canqui, H., Jasa, P. J., Ferguson, R. B., & Slater, G. (2017). Can cover crop use allow increased levels of corn residue removal for biofuel in irrigated and rainfed systems? *BioEnergy Research*, 10, 992–1004. <https://doi.org/10.1007/s12155-017-9858-z>
- Sainju, U. M., & Singh, B. P. (1997). Winter cover crops for sustainable agricultural systems. Influence on soil properties, water quality and crop yields. *Horticultural Science*, 32, 21–28.
- Sieverding, H., Kebreab, E., Johnson, J. M. F., Xu, H., Wang, M., Grosso, S. J. D., Bruggeman, S., Stewart, C. E., Westhoff, S., Ristau, J., Kumar, S., & Stone, J. J. (2020). A life cycle analysis (LCA) primer for the agricultural community. *Agronomy Journal*, 112, 3788–3788. <https://doi.org/10.1002/agj2.20279>
- Singer, J. W., Nusser, S. M., & Alf, C. J. (2007). Are cover crops being used in the US Corn Belt? *Journal of Soil and Water Conservation*, 62(5), 353–358.
- Snapp, S. S., Swinton, S. M., Labarta, R., Mutch, D., Black, J. R., Leep, R., Nyiraneza, J., & O'neil, K. (2005). Evaluating cover crops for benefits, costs and performance within cropping system niches. *Agronomy Journal*, 97, 322–332. <https://doi.org/10.2134/agronj2005.0322a>
- Thapa, R., Mirsky, S. B., & Tully, K. L. (2018). Cover crops reduce nitrate leaching in agroecosystems: A global meta-analysis. *Journal of Environmental Quality*, 47, 1400–1411. <https://doi.org/10.2134/jeq2018.03.0107>
- Thapa, R., Poffenbarger, H., Tully, K. L., Ackroyd, V. J., Kramer, M., & Mirsky, S. B. (2018). Biomass production and nitrogen accumulation by hairy vetch/cereal rye mixtures: A meta-analysis. *Agronomy Journal*, 110, 1197–1208. <https://doi.org/10.2134/agronj2017.09.0544>
- Thapa, R., Tully, K. L., Reberg-Horton, C., Cabrera, M., Davis, B. W., Fleisher, D., Gaskin, J., Hitchcock, R., Poncet, A., Schomberg, H. H., Seehaver, S. A., Timlin, D., & Mirsky, S. B. (2022). Cover crop residue decomposition in no-till cropping systems: Insights from multi-state on-farm litter bag studies. *Agriculture, Ecosystems, and Environment*, 326, 107823. <https://doi.org/10.1016/j.agee.2021.107823>
- Thapa, R., Cabrera, M., Reberg-Horton, C., Dann, C., Balkcom, K. S., Fleisher, D., Gaskin, J., Hitchcock, R., Poncet, A., Schomberg, H. H., Timlin, D., & Mirsky, S. B. (2022). Modeling surface residue decomposition and N release using the cover crop nitrogen calculator (CC-NCALC). *Nutrient Cycling in Agroecosystems*, 124(1), 81–99. <https://doi.org/10.1007/s10705-022-10223-3>
- Thorup-Kristensen, K., Magid, J., & Jensen, L. S. (2003). Catch crops and green manures as biological tools in nitrogen management in temperate zones. *Advances in Agronomy*, Academic Press, 79, 227–302. [https://doi.org/10.1016/S0065-2113\(02\)79005-6](https://doi.org/10.1016/S0065-2113(02)79005-6)
- Van Den Noortgate, W., López-López, J. A., Marín-Martínez, F., & Sánchez-Meca, J. (2013). Three level meta-analysis of dependent effect sizes. *Behavior Research Methods*, 45, 576–594. <https://doi.org/10.3758/s13428-012-0261-6>
- Vieira, F. C. B., Bayer, C., Zanatta, J. A., Mielniczuk, J., & Six, J. (2009). Building up organic matter in a subtropical Paleudult under legume cover-crop-based rotations. *Soil Science Society of America Journal*, 73(5), 1699–1706. <https://doi.org/10.2136/sssaj2008.0241>
- Wilhelm, W. W., Johnson, J. M. F., Voorhees, W. B., & Linden, D. R. (2004). Crop and soil productivity response to corn residue removal: A literature review. *Agronomy Journal*, 96, 1–17.
- Wilts, A. R., Reicosky, D. C., Allmaras, R. R., & Clapp, C. E. (2004). Long-term corn residue effects: Harvest alternatives, soil carbon turnover, and root-derived carbon. *Soil Science Society of America Journal*, 68(4), 1342–1351. <https://doi.org/10.2136/sssaj2004.1342>
- Xu, H., Sieverding, H., Kwon, H., Clay, D., Stewart, C., Johnson, J. M. F., Qin, Z., Karlen, D. L., & Wang, M. (2019). A global meta-analysis of soil organic carbon response to corn stover removal. *GCB Bioenergy*, 11, 1215–1233. <https://doi.org/10.1111/gcbb.12631>
- Yang, X.-M., & Wander, M. M. (1999). Tillage effects on soil organic carbon distribution and storage in a silt loam soil in Illinois. *Soil and Tillage Research*, 52, 1–9. [https://doi.org/10.1016/S0167-1987\(99\)00051-3](https://doi.org/10.1016/S0167-1987(99)00051-3)

## SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

**How to cite this article:** Joshi, D. R., Sieverding, H. L., Xu, H., Kwon, H., Wang, M., Clay, S. A., Johnson, J. M., Thapa, R., Westhoff, S., & Clay, D. E. (2023). A global meta-analysis of cover crop response on soil carbon storage within a corn production system. *Agronomy Journal*, 1–14. <https://doi.org/10.1002/agj2.21340>