

## ANATOMY

#### BALANCE

To maintain balance we use input from our vision (eyes), proprioception (muscles/joints), and vestibular system (inner ear).

### ARTICLE



DID THIS ARTICLE HELP YOU? SUPPORT VEDA @ VESTIBULAR.ORG

5018 NE 15th Ave. Portland, OR 97211 1-800-837-8428 info@vestibular.org vestibular.org



# **The Human Balance System:** A Complex Coordination of Central and Peripheral Systems

By Vestibular Disorders Association, with contributions by Mary Ann Watson, MA, F. Owen Black, MD, FACS, and Matthew Crowson, MD

Good balance is often taken for granted. Most people don't find it difficult to walk across a gravel driveway, transition from walking on a sidewalk to grass, or get out of bed in the middle of the night without stumbling. However, with impaired balance such activities can be extremely fatiguing and sometimes dangerous. Symptoms that accompany the unsteadiness can include dizziness, vertigo, hearing and vision problems, and difficulty with concentration and memory.

#### WHAT IS BALANCE?

Balance is the ability to maintain the body's center of mass over its base of support. 1 A properly functioning balance system allows humans to see clearly while moving, identify orientation with respect to gravity, determine direction and speed of movement, and make automatic postural adjustments to maintain posture and stability in various conditions and activities.

Balance is achieved and maintained by a complex set of sensorimotor control systems that include sensory input from vision (sight), proprioception (touch), and the vestibular system (motion, equilibrium, spatial orientation); integration of that sensory input; and motor output to the eye and body muscles. Injury, disease, certain drugs, or the aging process can affect one or more of these components. In addition to the contribution of sensory information, there may also be psychological factors that impair our sense of balance.

#### SENSORY INPUT

Maintaining balance depends on information received by the brain from three peripheral sources: eyes, muscles and joints, and vestibular organs (Figure 1 - see end of this article). All three of these information sources send signals to the brain in the form of nerve impulses from special nerve endings called sensory receptors.

#### INPUT FROM THE EYES

Sensory receptors in the retina are called rods and cones. Rods are

1

believed to be tuned better for vision in low light situations (e.g. at night time). Cones help with color vision, and the finer details of our world. When light strikes the rods and cones, they send impulses to the brain that provide visual cues identifying how a person is oriented relative to other objects. For example, as a pedestrian takes a walk along a city street, the surrounding buildings appear vertically aligned, and each storefront passed first moves into and then beyond the range of peripheral vision.

#### INPUT FROM MUSCLES AND JOINTS

Proprioceptive information from the skin, muscles, and joints involves sensory receptors that are sensitive to stretch or pressure in the surrounding tissues. For example, increased pressure is felt in the front part of the soles of the feet when a standing person leans forward. With any movement of the legs, arms, and other body parts, sensory receptors respond by sending impulses to the brain. Along with other information, these stretch and pressure cues help our brain determine where our body is in space.

The sensory impulses originating in the neck and ankles are especially important. Proprioceptive cues from the neck indicate the direction in which the head is turned. Cues from the ankles indicate the body's movement or sway relative to both the standing surface (floor or ground) and the quality of that surface (for example, hard, soft, slippery, or uneven).

#### INPUT FROM THE VESTIBULAR SYSTEM

Sensory information about motion, equilibrium, and spatial orientation is provided by the vestibular apparatus, which in each ear includes the utricle, saccule, and three semicircular canals. The utricle and saccule detect gravity (information in a vertical orientation) and linear movement. The semicircular canals, which detect rotational movement, are located at right angles to each other and are filled with a fluid called endolymph. When the head rotates in the direction sensed by a particular canal, the endolymphatic fluid within it lags behind because of inertia, and exerts pressure against the canal's sensory receptor. The receptor then sends impulses to the brain about movement from the specific canal that is stimulated. When the vestibular organs on both sides of the head are functioning properly, they send symmetrical impulses to the brain. (Impulses originating from the right side are consistent with impulses originating from the left side.)

#### INTEGRATION OF SENSORY INPUT

Balance information provided by the peripheral sensory organs-eyes, muscles and joints, and the two sides of the vestib-ular system-is sent to the brain stem. There, it is sorted out and integrated with learned information contributed by the cerebellum (the coordination center of the brain) and the cerebral cortex (the thinking and memory center). The cerebellum provides information about automatic move-ments that have been learned through repeated exposure to certain motions. For example, by repeatedly practicing serving a ball, a tennis player learns to optimize balance control during that movement. Contributions from the cerebral cortex include previously learned information; for example, because icy sidewalks are slippery, one is required to use a different pattern of movement in order to safely navigate them.



#### PROCESSING OF CONFLICTING SENSORY INPUT

A person can become disoriented if the sensory input received from his or her eyes, muscles and joints, or vestibular organs sources conflicts with one another. For example, this may occur when a person is standing next to a bus that is pulling away from the curb. The visual image of the large rolling bus may create an illusion for the pedestrian that he or she-rather than the bus-is moving. However, at the same time the proprioceptive information from his muscles and joints indicates that he is not actually moving. Sensory information provided by the vestibular organs may help override this sensory conflict. In addition, higher level thinking and memory might compel the person to glance away from the moving bus to look down in order to seek visual confirmation that his body is not moving relative to the pavement.

#### MOTOR OUTPUT

As sensory integration takes place, the brain stem transmits impulses to the muscles that control movements of the eyes, head and neck, trunk, and legs, thus allowing a person to both maintain balance and have clear vision while moving.

#### MOTOR OUTPUT TO MUSCLES AND JOINTS

A baby learns to balance through practice and repetition as impulses sent from the sensory receptors to the brain stem and then out to the muscles form a new pathway. With repetition, it becomes easier for these impulses to travel along that nerve pathway–a process called facilitation– and the baby is able to maintain balance during any activity. Strong evidence exists suggesting that such synaptic reorganiza-tion occurs throughout a person's lifetime of adjusting to changing motion environs.

This pathway facilitation is the reason dancers and athletes practice so arduously. Even very complex movements become nearly automatic over a period of time. This also means that if a problem with one sensory information input were to develop, the process of facilitation can help the balance system reset and adapt to achieve a sense of balance again. For example, when a person is turning cartwheels in a park, impulses transmitted from the brain stem inform the cerebral cortex that this particular activity is appropriately accompanied by the sight of the park whirling in circles. With more practice, the brain learns to interpret a whirling visual field as normal during this type of body rotation. Alternatively, dancers learn that in order to maintain balance while performing a series of pirouettes, they must keep their eyes fixed on one spot in the distance as long as possible while rotating their body.

#### MOTOR OUTPUT TO THE EYES

The vestibular system sends motor control signals via the nervous system to the muscles of the eyes with an automatic function called the vestibulo-ocular reflex (VOR). When the head is not moving, the number of impulses from the vestibular organs on the right side is equal to the number of impulses coming from the left side. When the head turns toward the right, the number of impulses from the right ear increases and the number from the left ear decreases. The difference in impulses sent from each side controls eye movements and stabilizes the gaze during active head movements (e.g., while running or watching a hockey game) and passive head movements (e.g., while sitting in a car that is accelerating or decelerating).

#### THE COORDINATED BALANCE SYSTEM

The human balance system involves a complex set of sensorimotor-control systems. Its interlacing feedback mechanisms can be disrupted by damage to one or more components through injury, disease, or the aging process. Impaired balance can be accompanied by other symptoms such as dizziness, vertigo, vision problems, nausea, fatigue, and concentration difficulties.

The complexity of the human balance system creates challenges in diagnosing and treating the underlying cause of imbalance. The crucial integration of information obtained through the vestibular, visual, and proprioceptive systems means that disorders affecting an individual system



#### **FIGURE 1**

can markedly disrupt a person's normal sense of balance. Vestibular dysfunction as a cause of imbalance offers a particularly intricate challenge because of the vestibular system's interaction with cognitive functioning, 2 and the degree of influence it has on the control of eye movements and posture.

#### REFERENCES

 Shumway-Cook A, Woollacott MH. Motor Control: Theory and Practical Applications. Philadelphia: Lippincott, Williams & Wilkins; 2001.

#### NOTES:

2. Hanes DA, McCollum G. Journal of Vestibular Research 2006;16(3):75-91.

©2016 Vestibular Disorders Association VeDA's publications are protected under copyright. For more information, see our permissions guide at vestibular.org. *This document is not intended as a substitute for professional health care.* 

#### VESTIBULAR DISORDERS ASSOCIATION

5018 NE 15th Ave. Portland, OR 97211 1-800-837-8428 info@vestibular.org vestibular.org

## Did this free publication from VeDA help you?

You can ensure that educational articles like this continue to be available to vestibular patients like you by making a tax-deductible gift to VeDA today.

#### SUPPORT VEDA

One-time gift:	\$40	\$50	\$75	\$100	\$250	🗌 other
Monthly gift:	\$10	\$15	\$25	\$35	\$50	other

Check this box if you prefer that your donation remain anonymous.

#### **PAYMENT INFORMATION**

Donations gladly accepted online at http://vestibular.org. Check or money order in US funds, payable to VeDA.

Visa	МС	Amex	Discover				
				Card number	Exp. date	CVV code	
Billing	g add	ress of	card (if diffe	rent from mailing information)			
MAIL	.ING	INFOR	MATION				

Name	Telephone	Email	
Address	City	State/Province	Zip
Country			