



SYMPTOMS

DIZZINESS

Dizziness, vertigo, and disequilibrium are common symptoms of vestibular disorders, which can arise from a number of causes.

ARTICLE

011

**DID THIS ARTICLE
HELP YOU?
SUPPORT VEDA @
VESTIBULAR.ORG**

5018 NE 15th Ave.
Portland, OR 97211
1-800-837-8428
info@vestibular.org
vestibular.org

Causes of Dizziness

By Vestibular Disorders Association, with edits by John King, PhD & Stacey Buckner, DPT

Dizziness, vertigo, and disequilibrium are common symptoms reported by adults during visits to their doctors. They are all symptoms that can result from a peripheral vestibular disorder (a dysfunction of the balance organs of the inner ear) or central vestibular disorder (a dysfunction of one or more parts of the central nervous system that help process balance and spatial information). Although these three symptoms can be linked by a common cause, they have different meanings, and describing them accurately can mean the difference between a successful diagnosis and one that is missed. Dizziness is a sensation of lightheadedness, faintness, or unsteadiness. Unlike dizziness, vertigo has a rotational, spinning component, and is the perception of movement, either of the self or surrounding objects. Disequilibrium simply means unsteadiness, imbalance, or loss of equilibrium that is often accompanied by spatial disorientation and dizziness, although it can occur independently.

Almost everyone experiences a few seconds of spatial disorientation at some point—for example, when a person watches a 3-D movie in the theater and momentarily perceives an illusion of moving or falling as the images rush past. However, frequent episodes of vertigo—whether lasting only for a few seconds or days on end—are a primary sign of a vestibular dysfunction, especially when linked to changes in head and body position. By contrast, dizziness can be a primary sign of a vestibular disorder in addition to a broad array of cardiovascular, neurological, metabolic, vision, and psychological problems. It is also quite possible that a person may have a combination of problems, such as a degenerative vestibular disorder along with a visual deficit such as cataracts or a neurological disorder such as a stroke.



Because of the many possible causes of dizziness, getting a correct diagnosis can be a long and frustrating experience.



DEFINITIONS

Dizziness

A sensation of lightheadedness, faintness, or unsteadiness. Dizziness does not involve a rotational component.

Vertigo

The perception of movement or whirling, either of the self or surrounding objects.

Disequilibrium

Unsteadiness, imbalance, or loss of equilibrium, often accompanied by disorientation.

Spatial disorientation

A sensation of not knowing where one's body is in relation to the vertical and horizontal planes.

DIZZINESS CAUSED BY VESTIBULAR DISORDERS

The body maintains balance with sensory information from three systems:

- **Vision**
- **Proprioception** (touch sensors in the feet, trunk, and spine)
- **Vestibular system** (inner ear)

Sensory input from these three systems is integrated and processed by the brain-stem. In response, feedback messages are sent to the eyes to help maintain steady vision and to the muscles to help maintain posture and balance.

A healthy vestibular system supplies the most reliable information about spatial orientation. Mixed signals from vision or proprioception can usually be tolerated. When sitting in a car at a railroad crossing, seeing a passing train may cause the sensation of drifting or moving, and feeling a soft, thick carpet underfoot as opposed to a solid wood floor can produce a floating sensation. However, compensating for vestibular system abnormalities is more problematic. Just as a courtroom judge rules between two sides presenting competing evidence, the vestibular system serves as the tie-breaker between conflicting forms of sensory information. When the vestibular system malfunctions, it can no longer help resolve moments of sensory conflict,

resulting in symptoms such as dizziness, vertigo, and disequilibrium.

SPECIFIC VESTIBULAR SYSTEM PROBLEMS

Vestibular dysfunction is most commonly caused by head injury, aging, and viral infection. Other illnesses, as well as genetic and environmental factors, may also cause or contribute to vestibular disorders. For more detailed information about specific vestibular system problems, please visit www.vestibular.org. Causes of dizziness related to vestibular system dysfunction are listed below.

- A **vestibular schwannoma**, also known as an acoustic neuroma, is a benign tumor growing on the vestibulo-cochlear nerve.
- **Autoimmune inner ear disease** occurs when the defense capabilities of a mal-functioning immune system harm the cells of the body that affect the ear. Specific diagnoses include Cogan's syndrome, Wegener's granulomatosis, systemic lupus, Sjogren's syndrome, and rheumatoid arthritis, among others.
- **Benign paroxysmal positional vertigo (BPPV)** is a condition resulting from loose debris (otoconia) that collect within a part of the inner ear. In addition to head injury, BPPV can occur due to the degeneration of inner-ear hair cells during the natural process of aging.
- **Cervicogenic dizziness** describes symptoms of dizziness and/or disequilibrium that arise from neck pain and issues that include cervical trauma, cervical arthritis, and other causes.
- **Cholesteatoma** is a skin growth that occurs in the middle ear behind the eardrum that can invade the surrounding skull and inner ear, thus causing lots of destruction.
- An **enlarged vestibular aqueduct** houses the fluid-filled endolymphatic duct, which is connected to the endo-lymphatic sac. The function of the duct and the sac are affected when the aqueduct is larger than normal.
- **Labyrinthitis and vestibular neuritis** are inflammations caused by a viral infection that can result in damage to hearing and vestibular function (labyrinthitis) or damage to vestibular function only (vestibular neuritis).
- **Mal de débarquement** is a sensation of rocking or movement that persists after a cruise or other form of travel.



- **Ménière's disease, or primary endolymphatic hydrops**, involves abnormalities in quantity, composition, or pressure of the endolymph (one of the fluids within the inner ear). It is a progressive condition.
- **Migraine associated vertigo (MAV)** may be characterized by head pain with symptoms associated with vestibular impairment such as dizziness, motion intolerance, spontaneous vertigo, sensitivity to light and sound, tinnitus, imbalance, and spatial disorientation.
- **Otitis media** is a bacterial infection of the middle ear and meningitis is a bacterial infection of the brain covering that may spread to the inner ear.
- **Otosclerosis** a condition in which the bones of the middle ear become fixed, leading to hearing loss. It can invade the inner ear over time.
- **Ototoxicity** is caused by exposure to certain drugs or chemicals (e.g., intra-venous aminoglycoside antibiotics) that damage the inner-ear nerve hair cells or the vestibulo-cochlear nerve.
- **Perilymph fistula**, caused by injury, is a tear or defect in the oval or round window, which are small, thin membranes that separate the middle ear from the fluid-filled inner ear.
- **Superior canal dehiscence** is an opening in the bone overlying the uppermost semicircular canal within the inner ear.
- **Secondary endolymphatic hydrops** involves abnormalities in quantity, composition, or pressure of the endolymph.
- **Vascular compression** of the vestibular nerve is an irritation of the vestibular portion of the vestibulo-cochlear nerve by a blood vessel.

NON-VESTIBULAR CAUSES OF DIZZINESS

Dizziness can be linked to a wide array of problems and is particularly commonly linked to blood-flow irregularities from cardiovascular problems. Non-vestibular causes of dizziness are listed below.

- An **aneurysm** is a weak spot in an artery wall that balloons out and allows blood to leak into the vessel walls. An aneurysm is a catastrophic event that can cause severe dizziness and difficulty with walking.
- An **arrhythmia** is an irregular or abnormal heartbeat and can result in low blood flow to



the brain, causing one to faint or feel faint.

- **Atherosclerosis** is hardening or narrowing of the vertebral arteries. In older people who have high blood pressure, plaque is sometimes deposited within the arteries. This narrows the interior of the arteries and impedes blood flow. Heredity may be a factor in development of this condition.
- **Carotid sinus reflex** works rapidly in younger people but sometimes is much slower in older people, especially those with circulatory problems. The carotid sinus is very sensitive to decreases in blood pressure in the carotid artery. With a drop in blood pressure, the reflex constricts blood vessels in the lower extremities and dilates vessels in the head to maintain a normal blood pressure in the head and adequate blood flow to the brain.
- A **defective heart valve** usually involves the aortic valve, which when shut down (aortic stenosis) prevents the proper amount of blood from flowing to the brain.
- **Dehydration** can produce light-headedness through its effect on multiple systems.
- People with **severe degenerative arthritis** of the spine can develop bone spurs that may press on the vertebral arteries and interfere with blood supply to the brain.
- **Embolism** can occur when an embolus, or blood clot, forms around a heart valve that is not working properly, or is released within the arteries to the brain, causing a stroke. The effects of a stroke may include temporary dizziness. However, if the embolus travels to the vestibular system, it can cause severe dizziness.
- A **heart attack** rarely causes dizziness; when it does, lack of blood to the brain is the cause.
- **Hyperventilation** is a condition resulting from rapid breathing, when more carbon dioxide



than normal is expelled. When this happens, the level of carbon dioxide in the blood falls and affects the function of brain cells, causing temporary dizziness.

- Certain **medications**, including some prescription and over-the-counter drugs, can cause temporary dizziness.
- **Nervous-system disorders** such as **peripheral neuropathies** (diminished nerve function in the legs or feet) and **multiple sclerosis** can cause unsteadiness.
- **Orthostatic hypotension** is common in older people, especially those with circulatory problems and diabetes. When a person has low blood pressure and pooling of blood in the lower part of the body while sitting or lying down, the process of standing up quickly can cause dizziness and fainting. Normally, body reflexes accommodate such position changes. However, when circulation problems impair these compensation mechanisms, faintness occurs.
- **Osteoarthritis** is a joint disease that can narrow the openings in the neck vertebrae (bones) through which blood vessels flow. Blockage of these vertebral arteries results in an inadequate blood supply to the base of the brain or brainstem—where the balance information is controlled. This causes symptoms of dizziness and lightheadedness. The condition is termed vertebral basilar insufficiency. If this arterial narrowing takes place gradually over time, other arteries may enlarge and take over some of the function of the affected vessels. This event, called development of a collateral blood supply, can't happen if the arterial narrowing occurs suddenly (for example, if an embolus completely shuts off the blood supply). In such cases, death by stroke may result.
- **Stress, tension, or fatigue** may cause the brain stem to function less efficiently, resulting in some loss of automatic reflex control of balance. This leads to elevated levels of activity for the cerebral cortex as it works to help maintain balance through the control of voluntary muscle movements. Lightheadedness and unsteadiness can result.
- A **tumor** may affect the brain stem, the cerebellum (the coordination center of the brain), or the part of the cerebral cortex that controls voluntary muscle movements.
- **Vasovagal syndrome** is a nervous-system response that causes sudden loss of muscle

tone in peripheral blood vessels.

- **Vision disturbances** can occur when a person adjusts to bifocals or a new eyeglass prescription, or must compensate for reduced vision due to cataracts.
- **Intracranial Hypertension** occurs when the pressure of cerebrospinal fluid (CSF) is too high.

DIZZINESS CAUSED BY MULTIPLE SENSORY DEFICITS

Maintaining balance and equilibrium can be very difficult when more than one health problem exists. A mild vestibular disorder can be much more problematic when accompanied by a visual deficit. The ability to compensate for a vestibular disorder is compounded when there is also a deficit with proprioception due to disease or an injury and severe dizziness can result. Careful evaluation, including a complete medical history noting all potential causes of dizziness, is essential to correct diagnosis and treatment.



©2014 Vestibular Disorders Association
VeDA's publications are protected under copyright.
For more information, see our permissions guide at
vestibular.org. ***This document is not intended as a
substitute for professional health care.***

[illegible]

5018 NE 15th Ave. Portland, OR 97211
1-800-837-8428 info@vestibular.org vestibular.org

You can ensure that educational articles like this continue to be available to vestibular patients like you by making a tax-deductible gift to VeDA today.

One-time gift: ☐ \$40 ☐ \$50 ☐ \$75 ☐ \$100 ☐ \$250 ☐ other

Monthly gift: ☐ \$10 ☐ \$15 ☐ \$25 ☐ \$35 ☐ \$50 ☐ other

☐ Check this box if you prefer that your donation remain anonymous.

Donations gladly accepted online at <http://vestibular.org>. Check or money order in US funds, payable to VeDA.

Visa	MC	Amex	Discover	<div><div></div><div>Card number</div></div>	<div><div></div><div>Exp. date</div></div>	<div><div></div><div>CVV code</div></div>
------	----	------	----------	--	--	---

Billing address of card (if different from mailing information)

Name _____ Telephone _____ Email _____
Address _____ City _____ State/Province _____ Zip _____
Country _____