


- Who can administer an auto injector in an anaphylaxis emergency?
 - Anyone officially trained in its use and recognising signs and symptoms
 - Only the casualty can self-administer the medication
 - Only medical professionals and the person to whom it is prescribed
 - Anyone can, it is not a complicated procedure
- Challenging child Charlie has broken gentle Jonny's jaw. What can you do for Jonny?
 - Bandage tightly around the jaw, over the head to immobilise area
 - Provide a cushioned cold pack for pain relief and transfer to hospital
 - Immobilise the jaw and neck using neck collar, call 999 for a paramedic
 - Place Jonny in the recovery position leaving Billy to look after him
- Six year old Ahava has a fading level of response, after a bad bleed. An ambulance was called 15 minutes ago, she is on her back, her legs are raised, what else do you do?
 - Move her to her side as she is about to pass out
 - There is no more you can do, the ambulance will be there soon
 - Update ambulance to the deterioration and prepare to move her to her side
 - Stimulate her to keep her awake using pain and splashes of water
- Polly has inhaled poisonous sulphuric gases in school. She is coughing slightly, but seems ok, what pro-active measures do you take?
 - Reprimand her for opening the fuming cupboard, as the room stinks now
 - Send her with a friend to sit outside, and send another child to get help
 - Tell her to let you know if she does not feel any better in a hour
 - Sit her by corridor entrance, so you can keep an eye on her
- Stacey has been stung by a bee, and has a little sack of poison still attached:
 - You flick it off and give her a drink of warm tea with honey
 - Do not remove it, apply ice and immobilise by bandaging around the sting
 - Wash the area with warm soapy water, and apply anti sting spray
 - Remove it without squeezing poison into her from the sting sack, and wash
- A baby is having a seizure because they are too hot, what do you do?
 - Call 999 and wait for help in a cool area with the casualty
 - Hold the baby close to you and gently place a wooden spoon in their mouth
 - Move the fitting baby into a sink and place in very cold water around them
 - Remove their clothes, cool them down, pad the area around the baby and time how long the seizure lasts
- Five year old Grace is frustrated as she has a grass seed in her eye, you can see it, her eye is not really hurting, but feels uncomfortable. What can you do?
 - Get a helper to hold her eye open, while you extract it using tweezers
 - Try flushing it out using saline solution or water, if no luck take her to hospital
 - Bandage both her eyes up so she cannot see, and take her to hospital
 - Leave it in there, and encourage her to cry more, it will eventually fall out

- Which is the best position for a conscious casualty with breathing problems?



- Six year old Sarah has a rash on her arm, a fever and a headache. What do you do?
 - Check rash disappears under a clear glass, call parents and send to a doctor
 - Send Sarah home and advise a good night's sleep
 - Call 999 and send to A&E department for meningitis
 - Let her continue playing and see how she is at lunch time
- A little Lego batman head is hiding deep in Bruce's ear, what is the best treatment?
 - Using water and warm oils, gently tease Batman out of the ear
 - Using tweezers and cocktail sticks, remove the Lego piece
 - Take Bruce to the A&E department, where they can remove it carefully
 - Call 999, but do nothing yourself, let the emergency services deal with it
- 8 year old Spencer has a splinter in his finger. It is small and still poking out. You would...
 - ... tell Spencer to go and wash his hands and hope it comes out
 - ... wash the area for Spencer and put a plaster on it. We are not allowed to remove splinters
 - ... carefully wash Spencer's finger, then using tweezers pull the splinter out. Make it bleed slightly and then wash again, finally covering with a plaster
 - .. call 111 and ask for advice, or 101 for less urgent assistance
- Susy has had a second seizure. She is now not moving. Can you help?
 - Yes, leave her in the recovery position and call ambulance. She might have further fits
 - No, she needs to sleep it off
 - Yes, try to give her some sugary food and drink ready for the next seizure
 - Yes, look for any medication she needs and get her to take it
- Athena has wheezing breathing problems, what's wrong with her and what can you do?
 - You diagnose a first asthma attack, lie her down on the floor
 - She is worried and panicking, help her breathe through a paper bag
 - You diagnose a first asthma attack, give her Zakie's asthma inhaler to use
 - You diagnose a first asthma attack, sit her up and call an ambulance
- Joseph is nine and feels very sick after lunch and looks pale. What can you do?
 - Give him Coca Cola™ as it helps settle the stomach
 - Ask SAMPLE questions including what he ate for lunch, be ready for him to vomit and send home
 - Encourage him to vomit as you suspect he has swollen poison
 - You think it might be poison, so lie him down and let him sleep it off



15. Three year old Timmy has bumped his mouth and a tooth came out, what can you do?
- Manage bleeding, reassure Timmy and send the tooth home for the fairies to deal with. It is only a baby tooth.
 - Gently slightly rinse the tooth with saline solution and insert it into the gum
 - Rub the tooth clean thoroughly, and then insert it into back into the gum
 - Send Timmy to a dentist or hospital
16. A dog has bitten Rebecca, leaving a small puncture wound. What can you do?
- Just clean the area. A dog's saliva is a natural antiseptic and so it will be ok
 - Apply a tight tourniquet to the area above the bite to stop infection spreading
 - Clean and cover the area, check for other injuries and then send to be checked at hospital
 - Call police to have the dog seized, and an ambulance for the bite
17. Beth's leg is completely trapped under a joist, and no blood is reaching her foot, and has been there for 10 minutes. Do you lift it off?
- No, it has been too long. She will need urgent medical specialists. Just call 999
 - Yes, it has not been too long, remove it slowly and carefully. Call 999
 - No, it might have punctured a blood vessel and may cause serious bleeding
 - Yes, but put a tourniquet above the area first to totally stop the blood flowing
18. The correct child CPR protocol is
- 2 rescue breathes, 30 compressions followed by 2 breaths again
 - 30 compressions followed by two breaths, then 30:2 repeated
 - 5 rescue breathes, 15 compressions, 2 breaths, then 15:2 repeated
 - 5 rescue breathes, 30 compressions followed by two breaths again
19. When can you give a child prescribed medication?
- If the child says it is their medication, you can give it to them
 - Only prescribed medication with parental written consent
 - Any medication prescribed to anyone if directed by a head teacher
 - Any lifesaving medication if you are sure it will help
20. A child's asthma medication Salbutamol is not working, why might that be?
- The child is not inhaling the medication, help them breathe it in
 - The inhaler is empty, find a replacement
 - It is the wrong inhaler, check they use the correct one
 - All of the above are possible



now check your answers!

Paediatric First Aid Assessment Paper 3

- As it is not a requirement for first aiders to be able to read or write English. Learners may be unfairly challenged by the paper. If this is the case please contact Learn2Rescue to discuss different assessment options.
- The paper should be undertaken in typical exam conditions.
- A maximum of 30 minutes should be allowed for candidates undertaking the assessment.
- A clock should be prominently displayed in the assessment room
- Learners should be spaced round the room and be sat at least 1.5 metres apart.
- Learners should not be allowed assistance to answer the questions, but help understanding certain words is permissible
- The assessment paper consists of 20 multiple choice questions.
- The pass mark of the paper is 15 correct answers
- Learners must demonstrate competence before commencing the practical assessments
- Any learner who fails to achieve 15 correct answers should be referred
- Do not write on the question paper
- Learn2Rescue has an appeal process for those that want to have their test results investigated

Good luck