



PROTECT THEIR PURR: FIGHT FELINE CYSTITIS

1 WHAT IS FELINE IDIOPATHIC CYSTITIS?

A build up of crystals in the urine without a presence of bacteria.



2 WHAT CAUSES FELINE IDIOPATHIC CYSTITIS?

Stress and pain can be a cause for Feline Idiopathic Cystitis. Stress in cats can be caused by many different things including cat conflict, building work, roadworks, fireworks, guests and more.




3 WHY CAN PAIN BE A CAUSE OF FELINE IDIOPATHIC CYSTITIS?

Pain and stress go hand in hand and each one exacerbates the other. Pain that affects their mobility can mean they struggle to get into litter trays or go outside to go to the toilet. It may also be uncomfortable for them to position themselves to pass urine so they start to stand to go or hold on until desperate.



4 WHY IS LITTER TRAY ACCESS IMPORTANT?

Providing extra toilet options even for cats who usually go outside can reduce stress. Creating specific outside spaces to toilet can help too. More options means less risk of house soiling and you can adapt to their individual preferences and needs.



5 INCREASE THEIR WATER INTAKE.

Increasing their water intake is key to help cats with Feline Idiopathic Cystitis. Providing extra places for them to get water and offering in raised bowls, via water fountains, on different floors of your home as well as providing safe options for outside.



**For more information on
Feline Idiopathic Cystitis head to our website.
www.purrfectfelinecare.co.uk**