



CTA-071-Undersea Landslide-GBA


Vic Ferguson

The World Federation for Coral Reef Conservation 281.971.7703 P.O. Box 311117 Houston Texas 77231

2.23.17

Largest Undersea Landslide Revealed on the Great Barrier Reef

ECO



15 FEBRUARY 2017

[PREVIOUS ARTICLE](#)Banned Chemicals in Deepest Reaches of the Ocean

[NEXT ARTICLE](#)Watch Extreme Waves in 3D via Electronic Eyes

Scientists have discovered the remnants of a massive undersea landslide on the Great Barrier Reef, approximately 30 times the volume of Uluru.

James Cook University's Dr Robin Beaman said the remains of the slip, known as the Gloria Knolls Slide, were discovered 75 kilometers off the north Queensland coast near the town of Innisfail while the scientists were working from the Marine National Facility's blue-water research ship *Southern Surveyor*.

"This is all that remains after a massive collapse of sediment of about 32 cubic kilometers' volume more than 300,000 years ago," he said.

Dr Beaman said a debris field of large blocks, or knolls, and numerous smaller blocks, lies scattered over 30 kilometers from the main landslide remains, into the Queensland Trough, to a depth of 1350 meters.



CTA-071-Undersea Landslide-GBA

Vic Ferguson

The World Federation for Coral Reef Conservation 281.971.7703 P.O. Box 311117 Houston Texas 77231

“We were amazed to discover this cluster of knolls while 3D multibeam mapping the deep GBR seafloor. In an area of the Queensland Trough that was supposed to be relatively flat were eight knolls, appearing like hills with some over 100 m high and 3 km long.”

Associate Professor Jody Webster from the University of Sydney likened the research to a detective story, first finding the knolls, then using later mapping to reveal the landslide source of the knolls.

A sediment sample from a knoll at a depth of 1170 meters identified a remarkable cold-water coral community of both living and fossil cold-water coral species, gorgonian sea whips, bamboo corals, mollusks and stalked barnacles.

“The oldest fossil corals recovered off the top of the knoll was 302 thousand years,” says Dr Angel Puga-Bernabéu at the University of Granada and lead author on the study, “which means the landslide event that caused these knolls must be older”.

Modelling the potential tsunami for a sudden ‘mass failure’ on this scale yields a three-dimensional tsunami wave elevation of about 27 meters. However, the wave would likely be dampened significantly by the presence of any coral reefs.

Considerably more seabed mapping and sampling is needed to fully assess the tsunami hazard to the Queensland coast posed by these types of underwater landslides.

The scientists said one-third of the Great Barrier Reef lies beyond the seaward edge of the shallower reefs, and the discovery of this prominent undersea landslide and its vast debris field in the deep Great Barrier Reef reveals a far more complex landscape than previously known.

This research is a collaborative effort between James Cook University, University of Sydney, University of Granada, University of Edinburgh and the Australian Nuclear Science and Technology Organization.

[Source](#)