




Haglund's Deformity

What Is Haglund's Deformity?

Haglund's deformity is a bony enlargement on the back of the heel. The soft tissue near the Achilles tendon becomes irritated when the bony enlargement rubs against shoes. This often leads to painful bursitis, which is an inflammation of the bursa (a fluid-filled sac between the tendon and bone).



Causes of Haglund's Deformity

Haglund's deformity is often called "pump bump" because the rigid backs of pump-style shoes can create pressure that aggravates the enlargement when walking. In fact, any shoes with a rigid back, such as ice skates, men's dress shoes or women's pumps, can cause this irritation.

To some extent, heredity plays a role in Haglund's deformity. Inherited foot structures that can make one prone to developing this condition include:

- A high-arched foot
- A tight Achilles tendon
- A tendency to walk on the outside of the heel.

Symptoms of Haglund's Deformity

Haglund's deformity can occur in one or both feet. The symptoms include:

- A noticeable bump on the back of the heel
- Pain in the area where the Achilles tendon attaches to the heel
- Swelling in the back of the heel
- Redness near the inflamed tissue

Diagnosis of Haglund's Deformity

After evaluating the patient's symptoms, the foot and ankle surgeon will examine the foot. In addition, x-rays will be ordered to help the surgeon evaluate the structure of the heel bone.

Nonsurgical Treatment

Nonsurgical treatment of Haglund's deformity is aimed at reducing the inflammation of the bursa. While these approaches can resolve the pain and inflammation, they will not shrink the bony protrusion. Nonsurgical treatment can include one or more of the following:

- **Medication.** Oral nonsteroidal anti-inflammatory drugs (NSAIDs), such as ibuprofen, may be recommended to reduce the pain and inflammation. Ice. To reduce swelling, apply an ice pack to the inflamed area, placing a thin towel between the ice and the skin. Use ice for 20 minutes and then wait at least 40 minutes before icing again.

- **Exercises.** Stretching exercises help relieve tension from the Achilles tendon. These exercises are especially important for the patient who has a tight heel cord.
- **Heel lifts.** Patients with high arches may find that heel lifts placed inside the shoe decrease the pressure on the heel.
- **Heel pads.** Pads placed inside the shoe cushion the heel and may help reduce irritation when walking.
- **Shoe modification.** Backless or soft backed shoes help avoid or minimize irritation.
- **Physical therapy.** Physical therapy modalities, such as ultrasound, can help to reduce inflammation.
- **Orthotic devices.** Custom arch supports control the motion in the foot.
- **Immobilization.** In some cases, casting may be necessary.

When Is Surgery Needed?

If nonsurgical treatment fails to provide adequate pain relief, surgery may be needed. The foot and ankle surgeon will determine the procedure that is best suited to your case. It is important to follow the surgeon's instructions for postsurgical care.

Prevention

To help prevent a recurrence of Haglund's deformity:

- wear appropriate shoes; avoid shoes with a rigid heel back
- use arch supports or orthotic devices
- perform stretching exercises to prevent the Achilles tendon from tightening
- avoid running on hard surfaces and running uphill

Why choose a foot and ankle surgeon?

Foot and ankle surgeons are the leading experts in foot and ankle care today. As doctors of podiatric medicine – also known as podiatrists, DPMs or occasionally “foot and ankle doctors” – they are the board-certified surgical specialists of the podiatric profession. Foot and ankle surgeons have more education and training specific to the foot and ankle than any other healthcare provider.

Foot and ankle surgeons treat all conditions affecting the foot and ankle, from the simple to the complex, in patients of all ages including Haglund's deformity. Their intensive education and training qualify foot and ankle surgeons to perform a wide range of surgeries, including any surgery that may be indicated for Haglund's deformity.