

From Peels to Plates: Unlocking the Hidden Nutritional Potential of Food By-products

Dr. Ankita Lakade^{1*} and Dr. Jyoti Nishad²
¹Amity Institute of Food Technology, Amity University, NOIDA-201313, Uttar Pradesh, India
²ICAR-Directorate of Cashew Research, Puttur, Dakshina Kannada-574202, Karnataka, India
*Corresponding author: ajlakade@amity.edu

Introduction

Every year, the global food processing industry generates enormous quantities of by-products in the form of peels, seeds, husks, pomace, and shells. It is estimated that fruit and vegetable processing alone produces by-products amounting to nearly 25 to 30 percent of the total raw material processed. Traditionally, these residues have been treated as waste, disposed of in landfills, or, at best, used as low-value animal feed. However, an emerging and rapidly growing body of research demonstrates that these by-products are far from worthless. They are, in fact, rich reservoirs of dietary fiber, vitamins, minerals, and bioactive compounds such as polyphenols, carotenoids, and flavonoids (Galanakis, 2012). The concept of transforming food by-products into value-added ingredients, popularly summarized as "from peels to plates," is gaining traction as a strategy that simultaneously addresses food waste management, environmental sustainability, and the growing consumer demand for functional and nutraceutical foods.

This article reviews the nutritional and functional potential of common food by-products, the extraction and processing technologies used to recover valuable compounds from them, their applications in the food industry, and the

challenges that must be overcome to enable large-scale commercial adoption.

The Hidden Nutritional Wealth of By-products

Fruit and vegetable peels, which are among the most commonly discarded by-products, often contain higher concentrations of certain nutrients than the edible pulp itself. Citrus peels, for instance, are rich sources of pectin, hesperidin, and naringin, compounds known for their antioxidant and other health benefits (Sharma et al., 2017). Banana peels contain notable amounts of dietary fiber, potassium, and phenolic compounds, which contribute to their antioxidant activity (Ayala-Zavala et al., 2010). Pomegranate peel, frequently discarded during juice extraction, is exceptionally rich in compounds associated with anti-inflammatory and anticancer properties (Ayala-Zavala et al., 2010).

Similarly, oilseed cakes remaining after oil extraction, such as those from groundnut, sesame, and flaxseed, retain substantial residual proteins, fibres, micronutrients and bioactive compounds (Vichare and Morya, 2024). Table 1 summarizes the nutritional composition of selected fruit and vegetable by-products commonly generated in food processing operations.

Table 1. Nutritional and Bioactive Composition of Selected Fruit and Vegetable By-products (Dry Weight Basis)

By-product	Dietary Fiber (g/100 g DW)	Total Polyphenols (g GAE/100 g DW)	Vitamin C (mg/100 g DW)	Key Bioactive Compounds (Major reported compounds)
Banana peel	45.0	2.0	12.0	Catechin, carotenoids, dopamine
Pomegranate peel	40.0	10.0	9.5	Ellagic acid, punicalagin, tannins
Tomato pomace	45.0	6.2	5.0	Lycopene, beta-carotene
Citrus peel	42.0	12.5	58.0	Hesperidin, naringin, pectin
Grape pomace	48.0	2.5	2.0	Resveratrol, anthocyanins, tannins

As shown in Table 1, citrus peel exhibits the highest dietary fiber content among the by-products compared, while pomegranate peel stands out for its comparatively high polyphenol content, underscoring the diversity of nutritional value locked within different types of food residues. It is important to note that these values are representative of the mid-range values reported in the literature. However, the actual composition can vary depending on factors such as the

cultivar, stage of ripening, extraction method, and analytical technique used.

Extraction and Processing Technologies

Recovering bioactive compounds from food by-products requires appropriate extraction and stabilization technologies. Conventional solvent extraction remains widely used, but it is increasingly being supplemented or replaced by

greener alternatives such as ultrasound-assisted extraction, microwave-assisted extraction, and supercritical fluid extraction, all of which improve yield while reducing solvent consumption and processing time based on the depends on the matrix and extraction conditions (Galanakis, 2012). Enzyme-assisted extraction is also gaining popularity for its ability to selectively break down plant cell walls and release bound phenolic compounds.

Once extracted, by-products are commonly converted into shelf-stable forms such as dried powders, flours, or concentrated extracts. Drying methods, including hot-air drying, freeze-drying, and spray drying, are selected based on the sensitivity of the target compounds to heat and the desired final application. Freeze-drying, although more expensive, is preferred when preserving heat-labile antioxidants such as vitamin C and certain flavonoids is critical.




Figure 1. Conceptual Pathway for the Valorization of Food Processing By-products

Table 2. Functional Applications of Selected Food By-products in the Food Industry

By-product	Processed Form	Food Application	Functional Benefit
Citrus peel	Dried powder / pectin extract	Bakery products, jams, jellies, confectionery	Gelling, thickening and stabilizing properties; dietary-fibre enrichment
Banana peel	Flour / powder	Biscuits, muffins, bakery products	Dietary-fibre and mineral enrichment; modification of functional and textural properties
Tomato pomace	Dried powder / extract	Sauces, pasta, bakery products and other formulated foods	Lycopene and antioxidant enrichment; potential colour enhancement
Grape pomace	Flour / powder / polyphenol-rich extract	Bread, bakery products and meat products	Polyphenol and dietary-fibre enrichment; antioxidant activity and improved oxidative stability
Pomegranate peel	Aqueous / hydroalcoholic extract or powder	Beverages, dairy products and other functional foods	Antioxidant enrichment; potential antimicrobial activity

Consumer Perception and Market Trends

Growing consumer awareness of sustainability and clean-label products has created favourable market conditions for by-product-derived ingredients. Upcycled food products, those made using ingredients that would otherwise go to waste, are increasingly being marketed as premium,


Applications in the Food Industry

The functional properties of by-product-derived ingredients make them attractive additives across a wide range of food products. Citrus pectin extracted from peels is extensively used as a natural gelling and thickening agent in jams, jellies, and confectionery (Sharma et al., 2017). Banana peel flour is incorporated into bakery products and extruded snacks to enhance fiber and mineral content while also improving water-holding capacity in dough formulations. Tomato pomace, rich in lycopene, is being explored as a natural colorant and nutritional fortificant in sauces, soups, and pasta products.

Grape pomace extracts, owing to their high antioxidant capacity, are being incorporated into bread and processed meat products to extend shelf life and reduce lipid oxidation, offering a natural alternative to synthetic preservatives (Deng et al., 2011). Pomegranate peel extracts have shown promise as antimicrobial agents in beverages and dairy products, consistent with their reported role in extending the shelf life of minimally processed foods (Ayala-Zavala et al., 2010). Table 2 outlines some of the key functional applications of these by-products across different food categories.

environmentally conscious choices rather than inferior substitutes. This shift in perception, combined with regulatory support for food waste reduction in many countries, is encouraging both established food companies and startups to invest in by-product valorization technologies.

Figure 2. Dietary Fiber and Polyphenol Content of Selected Fruit and Vegetable By-products



Challenges and Future Outlook

Despite their promise, several challenges hinder the widespread commercial utilization of food by-products. Seasonal and geographical variability in raw material composition can result in inconsistent nutrient and bioactive content, complicating standardization. Microbial contamination risk during collection and storage of wet by-products necessitates rapid processing or preservation, which can add to operational costs. Additionally, regulatory frameworks governing the use of by-product-derived ingredients in food products are still evolving in many regions, creating uncertainty for manufacturers.

Looking ahead, continued research into cost-effective extraction technologies, coupled with supportive policy frameworks and growing consumer acceptance, is likely to accelerate the mainstream adoption of by-product valorization. Collaborative efforts between food processors, researchers, and policymakers will be essential to build efficient supply chains that connect by-product generators with ingredient manufacturers, closing the loop in a truly circular food system.

Conclusion

Food by-products, long regarded as waste, represent an underutilized source of nutrition and functional ingredients with significant potential to benefit both the food industry and

public health. Peels, pomace, and other residues are rich in dietary fiber, vitamins, and bioactive compounds that can be recovered through appropriate extraction and processing technologies and incorporated into a wide range of food products. Realizing the full potential of this approach will require continued innovation in processing technology, supportive regulation, and consumer education, ultimately transforming what was once discarded into ingredients that genuinely go "from peels to plates."

Acknowledgement

The authors gratefully acknowledge the support extended by the Amity Institute of Food Technology, Amity University, Noida, and the ICAR-Directorate of Cashew Research, Puttur, for facilitating this review.

References

- Ayala-Zavala J. F., Rosas-Dominguez C., Vega-Vega V. and Gonzalez-Aguilar G. A. (2010). Antioxidant enrichment and antimicrobial protection of fresh-cut fruits using their own by-products: Looking for integral exploitation. *Journal of Food Science*, 75 (8), R175–R181.
- Deng Q., Penner M. H. and Zhao Y. (2011). Chemical composition of dietary fiber and polyphenols of five different varieties of wine grape pomace skins. *Food Research International*, 44 (9), 2712–2720.
- Galanakis C. M. (2012). Recovery of high added-value components from food wastes: Conventional, emerging technologies and commercialized applications. *Trends in Food Science and Technology*, 26 (2), 68–87.
- Sharma K., Mahato N., Cho M. H. and Lee Y. R. (2017). Converting citrus wastes into value-added products: Economic and environmentally friendly approaches. *Nutrition*, 34, 29–46.
- Vichare S. A. and Morya S. (2024). Exploring waste utilization potential: Nutritional, functional and medicinal properties of oilseed cakes. *Frontiers in Food Science and Technology*, 4, 1441029.
