

# Beating the Heat: Managing Summer Stress in Dairy Cattle

Shweta Choudhary, Kalpana Jorasia, Priyanka Meena, Sandeep Choudhary

Post Graduate Institute of Veterinary Education and Research, Jaipur, Rajasthan, India

Corresponding Author: [vet.shwetakhicher@gmail.com](mailto:vet.shwetakhicher@gmail.com)

## Introduction

Dairy farming plays a vital role in the agricultural economy, providing livelihood, nutrition, and income to millions of farmers. Summer can be a challenging season for dairy farmers. When temperatures rise and humidity climbs, dairy cattle face a serious threat that often goes underestimated—heat stress. Unlike humans, cattle are poorly equipped to cool themselves efficiently. Their large body mass, limited sweat glands, and high metabolic heat production from rumen fermentation make them particularly vulnerable to warm weather.

Therefore, understanding the causes, effects, and management strategies of heat stress is essential for maintaining animal welfare, sustaining productivity, and ensuring economic profitability during the summer season.

## Understanding Heat Stress in Dairy Cattle

### What Is Heat Stress?

Heat stress occurs when a cow's heat load exceeds her ability to dissipate it to the surrounding environment. Cattle generate significant metabolic heat, especially high-producing dairy cows that consume large quantities of feed to support lactation. When ambient temperature and humidity prevent this heat from escaping, the animal's core body temperature rises, triggering a cascade of physiological responses.

### The Temperature-Humidity Index (THI)

The most commonly used metric to assess heat stress risk is the Temperature-Humidity Index (THI). This index combines ambient temperature and relative humidity into a single value that reflects the animal's thermal comfort level.

THI Value	Stress Level	Expected Impact
< 68	No stress	Normal production and behavior
68-71	Mild stress	Slight drop in milk yield, increased respiration rate
72-78	Moderate stress	Noticeable decline in milk production and feed intake
79-83	Severe stress	Significant production losses, reproductive issues
Above 84	Emergency	Risk of organ failure and death

It is important to note that high-producing dairy cows can begin showing signs of heat stress at THI values as low as 65 to 68, lower than the traditional threshold of 72. This is because these animals generate considerably more metabolic heat due to their higher feed intake and milk output.

### Effects of Heat Stress on Dairy Cattle

1. Reduced Milk Production
2. Decreased Feed Intake and Digestive Efficiency
3. Impaired Reproductive Performance

Heat stress has profound effects on fertility in both cows and bulls:

- Estrus detection rates decline
- Conception rates fall
- Embryo mortality increases
- Bull fertility declines

4. Compromised Immune Function

Heat-stressed cattle are more susceptible to:

- Mastitis
- Respiratory diseases
- Retained placenta and metritis
- Lameness

### Heat Stress Management Strategies

#### **1. Housing Management**

Proper housing is one of the most effective ways to reduce heat stress:

- Provide adequate shade using trees, asbestos sheets, or thatched roofs
- Ensure proper orientation of sheds (east-west direction) to minimize sun exposure
- Maintain good ventilation for air circulation
- Install fans, exhaust systems, or tunnel ventilation in intensive systems

#### **2. Cooling Systems**

Cooling methods help in reducing body temperature:

- Use sprinklers and misters to wet the animal's body
- Combine sprinkling with fans for evaporative cooling
- Provide wallows or showers in hot climates

- Avoid excessive moisture accumulation to prevent hygiene issues

### 3. Feeding Management

Nutritional adjustments can help animals cope with heat stress:

- Feed during cooler hours (early morning or late evening)
- Provide high-quality, easily digestible feed
- Increase energy density by adding fats instead of carbohydrates
- Supplement minerals and vitamins (especially electrolytes, sodium, potassium)
- Avoid feeding spoiled or fermented feed

### 4. Water Management

Water is crucial during heat stress:

- Provide unlimited access to clean and cool drinking water
- Increase the number of water points to avoid crowding
- Ensure water troughs are placed in shaded areas
- Regular cleaning of water troughs to maintain hygiene

### 5. Herd Management Practices

- Avoid overcrowding in sheds
- Minimize animal handling during peak heat hours
- Schedule milking, vaccination, and transportation during cooler times

- Regularly observe animals for early signs of heat stress

### 6. Breeding and Genetic Management

- Prefer heat-tolerant breeds or crossbreeds
- Plan breeding programs during cooler seasons
- Use reproductive technologies carefully during summer

#### Recent Approaches and Innovations

- Use of climate-controlled housing systems
- Precision dairy farming tools for monitoring temperature and animal behavior
- Nutritional supplements such as yeast cultures and antioxidants
- Genetic selection for heat tolerance

#### Conclusion

Heat stress is a critical challenge affecting dairy cattle during the summer months, leading to significant losses in milk production, reproductive efficiency, and overall health. However, its adverse effects can be effectively minimized through proper management practices, including improved housing, efficient cooling systems, balanced nutrition, and adequate water supply.

Adopting scientific and practical heat stress management strategies not only enhances animal comfort and welfare but also ensures sustainable dairy production and economic stability for farmers. As climate change continues to increase the frequency of extreme heat events, proactive and adaptive management will become even more essential in modern dairy farming.

\*\*\*\*\*