


# Common Neurologic Consults in Pregnancy



Mary Angela O'Neal, M.D.

Assistant Professor, Department of Neurology, Harvard Medical School


Director of the Women's Neurology Program

Chief of General Neurology




BRIGHAM AND  
WOMEN'S HOSPITAL

# Disclosures



- I have no disclosures

# Objectives




Use Cases to:

- Review the pathophysiology of some common neurologic disorders seen in pregnancy
- Discuss the importance of planning pregnancy
- Describe treatment specific concerns around pregnancy


**Migraine, Stroke, IIH, Low pressure headache, Postpartum Neuropathy, Epilepsy**

## Issues to discuss when caring for Women in their Reproductive years




- Family planning
- Discussion of medication risks in pregnancy
- Effects of pregnancy on the underlying disease
- Effects of the underlying disease on pregnancy

# FDA Pharmaceutical Pregnancy Categories




- A** Adequate and well controlled human studies have failed to demonstrate a risk to the fetus in the first trimester of pregnancy ( and there is no risk in later trimesters).
- B** Animal reproduction studies have failed to demonstrate a risk to the fetus and there are no adequate and well controlled studies in pregnant women OR Animal studies have shown an adverse effect, but adequate and well-controlled studies in pregnant women have failed to demonstrate a risk to the fetus in any trimester.
- C** Animal reproduction studies have shown an adverse effect on the fetus and there are no adequate and well-controlled studies in humans, but potential benefits may warrant use of the drug in pregnant women despite potential risks.
- D** There is positive evidence of human fetal risk based on adverse reaction data from investigational or marketing experience or studies in humans, but potential benefits may warrant use of the drug in pregnant women despite potential risks
- X** Studies in animals or humans have demonstrated fetal abnormalities and/or there is positive evidence of human fetal risk based on adverse reaction data from investigational or marketing experience, and the risks involved in use of the drug in pregnant women clearly outweigh potential benefits

# Case 1




- A 35 year-old woman G1 PO at 31 and 5/7 weeks of gestation woke with a severe headache. She began seeing visual spots, and a half hour later she completely lost her vision.
- Shortly thereafter, she developed the worst headache of her life and blacked out. At the outside hospital her blood pressure was 170/120. She was transferred to our hospital.

Which one of these etiologies is the most likely cause?



- A. Migraine without aura
- B. Subdural hematoma
- C. Migraine with aura
- D. Preeclampsia
- E. Idiopathic intracranial hypertension


Answer is D



- A. Migraine without aura
- B. Subdural hematoma
- C. Migraine with aura
- D. Preeclampsia**
- E. Idiopathic intracranial hypertension





# Red Flags



- Change in headache character or pattern
- Headaches with characteristics of elevated ICP
- New headaches
- Associated with elevated BP
- Unusually severe headache
- Abnormal neurologic exam
- Headaches associated with systemic disorders


Historical features	Headache type	Helpful radiographic studies
Thunderclap onset	SAH, RCVS, CVT	MRI,MRV,MRA, CT, CTA, CTV
Postural	PDPH, IIH	Brain MRI with gadolinium (gadolinium is not usually used in pregnancy)
First Trimester	Likely migraine or Tension type	None (except if the headaches have any red flags)
Prior similar headache	Benign	Not needed
Hypertension, proteinuria	PEE/Eclampsia	MRI

# Frequency and Headache Type by Trimester



O'Neal MA. Headaches complicating pregnancy and the postpartum period. Pract Neurol 2017; 17:191-202



# Pre-eclampsia/Eclampsia Definition



Diagnosed when a pregnant woman develops

- High blood pressure (two separate readings taken at least six hours apart of 140 or more in systolic blood pressure and/or 90 or more in diastolic blood pressure)
- 300 mg of protein in a 24-hour urine sample  
(proteinuria- no longer required)
- Onset of seizures and change in mental status defines eclampsia


# Pathogenesis of Eclampsia



Nature Reviews | Immunology

Sibai B, Dekkar G, Kupfermanc M. Pre-eclampsia. Lancet 2005; 365:785-799

# Pre-eclampsia/ Eclampsia




- PEE is associated with both significant maternal and fetal morbidity and mortality
- Maternal complications include abruption placentae, disseminated coagulopathy, acute renal failure, stroke, hemorrhage, death and long- term cardiovascular morbidity
- Fetal complications include premature delivery, low birth weight, hypoxic neurologic injury, and death.

Khan KS, Wojdyla D, Say L, Gulmezoglu AM, Van Look PFA. WHO analysis of causes of maternal death: systematic review. *Lancet* 2006; 367: 1066–1074.


Task force on hypertension in pregnancy. *Hypertension in Pregnancy* 2013. American College of Obstetricians and Gynecologists

In this patient what is the most appropriate imaging?



- A. Head CT without contrast
- B. Brain MRI without gadolinium
- C. Head CT without contrast and CTA
- D. Head CT without contrast, CTA and CTV
- E. Brain MRI without gadolinium, MRA and MRV

## Answer is D



- A. Head CT without contrast
- B. Brain MRI without gadolinium
- C. Head CT without contrast and CTA
- D. Head CT without contrast, CTA and CTV**
- E. Brain MRI without gadolinium, MRA and MRV



# Pathogenesis of PEE

Genetic factors

Maladaptation to placental implantation

Impaired remodeling of spiral arteries

Increased free radicals

Endothelial damage-  
Maternal hypertension

Increased levels of antiangiogenic factors

Kidney

Brain-  
Eclampsia


Liver

Proteinuria

Seizures, visual changes, vasogenic edema, infarction and ICH

Activation of the coagulation system → HELLP

# Axial Flair MRI-PRES



Hinchey J, Chaves C, Appignani B, Breen J, Pao L et al. A Reversible Posterior Leukoencephalopathy Syndrome. NEJM 1996; 334:494-500

# Treatment of Pre-eclampsia/Eclampsia

- Blood Pressure control




- Magnesium Sulfate

Altman D, et al. Do women with pre-eclampsia, and their babies, benefit from magnesium sulphate? The Magpie Trial: a randomized placebo-controlled trial. *Lancet* 2002, 359(9321):1877-1890.


Lucas et al. A comparison of magnesium sulfate and phenytoin for the prevention of eclampsia. *N Engl J Med* 1995;333:201-5.

# Case




- A 23 year-old women with migraine without aura is now 8 weeks pregnant. Her migraines had been well controlled with sumatriptan.
- She's now having her usual headaches with nausea and vomiting 2-3 times a week.

Which of these migraine medications is safest to use throughout pregnancy?




- A. Sumatriptan
- B. Topiramate
- C. Dihydroergotamine
- D. Naprosyn
- E. Lasmiditan

Answer is A



- A. **Sumatriptan**
- B. Topiramate
- C. Dihydroergotamine
- D. Naprosyn
- E. Lasmiditan


# Migraine Treatment around Pregnancy



- Planning
- Symptomatic therapy
- Other



# Other Therapies




- Physical, chiropractic and massage therapy
- Trigger Management
- Acupuncture
- Occipital nerve blocks





# Migraine during Pregnancy



- 60-70 % migraines undergo remission
- Small percent of new onset migraine during pregnancy
- Increased risk of preeclampsia/eclampsia

Kvisvik EV, et al. Headache and migraine during pregnancy and puerperium: the MIGRA-study. J Headache Pain Aug 2011; 12(4):443-451

Adeney KL, et al. Risk of preeclampsia in relation to maternal history of migraine headaches. J Matern Fetal Neonatal Med 2005; 18(3):167-172


# Symptomatic Therapies

Generic Name	Level of Risk in Pregnancy	Breastfeeding- Hale Lactation Rating
Acetaminophen	B	L1
NSAIDS	B (D in 3 <sup>rd</sup> trimester)	L1-L2
Metoclopramide	B	L2
Prochlorperazine	C	L3
Dihydroergotamine	X	L4
Magnesium	A (D)	L1
Triptans	C	L3
Ditans; Lasmiditan	No data in humans (adverse effects noted in animals)	No data
Gepants; Ubrogapant, Rimegepant	No data in humans (adverse effects noted in animals)	No data

# Preventative Medications

Drug Class	Generic Name	Level of Risk in Pregnancy	Breastfeeding
Beta- blockers	Atenolol Propranolol	D C (D at term)	Caution Compatible
Antiepileptics	Gabapentin Topiramate Valproate	C D X	Compatible Caution Caution
Tricyclics	Amitriptyline	C	Compatible
SNRIs	Duloxetine Venlafaxine	C C	Little data Little data
CGRP inhibitors	Erenumab Fremanezumab Galcanezumab	No data	No data
Gepants	Rimegepant Eptinezumab	No data	No data
Vitamins	Magnesium Coenzyme Q10	Safe	Safe

# Triptans and Pregnancy




- Data from the Sumatriptan/Naratriptan Pregnancy Registry
- Data from Norwegian Mother and Child Cohort

Ephross SA, Sinclair SM. Final results from the 16-year sumatriptan, naratriptan, and treximet pregnancy registry. *Headache* 2014;54(7):1158–1172.

-Spielmann K, Kayser A, Beck E, et al. Pregnancy outcome after anti-migraine triptan use: a prospective observational cohort study. *Cephalalgia* 2019;38(6):1181–1092

-Nezvalová-Henriksen K, Spigset O, et al. Triptan safety during pregnancy: a Norwegian population registry study. *European Journal of Epidemiology* 2013; 28(9): 759–769.

# Emergency Treatment of Migraine in Pregnancy



A sequential algorithm for migraine treatment:

- IVF- suggest NS at 20-30 mg/kg over 1-2 hours and 500 mg- 1 gm of IV magnesium sulfate
- Metoclopramide 10 mg IV or Prochlorperazine 5-10 mg IV
- Methylprednisolone 1 gm IV or 6 mg SQ Sumatriptan
- Analgesics

1. Cete Y, Dora B, Ertan C et al. A randomized prospective placebo-controlled study of intravenous magnesium sulfate vs. metoclopramide in the management of acute migraine attacks in the emergency department. *Cephalalgia* 2005; 25(3): 199-204.

2. Nezalova-Henriksen et al. Triptan exposure during pregnancy and the risk of major congenital malformations and adverse pregnancy outcomes: results from the Norwegian Mother and Child Cohort Study. *Headache* 2010; 50(4): 563-575.

# Migraine and Lactation


---

- Triptans considered safe
- Several Preventative drugs also safe



Lactmed


## Case



- A 28-year-old woman G2P1 at 30 weeks gestation was last seen well at 10 am. She was found on the ground not speaking or moving her right side at 10:50 am.
- On initial exam at noon, she was mute with left gaze deviation and dense right hemiplegia.

NIHSS\* was 15.

# Question




Which of the following statements is true of ischemic stroke during pregnancy and postpartum?

- a. Brain MRI, Brain and Neck MRA are the best imaging modalities
- b. IV tPA should not be used in pregnancy due to fetal risk
- c. Head CT and CTA of neck and brain are the most appropriate imaging modalities
- d. The causes of ischemic stroke during pregnancy and postpartum are similar to those that occur outside of pregnancy



# Answer




Which of the following statements is true of ischemic stroke in pregnancy and postpartum?

- a. Brain MRI, Brain and Neck MRA are the best imaging modalities
- b. IV tPA should not be used in pregnancy due to fetal risk
- c. Head CT and CTA of neck and brain are the most appropriate imaging modalities**
- d. The causes of ischemic stroke during pregnancy and postpartum are similar to those that occur outside of pregnancy


The evaluation and treatment of stroke during pregnancy should be, in most cases, the same as in the non-pregnant state

# How should you treat this lady?




- Imaging- Head CT vs Brain MRI, Vessel Imaging
- IV t-PA
- Thrombectomy


# Radiation Exposure




- **DETERMINISTIC EFFECTS**





# Radiation Exposure



- **STOCHASTIC EFFECTS**







## Case Reports of IV tPA in Pregnancy

Author/yr	# Cases	Gestational age	Maternal Outcome	Fetal Outcome
Dapprich/2002	1	12 weeks	Near complete	Healthy
Weatherby/2003	1	9 weeks	Full	Healthy
Leonhardt/2006	1	23 weeks	Full	Healthy
Wiese/2006	1	13 weeks	NIHSS-4	Healthy
Murugappan/2006	4	12 weeks 4 weeks 6 weeks 37 weeks	Healthy Healthy Death-complic of angio Healthy	Pregnancy terminated Pregnancy terminated Death Healthy
Yamaguchi/2010	1	18 weeks	Full	Healthy
Hori/2012	1	13 weeks	NIHSS-1	Healthy
Tassi/2013	1	16 weeks	NIHSS-1	Healthy
Ritter/2014	1	36 weeks	NIHSS-3	Healthy
Ritchie/2015	1	39 weeks	Full	Healthy
Tversky/2016	1	5 weeks	Full	Not available
Festa/2017	1	5 weeks	NIHSS-1	Healthy
Khan/2017	1	9 weeks	NIHSS-4	Fetal demise
Jiang/2018	1	31 weeks	Full	Healthy
Laudais/2018	1	13 weeks	NIHSS-1	Healthy


## Case Reports of Thrombectomy for Pregnant Women

Author/yr	# of Cases	Gestational Age	Maternal Outcome	Fetal Outcome
Aaron/2016	2	3 <sup>rd</sup> trimester 37 weeks	NIHSS-1 NIHSS-4	Healthy Healthy
Bhokal/2017	2	24 weeks 25 weeks	NIHSS-1 Full	Healthy Not available

Dicpinigaitis AJ, et al. Endovascular Thrombectomy for Treatment of Acute Ischemic Stroke During Pregnancy and the Early Postpartum Period. Stroke. 2021 Sep 20.




# Causes of Ischemic Stroke in Pregnancy



- Cardiac emboli
- Dissection
- Pre- eclampsia/ Eclampsia
- Coagulopathy
- Cerebral Venous Thrombosis
- Reversible Cerebral Vasoconstriction Syndrome
- Other

## Case




22 year-old lady 26 weeks pregnant who comes in for evaluation of headaches. She has gained 47 lbs. from her pre- pregnancy weight. She is currently having headaches mostly in the morning. She denies any visual symptoms.

She has bilateral disc edema on exam.




# What tests would you do next?




- A. Head CT
- B. Brain MRI with gadolinium
- C. Brain MRI and MRA
- D. Brain MRI and MRV without gadolinium

Answer is D




- A. Head CT
- B. Brain MRI with gadolinium
- C. Brain MRI and MRA
- D. Brain MRI and MRV without gadolinium**

# Idiopathic Intracranial Hypertension




- Worsening headache with recumbency
  - Improves while upright and the worst upon awakening from sleep
  - Behaves like tumor associated headache (pseudotumor cerebri)
- CSF pressure greater than 250 mm water
- Transient visual obscurations
  - Sudden loss of vision lasting less than 30 seconds involving one or both eyes, and are followed by full visual recovery.
- Vomiting, often projectile, tends to improve headaches
- Pulsatile tinnitus can occur in 52% of IIH patients.<sup>11</sup>

# Epidemiology of IIH



- Up 21/100,000
- 90% of patients are obese
- Female predominance 4:1

# IIH in Pregnancy




- Onset during pregnancy - 14 weeks
- Reactivation of IIH later second trimester- 20 weeks
- No increased obstetric complications
- Visual outcomes the same as in the non-pregnant patients

The Brigham Leadership Program - Cohort VI  
Group VI

Digre K et al, Pseudotumor cerebri and pregnancy, Neurology 1984;34:721-9.


# Symptoms of IIH



- Headache 92%
- Transient visual obscurations 72%
- Pulsatile Tinnitus 60%
- Photopsia
- Retrobulbar pain
- Diplopia
- Sustained visual loss 26%




# Evaluation



- Brain MRI/ MRV
- Lumbar Puncture
- Ophthalmological Evaluation





# Idiopathic Intracranial Hypertension




- **MRI Findings**
  - Prominent subarachnoid space around the optic
  - Papilledema
    - ✦ flattening of the posterior sclera
    - ✦ intraocular protrusion of the optic nerve head
  - Enhancement of the prelaminar (intra-ocular) optic nerves
  - Partial empty sella turcica
  - slitlike ventricles (relatively uncommon compared to other findings)
  - Acquired tonsillar ectopia (mimicking Chiari I malformation)
  - Increased subcutaneous fat thickness in the scalp and neck <sup>17</sup>

# MRI findings in IIH



# IIH Management Issues in Pregnancy



- Weight control
- Medications
- Serial lumbar punctures
- Surgery




## **The Use of Acetazolamide During Pregnancy in Intracranial Hypertension Patients**

Julie Falardeau, MD, Brenna M. Lobb, MS, MPH, Sara Golden, MPH,  
Steven D. Maxfield, BS, Emanuel Tanne, MD

Results: 101 women with IIH (158 pregnancies)  
acetazolamide usage before 13 week of gestation 50  
pregnancies

1. Similar risk of spontaneous abortion
2. No major complication in the offspring


# Management of Labor




- C-section – for Obstetric concerns
- Adequate labor analgesia – Uterine contractions increase ICP
- Neuroaxial anesthesia preferred




# Case



- 26 year old lady 2 days after a vaginal delivery with epidural anesthesia is seen for a holocephalic headache which is worse with sitting or standing and relieved with lying down.




# Which of these are risk factors for low pressure headache?



- A. Female sex
- B. Size of the needle
- C. Operator experience
- D. Low BMI
- E. All of the above



Answer is E



- A. Female sex
- B. Size of the needle
- C. Operator experience
- D. Low BMI
- E. All of the above**

# Risk Factors

## **Nonmodifiable**


1. Age
2. Sex
3. Low BMI
4. History prior postdural puncture headache
5. History of chronic headache

**vs**


## **Modifiable**

1. Needle shape
2. Bevel orientation and angle of insertion
3. Size of spinal needle
4. Stylet replacement
5. Operator experience

# Clinical Features




- Headache
- Nausea/Vomiting
- Neck Pain
- Tinnitus
- Visual changes
- Vertigo/Gait problems



# Pathophysiology


---

**Decrease in CSF pressure**




# Diffuse Meningeal Enhancement

---




# MRI findings in low pressure headache

---



# Treatment of Post-lumbar Puncture Headache



- Conservative
- Caffeine
- Epidural Blood Patch




# Controlled trials of medication in treatment of PDPH

Study	Patient Number	Regimen	Outcome
Camann 1990	40 Randomized	Caffeine 300 mg Assessed 4 & 24 hours	4 hours- 30% improvement
Sechzer 1978	41 Randomized	Caffeine 500mg IV given and repeated if headache persisted	1-2 hours 75% improvement Overall 70 % improvement
Wu 2018	126 Randomized	Aminophylline 250 mg IV	8 hours significant improvement in headache severity




# Case




- ▶ The neurology service was asked to evaluate a 32-year-old woman G1P1 for right leg weakness two days postpartum. She had an uneventful vaginal delivery of a 3266-g baby with epidural analgesia. She noted right leg numbness and knee weakness. On the first postpartum day, her leg buckled and she fell when she stood to move to the bathroom. She had no back pain or leg pain.
- ▶ On exam: 4/5 weakness in right hip flexion and knee extension, diminished right patellar DTR, and sensory loss in her medial thigh and calf. Her back was not tender and back range of motion was normal. She had some tenderness over the right inguinal ligament.

# What is the most likely diagnosis?




- A. Epidural hematoma
- B. A sciatic nerve injury
- C. A lateral femoral cutaneous neuropathy
- D. A femoral nerve injury
- E. Epidural abscess

Answer is D




- A. Epidural hematoma
- B. A sciatic nerve injury
- C. A lateral femoral cutaneous neuropathy
- D. A femoral nerve injury**
- E. Epidural abscess


# Femoral Neuropathy




- ▶ Incidence 2.8/100,000
- ▶ 25% are bilateral
- ▶ Findings
- ▶ Risk factors




# Peripheral Nerves of the leg



- Lateral Femoral Cutaneous Neuropathy
- Femoral Neuropathy
- Peroneal Neuropathy
- Lumbosacral Plexopathy
- Obturator nerve injury



## Case




- A 27 year-old right-handed woman with a genetic generalized epilepsy well controlled on valproic acid and levetiracetam plans to start a family and presents to clinic to optimize her preconception regimen.
- Past medical history significant for major depression with several suicidal attempts in the past and her epilepsy was refractory for a while for unclear reasons. Reportedly had a rash to lamotrigine and also failed levetiracetam monotherapy. Therefore, valproic acid was added and she was kept on dual therapy. She enjoyed five years of seizure freedom and is hesitant to change her regimen.

# Preconception Planning

1. Ensure **adequate contraception**:
  1. Some AEDs may reduce the efficacy of hormonal forms of birth control
  2. Consider possible impact of the contraception on the AED level
2. **Clarify the diagnosis**: non-epileptic versus epileptic; focal versus generalized; known etiology?; surgical candidate?
3. Optimize **AED regimen** for seizure control and pregnancy outcome:
  1. Consider switching to an AED with a better pregnancy outcome profile
  2. Monotherapy preferred to polytherapy
  3. Reduce to minimal effective dose
  4. If polytherapy is necessary, some AEDs are preferable (LTG, LEV) while some should be avoided (VPA, PB, TPM)
4. Recommend **folic acid (1–4 mg daily)**


## Balancing Act: Benefits versus Risks of AEDs during Pregnancy



- Teratogenic effects on offspring significant, with increased risk for ***major congenital malformations (MCMs)***
- ***Neurodevelopmental defects*** common, with lifelong consequences
- Risk for adverse ***Ob and Neonatal outcomes*** may also be higher
- The effect of ***uncontrolled seizures*** on the fetus



# Comparative safety of anti-epileptic drugs during pregnancy



% MCM for different AED monotherapies as revealed by different registries: NAAPR, North American AED Pregnancy Registry (Hernandez-Díaz et al, *Neurology* 2012); EURAP, an International Registry of Antiepileptic drugs and Pregnancy, UK Epilepsy and Pregnancy Register (Tomson et Battino, *Lancet Neurol* 2012 and *Seizure* 2015). The total sample size (n) included in each treatment is also presented; if n < 50, the information was not included.


## 2009 AAN/AES PP Update on Neurodevelopmental Outcomes

- Cognition is *probably not reduced* in children of *untreated WWE*.
- *CBZ probably* does *not* increase poor cognitive outcomes compared to unexposed controls.
- Monotherapy exposure to *VPA probably* reduces cognitive outcomes, and monotherapy exposure to *PHT or PB possibly* reduces cognitive outcomes.
- *AED polytherapy* exposure *probably* reduces cognitive outcomes as compared to AED monotherapy.



1. Gaily E, et al. *Neurology*. 2004;62(1):28-32.
2. Holmes LB, et al. *Teratology*. 2000;61(3):196-202.
3. Reinisch JM, et al. *JAMA*. 1995;274(19):1518-1525.
4. Adab N, et al. *J Neurol Neurosurg Psychiatry*. 2004;75(11):1575-1583.
5. Vinten J, et al. *Neurology*. 2005;64(6):949-954.
6. Adab N, et al. *J Neurol Neurosurg Psychiatry*. 2001;70(1):15-21.

# Preferential selection of AEDs for WWE of childbearing age





*“This is a teaching hospital”*