

WARNING

KEEP OUT OF REACH OF CHILDREN

DO NOT SWALLOW. The spray from this product may act as an irritant. Avoid inhalation and contact with the eyes and skin. Users are encouraged to seek further advice.

STORAGE:

DO NOT STORE product in direct sunlight.

DO NOT STORE product in temperature below 5°C



AfriKelp™ LG-1

(Ecklonia maxima)

PACKAGE WEIGHTS

VOLUME	NET WEIGHT	GROSS WEIGHT
1000 Litres	1080.0 kg	1180.0 kg
200 Litres	216.0 kg	227.0 kg
20 Litres	21.6 kg	23.6 kg

TYPICAL ANALYSIS

Effective Extracts		11 mg / L
Analysis:	w/v	(weight/volume)
Product:	pH	4.7
Specific Gravity:		1.08



Scan for
product SDS

Distributed by Advanced Nutrients Pty Ltd
Address: 27 Westbourne Drive, Wights Mountain, QLD 4520
Phone: 1800 244 009
Email: info@advancednutrients.com.au
Web: www.advancednutrients.com.au

This label complies with the National Code of Practice for fertiliser description and labelling.

Crop	Application	Rate	Water & Comments	Application	Rate	Water & Comments
Tomato	Foliar	2 L/ha	Foliar spray 14 days after plantation. Repeat 2-3 times every 14 days. <i>Apply first foliar spray at 3-4 leaf stage for directly sown plants.</i>	Dip	1:100	Dip seedlings directly before planting
Potato	Foliar	2 L/ha	Foliar spray at 3-4 leaf stage. Repeat after 2-3 weeks. <i>Do not apply after tuber initiation.</i>	Dip	1:300	Dip seed tubers directly before planting.
Vines	Drippers	1:500	Apply through drippers at start of new growth (for root stimulation and early growth)	Foliar	2L/ha	Foliar spray at flowering & repeat after 10-12 days. (1:300 with water)
Table Grapes	Drippers	1:500	Apply through drippers at start of new growth (for root stimulation and early growth)	Foliar	3 L/ha	Spray after full bloom and repeat 2 to 3 times at 10-12 day intervals
Banana	Drench	1:500	Soil drench 14 days after nematicide application to the base of the sucker and repeat after 6 months	Foliar	4 L/ha	Foliar spray at pre-flowering (5- 6 months pre harvest) repeat 3 times at monthly intervals
Cucurbits	Dip	1:100	Dip tray with seedlings in solution or wet seedling tray with watering can before transplant and spray 14 days later, repeat once or twice	Foliar	2-3 L/ha	14-21 day intervals. Spray at 3-4 leaf stage for directly seeded plants. Not recommended after flowering
Canola & Corn	Foliar	2 L/ha	Foliar spray at 3-4 leaf stage.	APPLICATION GUIDE		
Almonds	Foliar	3 L/ha	Spray at 50% bloom and repeat twice at 14 days intervals	Apply through boom spray, aerial, fertigation or overhead irrigation. DO NOT apply when conditions are extremely hot. Best applied in temperatures in the range of 5 - 25°C.		
Avocados	Dip	1:100	Dip roots of seedling before planting into nursery bag	MIXING INSTRUCTIONS		
	Foliar	3L/ha	1:200 Spray 4 weeks after planting in nursery bag. Mature trees Spray at pre-bloom, full bloom, fruit set and, optional, 14-21 days later.	1. Half fill tank to the required volume with water.		
	Drench	1:200	Drench plant before planting out.	2. Add the required amount of AfriKelp™ LG-1		
Fruit Trees	Dip	1:100	Dip bare roots for 5 minutes before transplant	3. Add appropriate amount of wetting agent as per manufacturer's instructions.		
	Soak seedling	1:200	Soak seedling bag before transplant	4. Fill the remainder of tank with water and maintain agitation.		
	Soak soil	1:300	Or soak soil around trees after plant-out or apply via irrigation system	COMPATIBILITY		
	Foliar	3 L/ha	Spray 3 to 5 times during early active growth at 21 day intervals. Deciduous: Spray at full bloom, fruit set & again 14 days later (14-21 days intervals) Evergreen Spray at pre-bloom, full bloom, fruit set and, optional, 14-21 days later (if water volume exceeds 1000 L/ha, increase AfriKelp rate to retain a 1:300 dilution).	Product may be compatible with herbicides and insecticides, please check with the manufacturer. Please be aware that physical compatibility does not guarantee product efficacy.		
Vegetables	Dip	1:100	Dip tray with seedlings in solution or wet with watering can before transplant and spray 14 days after transplant and repeat once	CONDITIONS OF SALE		
	Foliar	2-3 L/ha	14 to 21 day intervals. Directly seeded plants receive foliar spray at 3 to 5-leaf stage. Repeat once or twice at 14 to 21 day intervals	These goods are manufactured and tested to specific standards and are sold in good faith, however, no guarantee is given as conditions and methods under which they are used are beyond the control of Advanced Nutrients Pty Ltd. Before using, user should determine the suitability of the product for his or her intended use and user assumes the risk and liability in connection therewith.		