

## WARNING

KEEP OUT OF REACH OF CHILDREN

DO NOT SWALLOW. The spray from this product may act as an irritant. Avoid inhalation and contact with the eyes and skin. Users are encouraged to seek further advice.

## STORAGE:

DO NOT STORE product in direct sunlight.

DO NOT STORE product in temperature below 5°C



# Bang CalciN™

NPKS (12-0-0-0) + 16% Ca  
+ Organic Catalysts



Scan for  
product SDS

## GUARANTEED ANALYSIS

|             |    |            |       |         |
|-------------|----|------------|-------|---------|
| Nitrogen    | N  | as nitrate | 12.0% | 120 g/L |
| Calcium     | Ca | as nitrate | 16.0% | 160 g/L |
| Fulvic Acid | FA | as fulvate | 0.16% | 1.6 g/L |

|                   |     |                 |
|-------------------|-----|-----------------|
| Analysis:         | w/v | (weight/volume) |
| Product:          | pH  | 2.0 - 3.0       |
| Specific Gravity: |     | 1.41            |

Distributed by Advanced Nutrients Pty Ltd  
Address: 27 Westbourne Drive, Wights Mountain, QLD 4520  
Phone: 1800 244 009  
Email: [info@advancednutrients.com.au](mailto:info@advancednutrients.com.au)  
Web: [www.advancednutrients.com.au](http://www.advancednutrients.com.au)

This label complies with the National Code of Practice for fertiliser description and labelling.

# Bang CalciN™

NPKS (12-0-0-0) + 16% Ca, + Organic Catalysts

| BROADACRE                       | RATE L/Ha | WATER     | COMMENTS                                            |
|---------------------------------|-----------|-----------|-----------------------------------------------------|
| Cotton                          | 2 - 6     | 1:20 - 50 | Apply prior to flowering & again as boll fills      |
| Wheat                           | 4 - 7     | 1:20 - 50 | Apply prior to flowering                            |
| Pasture                         | 6 - 10    | 1:20 - 50 | Apply 3-4 weeks after emergence                     |
| Soil Application                | 30 - 70   |           |                                                     |
| VITICULTURE                     | RATE L/Ha | WATER     | COMMENTS                                            |
| Wine/Table Grapes               | 3 - 5     | 1:100     | Prior to flowering and at 2 weekly intervals        |
| Soil Application                | 20 - 50   |           | Apply 2-3 application from bud burst to flowering   |
| TREE CROPS                      | RATE L/Ha | WATER     | COMMENTS                                            |
| Stone/Pome Fruit                | 4 - 7     | 1:100     | Apply after petal fall & at 2 weekly intervals      |
| Citrus/Olives                   | 5 - 10    | 1:100     | Apply after flowering & at 2 weekly intervals       |
| Avocado                         | 5 - 10    | 1:100     | Apply prior to flowering and after fruit set        |
| Nut Trees                       | 5 - 10    | 1:100     | Apply at fruit development                          |
| Soil Application                | 20 - 60   |           | Apply 2-4 application from flowering to harvest     |
| HORTICULTURE                    | RATE L/Ha | WATER     | COMMENTS                                            |
| Brassica                        | 5 - 7     | 1:100     | Apply before head formation & at 2 weekly intervals |
| Carrots                         | 3 - 5     | 1:100     | Apply after seedling and vegetative stages          |
| Lettuce                         | 3 - 5     | 1:100     | Apply prior to head formation                       |
| Potatoes                        | 4 - 7     | 1:100     | Apply after flowering & at 2 weekly intervals       |
| Strawberries/<br>Tomatoes/Beans | 3 - 5     | 1:100     | Apply after flowering & at 2 weekly intervals       |
| Soil Application                | 30 - 50   |           | Apply 2-4 application from flowering to harvest     |

## APPLICATION GUIDE

Apply through boom spray, aerial, fertigation or overhead irrigation. DONOT apply when conditions are extremely hot. Best applied in temperatures in the range of 5 - 25°C.

## MIXING INSTRUCTIONS

1. Half fill tank to the required volume with water.
2. Add the required amount of Bang CalciN™.
3. Add appropriate amount of wetting agent as per manufacturer's instructions.
4. Fill the remainder of tank with water and maintain agitation.

## PACKAGE WEIGHTS

| VOLUME      | NET WEIGHT | GROSS WEIGHT |
|-------------|------------|--------------|
| 1000 Litres | 1410.0 kg  | 1510.0 kg    |
| 200 Litres  | 282.0 kg   | 294.0 kg     |
| 20 Litres   | 28.2 kg    | 30.2 kg      |

## CONDITIONS OF SALE

These goods are manufactured and tested to specific standards and are sold in good faith, however, no guarantee is given as conditions and methods under which they are used are beyond the control of Advanced Nutrients Pty Ltd. Before using, user should determine the suitability of the product for his or her intended use and user assumes the risk and liability in connection therewith.

## COMPATIBILITY

Product may be compatible with herbicides and insecticides, please check with the manufacturer. Please be aware that physical compatibility does not guarantee product efficacy.

**Office:** 27 Westbourne Drive, Wights Mountain, QLD 4520

**Phone:** 1800 244 009

**Email:** info@advancednutrients.com.au

**Web:** www.advancednutrients.com.au