

Beck Anxiety Inventory

Below is a list of common symptoms of anxiety. Please carefully read each item in the list. Indicate how much you have been bothered by that symptom during the past month, including today, by circling the number in the corresponding space in the column next to each symptom.

	Not At All	Mildly but it didn't bother me much.	Moderately - it wasn't pleasant at times	Severely – it bothered me a lot
Numbness or tingling	0	1	2	3
Feeling hot	0	1	2	3
Wobbliness in legs	0	1	2	3
Unable to relax	0	1	2	3
Fear of worst happening	0	1	2	3
Dizzy or lightheaded	0	1	2	3
Heart pounding/racing	0	1	2	3
Unsteady	0	1	2	3
Terrified or afraid	0	1	2	3
Nervous	0	1	2	3
Feeling of choking	0	1	2	3
Hands trembling	0	1	2	3
Shaky / unsteady	0	1	2	3
Fear of losing control	0	1	2	3
Difficulty in breathing	0	1	2	3
Fear of dying	0	1	2	3
Scared	0	1	2	3
Indigestion	0	1	2	3
Faint / lightheaded	0	1	2	3
Face flushed	0	1	2	3
Hot/cold sweats	0	1	2	3
Column Sum				

Scoring - Sum each column. Then sum the column totals to achieve a grand score. Write that score here _____.

Interpretation

A grand sum between **0 – 21** indicates very low anxiety. That is usually a good thing. However, it is possible that you might be unrealistic in either your assessment which would be denial or that you have learned to “mask” the symptoms commonly associated with anxiety. Too little “anxiety” could indicate that you are detached from yourself, others, or your environment.

A grand sum between **22 – 35** indicates moderate anxiety. Your body is trying to tell you something. Look for patterns as to when and why you experience the symptoms described above. For example, if it occurs prior to public speaking and your job requires a lot of presentations you may want to find ways to calm yourself before speaking or let others do some of the presentations. You may have some conflict issues that need to be resolved. Clearly, it is not “panic” time but you want to find ways to manage the stress you feel.

A grand sum that **exceeds 36** is a potential cause for concern. Again, look for patterns or times when you tend to feel the symptoms you have circled. Persistent and high anxiety is not a sign of personal weakness or failure. It is, however, something that needs to be proactively treated or there could be significant impacts to you mentally and physically. You may want to consult a physician or counselor if the feelings persist.

Adult ADHD Self-Report Scale (ASRS-v1.1) Symptom Checklist

Please answer the questions below, rating yourself on each of the criteria shown using the scale on the right side of the page. As you answer each question, place an X in the box that best describes how you have felt and conducted yourself over the past 6 months.		Never	Rarely	Sometimes	Often	Very Often
1.	How often do you have trouble wrapping up the final details of a project, once the challenging parts have been done?					
2.	How often do you have difficulty getting things in order when you have to do a task that requires organization?					
3.	How often do you have problems remembering appointments or obligations?					
4.	When you have a task that requires a lot of thought, how often do you avoid or delay getting started?					
5.	How often do you fidget or squirm with your hands or feet when you have to sit down for a long time?					
6.	How often do you feel overly active and compelled to do things, like you were driven by a motor?					

Part A

7.	How often do you make careless mistakes when you have to work on a boring or difficult project?					
8.	How often do you have difficulty keeping your attention when you are doing boring or repetitive work?					
9.	How often do you have difficulty concentrating on what people say to you, even when they are speaking to you directly?					
10.	How often do you misplace or have difficulty finding things at home or at work?					
11.	How often are you distracted by activity or noise around you?					
12.	How often do you leave your seat in meetings or other situations in which you are expected to remain seated?					
13.	How often do you feel restless or fidgety?					
14.	How often do you have difficulty unwinding and relaxing when you have time to yourself?					
15.	How often do you find yourself talking too much when you are in social situations?					
16.	When you're in a conversation, how often do you find yourself finishing the sentences of the people you are talking to, before they can finish them themselves?					
17.	How often do you have difficulty waiting your turn in situations when turn taking is required?					
18.	How often do you interrupt others when they are busy?					

Part B

How old were you when these problems first began to occur? _____

Patient Health Questionnaire (PHQ-9)

Patient Name: _____

Date: _____

	Not at all	Several days	More than half the days	Nearly every day
1. Over the <i>last 2 weeks</i>, how often have you been bothered by any of the following problems?				
a. Little interest or pleasure in doing things	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. Feeling down, depressed, or hopeless	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c. Trouble falling/staying asleep, sleeping too much	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
d. Feeling tired or having little energy	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
e. Poor appetite or overeating	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
f. Feeling bad about yourself or that you are a failure or have let yourself or your family down	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
g. Trouble concentrating on things, such as reading the newspaper or watching television.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
h. Moving or speaking so slowly that other people could have noticed. Or the opposite; being so fidgety or restless that you have been moving around a lot more than usual.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
i. Thoughts that you would be better off dead or of hurting yourself in some way.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
2. If you checked off any problem on this questionnaire so far, how difficult have these problems made it for you to do your work, take care of things at home, or get along with other people?				
	Not difficult at all	Somewhat difficult	Very difficult	Extremely difficult
	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>