

## Integral Calculus and Its Applications

1. Reduction formulae. 2. Reduction formulae for  $\int \sin^n x \, dx$ ,  $\int \cos^n x \, dx$  and evaluation of  $\int_0^{\pi/2} \sin^n x \, dx$ ,  $\int_0^{\pi/2} \cos^n x \, dx$ . 3. Reduction formula for  $\int \sin^m x \cos^n x \, dx$  and evaluation of  $\int_0^{\pi/2} \sin^m x \cos^n x \, dx$ . 4. Reduction formulae for  $\int \tan^n x \, dx$ ,  $\int \cot^n x \, dx$ . 5. Reduction formulae for  $\int \sec^n x \, dx$ ,  $\int \operatorname{cosec}^n x \, dx$ . 6. Reduction formulae for  $\int x^n e^{ax} \, dx$ ,  $\int x^m (\log x)^n \, dx$ . 7. Reduction formulae for  $\int x^n \sin mx \, dx$ ,  $\int x^n \cos nx \, dx$  and  $\int \cos^m x \sin nx \, dx$ . 8. Definite integrals. 9. Integral as the limit of a sum. 10. Areas of curves. 11. Lengths of curves. 12. Volumes of revolution. 13. Surface areas of revolution. 14. Objective Type of Questions.

### 6.1 REDUCTION FORMULAE

The reader is already familiar with some standard methods of integrating functions of a single variable. However, there are some integrals which cannot be evaluated by the afore-said methods. In such cases, the method of reduction formulae proves useful. A reduction formula connects an integral with another of the same type but of lower order. The successive application of the reduction formula enables us to evaluate the given integral. Now we shall derive some standard reduction formulae.

#### 6.2 (1) REDUCTION FORMULAE for

$$(a) \int \sin^n x \, dx \quad (b) \int \cos^n x \, dx.$$

$$(a) \int \sin^n x \, dx = \int \sin^{n-1} x \cdot \sin x \, dx \quad [\text{Integrated by parts}]$$

$$\begin{aligned} &= \sin^{n-1} x \cdot (-\cos x) - \int (n-1) \sin^{n-2} x \cos x (-\cos x) \, dx \\ &= -\sin^{n-1} x \cos x + (n-1) \int \sin^{n-2} x (1 - \sin^2 x) \, dx \\ &= -\sin^{n-1} x \cos x + (n-1) \int \sin^{n-2} x \, dx - (n-1) \int \sin^n x \, dx \end{aligned}$$

Transposing

$$n \int \sin^n x \, dx = -\sin^{n-1} x \cos x + (n-1) \int \sin^{n-2} x \, dx$$

$$\text{or} \quad \int \sin^n x \, dx = -\frac{\sin^{n-1} x \cos x}{n} + \frac{n-1}{n} \int \sin^{n-2} x \, dx \quad \dots(i)$$

(b) Similarly, 
$$\int \cos^n x \, dx = \frac{\sin x \cos^{n-1} x}{n} + \frac{n-1}{n} \int \cos^{n-2} x \, dx$$

Thus we have the required reduction formulae.

**Obs. To integrate  $\int \sin^n x \, dx$  or  $\int \cos^n x \, dx$ ,**

(a) when the index of  $\sin x$  is odd put  $\cos x = t$

when the index of  $\cos x$  is odd, put  $\sin x = t$

(b) when the index is an even positive integer, express the integrand as a series of cosines of multiple angles and integrate term by term if  $n$  is small, otherwise use the method of reduction formulae.

(2) To show that 
$$\int_0^{\pi/2} \sin^n x \, dx = \int_0^{\pi/2} \cos^n x \, dx$$
  

$$= \frac{(n-1)(n-3)(n-5)\dots}{n(n-2)(n-4)\dots} \times \left(\frac{\pi}{2}, \text{ only if } n \text{ is even}\right)$$

From (i), we have

$$I_n = \int_0^{\pi/2} \sin^n x \, dx = - \left| \frac{\sin^{n-1} x \cos x}{n} \right|_0^{\pi/2} + \frac{n-1}{n} \int_0^{\pi/2} \sin^{n-2} x \, dx$$

i.e.

$$I_n = \frac{n-1}{n} I_{n-2}$$

**Case I. When  $n$  is odd**

Similarly 
$$I_{n-2} = \frac{n-3}{n-2} I_{n-4}, \quad I_{n-4} = \frac{n-5}{n-4} I_{n-6}$$

$$I_5 = \frac{4}{5} I_3, \quad I_3 = \frac{2}{3} I_1 = \frac{2}{3} \int_0^{\pi/2} \sin x \, dx = \frac{2}{3} \left| -\cos x \right|_0^{\pi/2} = \frac{2}{3}.$$

Form these, we get 
$$I_n = \frac{(n-1)(n-3)(n-5)\dots 2}{n(n-2)(n-4)\dots 3} \dots (ii)$$

**Case II. When  $n$  is even**

We have 
$$I_{n-2} = \frac{n-3}{n-2} I_{n-4}, \quad I_{n-4} = \frac{n-5}{n-4} I_{n-6}$$

$$I_4 = \frac{3}{4} I_2, \quad I_2 = \frac{1}{2} I_0 = \int_0^{\pi/2} \sin^0 x \, dx = \frac{1}{2} \int_0^{\pi/2} dx = \frac{1}{2} \cdot \frac{\pi}{2}.$$

Form these, we obtain 
$$I_n = \frac{(n-1)(n-3)(n-5)\dots 3 \cdot 1}{n(n-2)(n-4)\dots 4 \cdot 2} \cdot \frac{\pi}{2} \dots (iii)$$

Combining (ii) and (iii), we get the required result for  $\int_0^{\pi/2} \sin^n x \, dx$ .

Proceeding exactly as above, we get the result for  $\int_0^{\pi/2} \cos^n x \, dx$ .

**Example 6.1.** Integrate (i)  $\int \sin^4 x \, dx$  (ii)  $\int_0^{\pi/2} \cos^6 x \, dx$ .

**Solution.** (i) We have the reduction formula

$$\int \sin^n x \, dx = \frac{\sin^{n-1} x \cos x}{n} + \frac{n-1}{n} \int \sin^{n-2} x \, dx$$

Putting  $n = 4, 2$  successively,

$$\int \sin^4 x \, dx = - \frac{\sin^3 x \cos x}{4} + \frac{3}{4} \int \sin^2 x \, dx \dots (\alpha)$$

$$\int \sin^2 x \, dx = -\frac{\sin x \cos x}{2} + \frac{1}{2} \int (\sin x)^0 \, dx$$

But  $\int (\sin x)^0 \, dx = \int dx = x. \quad \therefore \int \sin^2 x \, dx = -\frac{\sin x \cos x}{2} + \frac{x}{2}$

Substituting this in (α), we get

$$\int \sin^4 x \, dx = -\frac{\sin^3 x \cos x}{4} + \frac{3}{4} \left( -\frac{\sin x \cos x}{2} + \frac{x}{2} \right)$$

(ii) We know that  $\int_0^{\pi/2} \cos^n x \, dx = \frac{(n-1)(n-3)(n-5)\dots}{n(n-2)(n-4)\dots} \left( \frac{\pi}{2} \text{ if } n \text{ is even} \right)$

Putting  $n = 6$ , we get

$$\int_0^{\pi/2} \cos^6 x \, dx = \frac{5.3.1}{6.4.2} \frac{\pi}{2} = \frac{5\pi}{16}.$$

**Example 6.2.** Evaluate

(i)  $\int_0^a \frac{x^7 \, dx}{\sqrt{(a^2 - x^2)}}$  (V.T.U., 2006) (ii)  $\int_0^{\pi} \frac{\sqrt{(1 - \cos x)}}{1 + \cos x} \sin^2 x \, dx$  (iii)  $\int_0^{\infty} \frac{dx}{(a^2 + x^2)^n}.$

**Solution.** (i)  $\int_0^a \frac{x^7}{\sqrt{(a^2 - x^2)}} \, dx$  Put  $x = a \sin \theta$ , so that  $dx = a \cos \theta \, d\theta$   
Also when  $x = 0$ ,  $\theta = 0$ , when  $x = a$ ,  $\theta = \pi/2$

$$= \int_0^{\pi/2} \frac{a^7 \sin^7 \theta}{a \cos \theta} \cdot a \cos \theta \, d\theta = a^7 \int_0^{\pi/2} \sin^7 \theta \, d\theta = a^7 \cdot \frac{6.4.2}{7.5.3.1} = \frac{16}{35} a^7$$

(ii) Putting  $x = 2\theta$ , we get

$$\begin{aligned} \int_0^{\pi} \frac{\sqrt{(1 - \cos x)}}{1 + \cos x} \sin^2 x \, dx &= \int_0^{\pi/2} \frac{\sqrt{(1 - \cos 2\theta)}}{1 + \cos 2\theta} \sin^2 2\theta \cdot 2 \, d\theta \\ &= 2 \int_0^{\pi/2} \frac{\sqrt{2} \sin \theta}{2 \cos^2 \theta} \cdot (2 \sin \theta \cos \theta)^2 \, d\theta = 4\sqrt{2} \int_0^{\pi/2} \sin^3 \theta \, d\theta = 4\sqrt{2} \cdot \frac{2}{3} = \frac{8\sqrt{2}}{3}. \end{aligned}$$

(iii)  $\int_0^{\infty} \frac{dx}{(a^2 + x^2)^n}$  Put  $x = a \tan \theta$ , so that  $dx = a \sec^2 \theta \, d\theta$   
Also when  $x = 0$ ,  $\theta = 0$ , when  $x = \infty$ ,  $\theta = \pi/2$

$$= \int_0^{\pi/2} \frac{a \sec^2 \theta \, d\theta}{a^{2n} \sec^{2n} \theta} = \frac{1}{a^{2n-1}} \int_0^{\pi/2} \cos^{2n-2} \theta \, d\theta = \frac{1}{a^{2n-1}} \cdot \frac{(2n-3)(2n-5)\dots 3.1}{(2n-2)(2n-4)\dots 4.2} \cdot \frac{\pi}{2}.$$

**Example 6.3.** Evaluate  $\int_0^a \frac{x^n}{\sqrt{(a^2 - x^2)}} \, dx$ . Hence find the value of  $\int_0^1 x^n \sin^{-1} x \, dx$ .

**Solution.** Putting  $x = a \sin \theta$ , we get

$$\begin{aligned} \int_0^a \frac{x^n}{\sqrt{(a^2 - x^2)}} \, dx &= \int_0^{\pi/2} \frac{(a \sin \theta)^n}{a \cos \theta} (a \cos \theta) \, d\theta = a^n \int_0^{\pi/2} \sin^n \theta \, d\theta \\ &= \frac{(n-1)(n-3)\dots 2}{n(n-2)\dots 3} a^n, \text{ if } n \text{ is odd} \\ &= \frac{(n-1)(n-3)\dots 1}{n(n-2)\dots 2} \cdot \frac{\pi}{2} a^n, \text{ if } n \text{ is even} \end{aligned} \quad \dots(i)$$

Now integrating by parts, we have

$$\int_0^1 x^n \sin^{-1} x \, dx = \left| (\sin^{-1} x) \cdot \frac{x^{n+1}}{n+1} \right|_0^1 - \int_0^1 \frac{x^{n+1}}{n+1} \cdot \frac{1}{\sqrt{(1-x^2)}} \, dx$$



$$\begin{aligned}
 &= \frac{1}{(n+1)} \left[ \frac{\pi}{2} - \int_0^1 \frac{x^{n+1}}{(1-x^2)} \right] && \text{[Using (i) p. 241]} \\
 &= \frac{1}{n+1} \left\{ \frac{\pi}{2} - \frac{n(n-2)(n-4) \dots 1}{(n+1)(n-1)(n-3) \dots 2} \frac{\pi}{2} \right\} && \text{when } n \text{ is odd} \\
 &= \frac{1}{n+1} \left\{ \frac{\pi}{2} - \frac{n(n-2)(n-4) \dots 2}{(n+1)(n-1)(n-3) \dots 3} \right\} && \text{when } n \text{ is even}
 \end{aligned}$$

**Evaluate 6.4.** Evaluate  $I_n = \int_0^a (a^2 - x^2)^n dx$  where  $n$  is a positive integer. Hence show that

$$I_n = \frac{2n}{2n+1} a^2 I_{n-1}$$

**Solution.** Putting  $n = a \sin \theta$ , we get

$$\begin{aligned}
 I_n &= \int_0^a (a^2 - x^2)^n dx = \int_0^{\pi/2} (a^2 - a^2 \sin^2 \theta)^n a \cos \theta d\theta = a^{2n+1} \int_0^{\pi/2} \cos^{2n+1} \theta d\theta \\
 &= a^{2n+1} \cdot \frac{(2n)(2n-2)(2n-4) \dots 4 \cdot 2}{(2n+1)(2n-1)(2n-3) \dots 5 \cdot 3} && [\because (2n+1) \text{ is always odd}]
 \end{aligned}$$

Now replacing  $n$  by  $n-1$ , we get

$$I_{n-1} = a^{2n-1} \frac{(2n-2)(2n-4) \dots 4 \cdot 2}{(2n-1)(2n-3) \dots 5 \cdot 3} \quad \therefore \quad \frac{I_n}{I_{n-1}} = a^2 \cdot \frac{2n}{2n+1} \quad \text{or} \quad I_n = \frac{2n}{2n+1} a^2 I_{n-1}$$

which is the second desired result.

### 6.3 (1) REDUCTION FORMULAE for $\int \sin^m x \cos^n x dx$

$$\int \sin^m x \cos^n x dx = \int \sin^{m-1} x \cdot \cos^n x \cdot \sin x dx \quad \text{[Integrate by parts]}$$

$$\begin{aligned}
 &= \sin^{m-1} x \cdot \left( -\frac{\cos^{n+1} x}{n+1} \right) - \int (m-1) \sin^{m-2} x \cos x \cdot \left( -\frac{\cos^{n+1} x}{n+1} \right) dx \\
 &= -\frac{\sin^{m-1} x \cos^{n+1} x}{n+1} + \frac{m-1}{n+1} \int \sin^{m-2} x (1 - \sin^2 x) \cos^n x dx \\
 &= -\frac{\sin^{m-1} x \cos^{n+1} x}{n+1} + \frac{m-1}{n+1} \int \sin^{m-2} x \cos^n x dx - \frac{m-1}{n+1} \int \sin^m x \cos^n x dx
 \end{aligned}$$

Transposing the last term to the left and dividing by  $1 + (m-1)/(n+1)$ , i.e.,  $(m+n)/(n+1)$ , we obtain the reduction formula

$$\int \sin^m x \cos^n x dx = -\frac{\sin^{m-1} x \cos^{n+1} x}{m+n} + \frac{m-1}{m+n} \int \sin^{m-2} x \cos^n x dx \quad \dots(1)$$

**Obs. To integrate  $\int \sin^m x \cos^n x dx$ ,**

(a) when  $m$  is odd, put  $\cos x = t$

when  $n$  is odd, put  $\sin x = t$

(b) when  $m$  and  $n$  both are even integers, express the integrand as a series of cosines of multiple angles and integrate term by term if  $m$  and  $n$  are small, otherwise use the method of reduction formulae.

**(2) To show that**

$$\int_0^{\pi/2} \sin^m x \cos^n x dx = \frac{(m-1)(m-3) \dots \times (n-1)(n-3) \dots}{(m+n)(m+n-2)(m+n-4) \dots} \times \left( \frac{\pi}{2} \right), \text{ only if both } m \text{ and } n \text{ are even}$$



From (i), we have

$$I_{m,n} = \int_0^{\pi/2} \sin^m x \cos^n x dx = \left| -\frac{\sin^{m-1} x \cos^{n+1} x}{m+n} \right|_0^{\pi/2} + \frac{m-1}{m+n} \int_0^{\pi/2} \sin^{m-2} x \cos^n x dx$$

i.e., 
$$I_{m,n} = \frac{m-1}{m+n} I_{m-2,n}$$

**Case I. When  $m$  is odd**

Similarly, 
$$I_{m-2,n} = \frac{m-3}{m+n-2} I_{m-4,n}, I_{m-4,n} = \frac{m-5}{m+n-4} I_{m-6,n}$$

.....

$$I_{5,n} = \frac{4}{n+5} I_{3,n}$$

Finally 
$$I_{3,n} = \frac{2}{n+3} I_{1,n} = \frac{2}{n+3} \int_0^{\pi/2} \sin x \cos^n x dx$$
  

$$= \frac{2}{n+3} \left| -\frac{\cos^{n+1} x}{n+1} \right|_0^{\pi/2} = \frac{2}{(n+3)(n+1)} \quad \dots(ii)$$

From these, we obtain

$$I_{m,n} = \frac{(m-1)(m-3)(m-5) \dots 4.2}{(m+n)(m+n-2)(m+n-4) \dots (n+3)(n+1)}$$

**Case II. When  $m$  is even**

We have, 
$$I_{m-2,n} = \frac{m-3}{m+n-2} I_{m-4,n}, I_{m-4,n} = \frac{m-5}{m+n-4} I_{m-6,n}$$

.....

$$I_{4,n} = \frac{3}{n+4} I_{2,n}, I_{2,n} = \frac{1}{n+2} I_{0,n} = \frac{1}{n+2} \int_0^{\pi/2} \cos^n x dx$$

From these, we have 
$$I_{m,n} = \frac{(m-1)(m-3)(m-5) \dots 1}{(m+n)(m+n-2)(m+n-4) \dots (n+2)} \int_0^{\pi/2} \cos^n x dx$$
  

$$= \frac{(m-1)(m-3) \dots 1}{(m+n)(m+n-2) \dots (n+2)} \cdot \frac{(n-1)(n-3) \dots 1}{n(n-2) \dots} \times (\pi/2 \text{ only if } n \text{ is even}) \quad \dots(iii)$$

Combining (ii) and (iii), we get the desired result.

**Example 6.5.** Integrate (i)  $\int \sin^4 x \cos^2 x dx$  (Raipur, 2005)

(ii)  $\int_0^\infty \frac{t^6}{(1+t^2)^7} dt$  (iii)  $\int_0^\infty \frac{x^2}{(1+x^2)^{7/2}} dx$  (V.T.U., 2010 S)

**Solution.** (i) Taking  $n = 2$ , in (i) of page 241, we have the reduction formula :

$$\int \sin^m x \cos^2 x dx = \frac{\sin^{m-1} x \cos^3 x}{m+2} + \frac{m-1}{m+2} \int \sin^{m-2} x \cos^2 x dx$$

Putting  $m = 4, 2$  successively,

$$\int \sin^4 x \cos^2 x dx = -\frac{\sin^3 x \cos^3 x}{6} + \frac{3}{6} \int \sin^2 x \cos^2 x dx \quad \dots(1)$$

$$\int \sin^2 x \cos^2 x dx = -\frac{\sin x \cos^3 x}{4} + \frac{1}{4} \int \cos^2 x dx$$

But 
$$\int \cos^2 x dx = \frac{1}{2} \int (1 + \cos 2x) dx = \frac{1}{2} \left( x + \frac{1}{2} \sin 2x \right)$$

$$\therefore \int \sin^2 x \cos^2 x \, dx = -\frac{\sin x \cos^3 x}{4} + \frac{1}{16} (2x + \sin 2x)$$

Substituting this in (1), we get

$$\int \sin^4 x \cos^2 x \, dx = -\frac{\sin^3 x \cos^3 x}{6} + \frac{1}{2} \left\{ -\frac{\sin x \cos^3 x}{4} + \frac{1}{16} (2x + \sin 2x) \right\}$$

(ii) Putting  $t = \tan \theta$ , so that

$$\int_0^\infty \frac{t^6}{(1+t^2)^7} dt = \int_0^{\pi/2} \frac{\tan^6 \theta}{\sec^{14} \theta} \sec^2 \theta \, d\theta = \int_0^{\pi/2} \sin^6 \theta \cos^6 \theta \, d\theta = \frac{5 \cdot 3 \cdot 1 \times 5 \cdot 3 \cdot 1}{12 \cdot 10 \cdot 8 \cdot 6 \cdot 4 \cdot 2} \times \frac{\pi}{2} = \frac{5\pi}{2048}.$$

(iii) Putting  $x = \tan \theta$ , so that

$$\int_0^\infty \frac{x^2}{(1+x^2)^{7/2}} dx = \int_0^{\pi/2} \frac{\tan^2 \theta}{\sec^7 \theta} \sec^2 \theta \, d\theta = \int_0^{\pi/2} \sin^2 \theta \cos^3 \theta \, d\theta = \frac{1.2}{53.1} = \frac{2}{15}.$$

**Example 6.6.** Evaluate : (i)  $\int_0^{\pi/6} \cos^4 3\theta \sin^3 6\theta \, d\theta$  (V.T.U., 2003 S)

(ii)  $\int_0^1 x^4 (1-x^2)^{3/2} dx$  (iii)  $\int_0^{2a} x^2 \sqrt{(2ax-x^2)} \, dx$ . (V.T.U., 2010)

**Solution.** (i)  $\int_0^{\pi/6} \cos^4 3\theta \sin^3 6\theta \, d\theta = \int_0^{\pi/6} \cos^4 3\theta (2 \sin 3\theta \cos 3\theta)^3 \, d\theta$

$$= 8 \int_0^{\pi/6} \sin^3 3\theta \cos^7 3\theta \, d\theta$$

$$= \frac{8}{3} \int_0^{\pi/2} \sin^3 x \cos^7 x \, dx$$

$$= \frac{8}{3} \cdot \frac{2 \times 6 \cdot 4 \cdot 2}{10 \cdot 8 \cdot 6 \cdot 4 \cdot 2} = \frac{1}{15}.$$

Put  $3\theta = x$   
so that  $3d\theta = dx$

Also when  $\theta = 0, x = 0$ ;  
when  $\theta = \pi/6, x = \pi/2$ .

(ii)  $\int_0^1 x^4 (1-x^2)^{3/2} dx$

$$= \int_0^{\pi/2} \sin^4 t (\cos^2 t)^{3/2} \cdot \cos t \, dt = \int_0^{\pi/2} \sin^4 t \cos^4 t \, dt$$

$$= \frac{3 \cdot 1 \times 3 \cdot 1}{8 \cdot 6 \cdot 4 \cdot 2} \cdot \frac{\pi}{2} = \frac{3\pi}{256}.$$

Put  $x = \sin t$  so that  $dx = \cos t \, dt$   
When  $x = 0, t = 0$ ; when  $x = 1, t = \pi/2$

(iii)  $\int_0^{2a} x^2 \sqrt{(2ax-x^2)} \, dx$

$$= \int_0^{\pi/2} x^{5/2} \sqrt{(2a-x)} \, dx$$

$$= \int_0^{\pi/2} (2a \sin^2 \theta)^{5/2} \sqrt{(2a)} \cos \theta \cdot 4a \sin \theta \cos \theta \, d\theta$$

$$= 2^5 a^4 \int_0^{\pi/2} \sin^6 \theta \cos^2 \theta \, d\theta = 32 a^4 \cdot \frac{5 \cdot 3 \cdot 1 \times 1}{8 \cdot 6 \cdot 4 \cdot 2} \cdot \frac{\pi}{2} = \frac{5\pi a^4}{8}.$$

Put  $x = 2a \sin^2 \theta$   
 $\therefore dx = 4a \sin \theta \cos \theta \, d\theta$

### PROBLEMS 6.1

Evaluate :

1. (i)  $\int_0^{\pi/2} \cos^9 x \, dx$

(ii)  $\int_0^{\pi/6} \sin^5 3\theta \, d\theta$

2. (i)  $\int_0^1 \frac{x^9}{\sqrt{(1-x^2)}} \, dx$

(ii)  $\int_0^1 x^5 \sin^{-1} x \, dx$

3. (i)  $\int_0^\infty \frac{dx}{(1+x^2)^n} (n > 1)$  (V.T.U., 2008 S)

(ii)  $\int_0^{\pi/4} \sin^2 x \cos^4 x \, dx$ . (J.N.T.U., 2003)

4. If  $I_{m,n} = \int_0^{\pi/2} \sin^m x \cos^n x dx$  ( $m > 0, n > 0$ ), show that  $I_{m,n} = \frac{m-1}{m+n} I_{m-2,n}$ .

Hence evaluate  $\int_0^{\pi/2} \sin^4 x \cos^8 x dx$

Evaluate :

5. (i)  $\int_0^{\pi/2} \sin^4 x \cos^6 x dx$  (Cochin, 2005)

(ii)  $\int_0^{\pi/2} \sin^{15} x \cos^3 x dx$

6. (i)  $\int_0^1 x^6 \sqrt{1-x^2} dx$

(ii)  $\int_0^{\pi/2} \cos^4 3\theta \sin^3 6\theta d\theta$

7. (i)  $\int_0^{2a} x^{7/2} (2a-x)^{-1/2} dx$

(ii)  $\int_0^{2a} \frac{x^3 dx}{\sqrt{(2ax-x^2)}}$  (Madras, 2000 S)

8. (i)  $\int_0^2 x^{5/2} \sqrt{2-x} dx$

(ii)  $\int_0^4 x^3 \sqrt{4x-x^2} dx$  (V.T.U., 2004)

9. If  $I_n = \int x^n \sqrt{a-x} dx$ , prove that  $(2n+3) I_n = 2an I_{n-1} - 2x^n (a-x)^{3/2}$  (Marathwada, 2008)

10. If  $n$  is a positive integer, show that  $\int_0^{2a} x^n \sqrt{(2ax-x^2)} dx = \frac{2n+1}{(n+2)n!} \cdot \frac{a^{n+2}}{2n} \pi$  (V.T.U., 2007)

#### 6.4 REDUCTION FORMULAE for (a) $\int \tan^n x dx$ (b) $\int \cot^n x dx$

(a) Let 
$$I_n = \int \tan^n x dx = \int \tan^{n-2} x \cdot \tan^2 x dx = \int \tan^{n-2} x \cdot (\sec^2 x - 1) dx$$
$$= \int \tan^{n-2} x \sec^2 x dx - \int \tan^{n-2} x dx$$

Thus, 
$$I_n = \frac{\tan^{n-1} x}{n-1} - I_{n-2}$$
 which is the required reduction formula.

(b) Let 
$$I_n = \int \cot^n x dx = \int \cot^{n-2} x \cot^2 x dx = \int \cot^{n-2} x (\operatorname{cosec}^2 x - 1) dx$$
$$= \int \cot^{n-2} x \operatorname{cosec}^2 x dx - \int \cot^{n-2} x dx$$

Thus 
$$I_n = -\frac{\cot^{n-1} x}{n-1} - I_{n-2}$$

which is the required reduction formula.

**Example 6.7.** Evaluate (i)  $\int \tan^5 x dx$  (ii)  $\int \cot^6 x dx$ .

**Solution.** (i) Putting  $n = 5, 3$  successively in the reduction formula for  $\int \tan^n x dx$ , we get

$$I_5 = \frac{1}{4} \tan^4 x - I_3; \quad I_3 = \frac{1}{2} \tan^2 x - I_1$$

Thus 
$$I_5 = \frac{1}{4} \tan^4 x - \frac{1}{2} \tan^2 x + I_1$$

i.e., 
$$\int \tan^5 x dx = \frac{1}{4} \tan^4 x - \frac{1}{2} \tan^2 x + \int \tan x dx = \frac{1}{4} \tan^4 x - \frac{1}{2} \tan^2 x - \log \cos x.$$

(ii) Putting  $n = 6, 4, 2$  successively in the reduction formula for  $\int \cot^n x dx$ , we get

$$I_6 = -\frac{1}{5} \cot^5 x - I_4; \quad I_4 = -\frac{1}{3} \cot^3 x - I_2; \quad I_2 = -\cot x - I_0$$

Thus 
$$I_6 = -\frac{1}{5} \cot^5 x + \frac{1}{3} \cot^3 x - \cot x - \int dx$$



i.e., 
$$\int \cot^6 x \, dx = -\frac{1}{5} \cot^5 x + \frac{1}{3} \cot^3 x - \cot x - x.$$

**Example 6.8.** If  $I_n = \int_0^{\pi/4} \tan^n \theta \, d\theta$ , prove that  $n(I_{n-1} + I_{n+1}) = 1$ .

(V.T.U., 2003)

**Solution.** The reduction formula for  $\int_0^{\pi/4} \tan^n \theta \, d\theta$  is

$$I_n = \frac{1}{n-1} \left| \tan^n x \right|_0^{\pi/4} - I_{n-2} = \frac{1}{n-1} - I_{n-2} \quad \text{or} \quad I_n + I_{n-2} = \frac{1}{n-1}$$

Changing  $n$  to  $n+1$ , we obtain

$$I_{n+1} + I_{n-1} = \frac{1}{(n+1)} \quad \text{or} \quad (n+1)(I_{n+1} + I_{n-1}) = 1.$$

### 6.5 REDUCTION FORMULAE for (a) $\int \sec^n x \, dx$ (b) $\int \operatorname{cosec}^n x \, dx$

(a) Let 
$$I_n = \int \sec^n x \, dx = \int \sec^{n-2} x \cdot \sec^2 x \, dx$$

Integrating by parts, we have

$$\begin{aligned} I_n &= \sec^{n-2} x \cdot \tan x - \int [(n-2) \sec^{n-3} x \cdot \sec x \tan x] \tan x \, dx \\ &= \sec^{n-2} x \tan x - (n-2) \int \sec^{n-2} x \cdot \tan^2 x \, dx \\ &= \sec^{n-2} x \tan x - (n-2) \int \sec^{n-2} x \cdot (\sec^2 x - 1) \, dx \\ &= \sec^{n-2} x \tan x - (n-2) I_n + (n-2) I_{n-2} \end{aligned}$$

Transposing, we have

$$(n-1)I_n = \sec^{n-2} x \tan x + (n-2)I_{n-2}$$

Thus 
$$I_n = \frac{\sec^{n-2} x \tan x}{n-1} + \frac{n-2}{n-1} I_{n-2} \text{ which is the desired reduction formula.}$$

(b) Let 
$$I_n = \int \operatorname{cosec}^n x \, dx = \int \operatorname{cosec}^{n-2} x \cdot \operatorname{cosec}^2 x \, dx$$

Integrating by parts, we have

$$\begin{aligned} I_n &= \operatorname{cosec}^{n-2} x \cdot (-\cot x) - \int [(n-2) \operatorname{cosec}^{n-3} x \cdot (-\operatorname{cosec} x \cot x) \cdot (-\cot x) \, dx \\ &= -\cot x \operatorname{cosec}^{n-2} x - (n-2) \int \operatorname{cosec}^{n-2} x (\operatorname{cosec}^2 x - 1) \, dx \\ &= -\cot x \operatorname{cosec}^{n-2} x - (n-2) I_n + (n-2) I_{n-2} \end{aligned}$$

or 
$$[1 + (n-2)]I_n = -\cot x \operatorname{cosec}^{n-2} x + (n-2)I_{n-2}$$

Thus 
$$I_n = -\frac{\cot x \operatorname{cosec}^{n-2} x}{n-1} + \frac{n-2}{n-1} I_{n-2}$$

which is the required reduction formula.

**Example 6.9.** Evaluate (i)  $\int_0^{\pi/4} \sec^4 x \, dx$  (ii)  $\int_{\pi/3}^{\pi/2} \operatorname{cosec}^3 \theta \, d\theta$ .

(V.T.U., 2008)

**Solution.** (i) Putting  $n = 4$  in the reduction formula for  $\int \sec^n x \, dx$ , we get  $I_4 = \frac{\sec^2 x \tan x}{3} + \frac{2}{3} I_2$

$$\begin{aligned} \therefore \int_0^{\pi/4} \sec^4 x \, dx &= \left| \frac{\sec^2 x \tan x}{3} \right|_0^{\pi/4} + \frac{2}{3} \int_0^{\pi/4} \sec^2 x \, dx \\ &= \frac{2}{3} + \frac{2}{3} \left| \tan x \right|_0^{\pi/4} = \frac{2}{3} (1+1) = 4/3. \end{aligned}$$

(ii) Putting  $n = 3$  in the reduction formula for  $\int \operatorname{cosec}^n x \, dx$ , we get

$$\begin{aligned} I_3 &= -\frac{1}{2} \cot x \operatorname{cosec} x + \frac{1}{2} I_1 \\ \therefore \int_{\pi/3}^{\pi/2} \operatorname{cosec}^3 x \, dx &= -\frac{1}{2} \left[ \cot x \operatorname{cosec} x \right]_{\pi/3}^{\pi/2} + \frac{1}{2} \int_{\pi/3}^{\pi/2} \operatorname{cosec} x \, dx \\ &= -\frac{1}{2} \left( 0 - \frac{2}{3} \right) + \frac{1}{2} \left[ \log (\operatorname{cosec} x - \cot x) \right]_{\pi/3}^{\pi/2} \\ &= \frac{1}{3} + \frac{1}{2} \left[ \log 1 - \log \left( \frac{2}{\sqrt{3}} - \frac{1}{\sqrt{3}} \right) \right] = \frac{1}{3} + \frac{1}{4} \log 3. \end{aligned}$$

### PROBLEMS 6.2

- Evaluate (i)  $\int \tan^6 x \, dx$  (V.T.U., 2007) (ii)  $\int \cot^5 x \, dx$ .
- Show that  $\int_0^{\pi/4} \tan^7 x \, dx = \frac{1}{12} (5 - 6 \log 2)$
- If  $I_n = \int_0^{\pi/4} \tan^n x \, dx$ , prove that  $(n-1)(I_n + I_{n-2}) = 1$ . (V.T.U., 2009)  
Hence evaluate  $I_5$ . (Madras, 2000)
- If  $I_n = \int_{\pi/4}^{\pi/2} \cot^n \theta \, d\theta$  ( $n > 2$ ), prove that  $I_n = \frac{1}{n-1} - I_{n-1}$ . Hence evaluate  $I_4$ . (Marathwada, 2008)
- Obtain the reduction formula for  $\int_0^{\pi/4} \sec^n \theta \, d\theta$ . (V.T.U., 2010 S)
- Evaluate (i)  $\int \sec^6 \theta \, d\theta$  (ii)  $\int_{\pi/6}^{\pi/2} \operatorname{cosec}^5 d\theta$ . 7. Evaluate  $\int_0^a (a^2 + x^2)^{5/2} \, dx$ .
- If  $I_n = \int \frac{t^n}{1+t^2} \, dt$ , show that  $I_{n+2} = \frac{t^{n+1}}{n+1} - I_n$ . Hence evaluate  $I_6$ .

### 6.6 REDUCTION FORMULAE for

(a)  $\int x^n e^{ax} \, dx$

(b)  $\int x^m (\log x)^n \, dx$ .

(a) Let  $I_n = \int x^n e^{ax} \, dx$

Integrating by parts, we have

$$I_n = x^n \cdot \frac{e^{ax}}{a} - \int nx^{n-1} \cdot \frac{e^{ax}}{a} \, dx$$

or  $I_n = \frac{x^n e^{ax}}{a} - \frac{n}{a} I_{n-1}$  which is the required reduction formula. (Madras, 2006)

(b) Let  $I_{m,n} = \int x^m (\log x)^n \, dx = \int (\log x)^n \cdot x^m \, dx$

Integrating by parts, we have

$$\begin{aligned} I_{m,n} &= (\log x)^n \cdot \frac{x^{m+1}}{m+1} - \int n(\log x)^{n-1} \cdot \frac{1}{x} \cdot \frac{x^{m+1}}{m+1} \, dx \\ &= \frac{x^{m+1}}{m+1} (\log x)^n - \frac{n}{m+1} \int x^m (\log x)^{n-1} \, dx \quad \text{or} \quad I_{m,n} = \frac{x^{m+1}}{m+1} (\log x)^n - \frac{n}{m+1} I_{m,n-1} \end{aligned}$$

which is the desired reduction formula.

### 6.7 REDUCTION FORMULAE for

(a)  $\int x^n \sin mx \, dx$

(b)  $\int x^n \cos mx \, dx$

(c)  $\int \cos^m x \sin nx \, dx$

(a) Let  $I_n = \int x^n \sin mx \, dx$

Integrating by parts, we get

$$\begin{aligned} I_n &= x^n \left( \frac{-\cos mx}{m} \right) - \int n x^{n-1} \left( \frac{-\cos mx}{m} \right) dx \\ &= -\frac{x^n \cos mx}{m} + \frac{n}{m} \int x^{n-1} \cos mx \, dx \quad [\text{Again integrate by parts}] \\ &= -\frac{x^n \cos mx}{m} + \frac{n}{m} \left\{ x^{n-1} \cdot \frac{\sin mx}{m} - \left\{ \int (n-1)x^{n-2} \cdot \frac{\sin mx}{m} dx \right\} \right\} \end{aligned}$$

or 
$$I_n = -\frac{x^n \cos mx}{m} + \frac{n}{m^2} x^{n-1} \sin mx - \frac{n(n-1)}{m^2} I_{n-2}$$

which is the desired reduction formula.

(Madras, 2003)

(b) Let  $I_n = \int x^n \cos mx \, dx$

Integrating twice by parts as above, we get

$$I_n = \frac{x^n \sin mx}{m} + \frac{n}{m^2} x^{n-1} \cos mx - \frac{n(n-1)}{m^2} I_{n-2}$$

(c) Let  $I_{m,n} = \int \cos^m x \sin nx \, dx$

Integrating by parts,

$$\begin{aligned} I_{m,n} &= -\cos^m x \cdot \frac{\cos nx}{n} - \int m \cos^{m-1} x (-\sin x) \cdot \left( \frac{-\cos nx}{n} \right) dx \\ &= -\frac{1}{n} \cos^m x \cos nx - \frac{m}{n} \int \cos^{m-1} x \cdot \cos nx \sin x \, dx \\ & \quad \left[ \because \sin(n-1)x = \sin nx \cos x - \cos nx \sin x \right. \\ & \quad \left. \text{or } \cos nx \sin x = \sin nx \cos x - \sin(n-1)x \right] \\ &= -\frac{1}{n} \cos^m x \cos nx - \frac{m}{n} \int \cos^{m-1} x \{ \sin nx \cos x - \sin(n-1)x \} dx \\ &= -\frac{1}{n} \cos^m x \cos nx - \frac{m}{n} (I_{m,n} - I_{m-1,n-1}) \end{aligned}$$

Transposing, we get

$$\left( 1 + \frac{m}{n} \right) I_{m,n} = -\frac{1}{n} \cos^m x \cos nx + \frac{m}{n} I_{m-1,n-1}$$

or 
$$I_{m,n} = -\frac{\cos^m x \cos nx}{m+n} + \frac{m}{m+n} I_{m-1,n-1}$$

which is the desired reduction formula.

**Example 6.10.** Show that  $\int_0^{\pi/2} \cos^m x \cos nx \, dx = \frac{m}{m+n} \int_0^{\pi/2} \cos^{m-1} x \cos(n-1)x \, dx$

Hence deduce that  $\int_0^{\pi/2} \cos^n x \cos nx \, dx = \frac{\pi}{2^{n+1}}$ . (S.V.T.U., 2008)

**Solution.** Let  $I_{m,n} = \int_0^{\pi/2} \cos^m x \cdot \cos nx \, dx$

Integrating by parts

$$I_{m,n} = \left[ \cos^m x \cdot \frac{\sin nx}{n} \right]_0^{\pi/2} - \int_0^{\pi/2} m \cos^{m-1} x (-\sin x) \times \frac{\sin nx}{n} dx$$



$$\begin{aligned}
 &= \frac{m}{n} \int_0^{\pi/2} \cos^{m-1} x \sin nx \sin x \, dx \quad \left[ \begin{array}{l} \because \cos(n-1)x = \cos nx \cos x + \sin nx \sin x \\ \text{or } \sin nx \sin x = \cos(n-1)x - \cos nx \cos x \end{array} \right] \\
 &= \frac{m}{n} \int_0^{\pi/2} \cos^{m-1} x [\cos(n-1)x - \cos nx \cos x] \, dx = \frac{m}{n} (I_{m-1, n-1} - I_{m, n})
 \end{aligned}$$

Transposing and dividing by  $(1 + m/n)$ , we get

$$I_{m, n} = \frac{m}{m+n} I_{m-1, n-1}$$

which is the required result.

Putting  $m = n$ ,  $I_n \left( = \int_0^{\pi/2} \cos^n x \cos nx \, dx \right) = \frac{1}{2} I_{n-1}$

Changing  $n$  to  $n-1$ ,

$$I_{n-1} = \frac{1}{2} I_{n-2}$$

$$\therefore I_n = \frac{1}{2} \left( \frac{1}{2} I_{n-2} \right) = \frac{1}{2^2} I_{n-2} = \frac{1}{2^3} I_{n-3} \dots = \frac{1}{2^n} I_{n-n} = \frac{1}{2^n} \cdot \int_0^{\pi/2} (\cos x)^0 \, dx$$

Hence  $I_n = \frac{1}{2^n} \cdot \frac{\pi}{2} = \frac{\pi}{2^{n+1}}$ .

**Example 6.11.** Find a reduction formula for  $\int e^{ax} \sin x \, dx$ . Hence evaluate  $\int e^x \sin^3 x \, dx$ .

**Solution.** Let  $I_n = \int e^{ax} \sin^n x \, dx = \int \frac{\sin^n x}{I} \cdot \frac{e^{ax}}{II} \, dx$

Integrating by parts,

$$\begin{aligned}
 I_n &= \sin^n x \cdot \frac{e^{ax}}{a} - \int (n \sin^{n-1} x \cos x) \cdot \frac{e^{ax}}{a} \, dx \\
 &= \frac{e^{ax} \sin^n x}{a} - \frac{n}{a} \int (\sin^{n-1} x \cos x) \cdot e^{ax} \, dx \quad [\text{Again integrating by parts}] \\
 &= \frac{e^{ax} \sin^n x}{a} - \frac{n}{a} \left[ \sin^{n-1} x \cos x \cdot \frac{e^{ax}}{a} - \int \{(n-1) \sin^{n-2} x \right. \\
 &\quad \left. \times \cos x \cdot \cos x + \sin^{n-1} x (-\sin x)\} \frac{e^{ax}}{a} \, dx \right] \\
 &= \frac{e^{ax} \sin^{n-1} x}{a^2} (a \sin x - n \cos x) + \frac{n}{a^2} \int [(n-1) \sin^{n-2} x \times (1 - \sin^2 x) - \sin^n x] e^{ax} \, dx \\
 &= \frac{e^{ax} \sin^{n-1} x}{a} (a \sin x - n \cos x) + \frac{n(n-1)}{a^2} I_{n-2} - \frac{n^2}{a^2} I_n
 \end{aligned}$$

Transposing and dividing by  $(1 + n^2/a^2)$ , we get

$$I_n = \frac{e^{ax} \sin^{n-1} x (a \sin x - n \cos x)}{a^2 + n^2} + \frac{n(n-1)}{a^2 + n^2} I_{n-2}$$

which is the required reduction formula.

Putting  $a = 1$  and  $n = 3$ , we get

$$I_3 = \frac{e^x \sin^2 x (\sin x - 3 \cos x)}{1^2 + 9} + \frac{3 \cdot 2}{1^2 + 9} I_1$$

But  $I_1 = \int e^x \sin x \, dx = \frac{e^x}{\sqrt{2}} \sin(x - \tan^{-1} 1)$ .

$$\therefore I_3 = \frac{e^x \sin^2 x (\sin x - 3 \cos x)}{10} + \frac{3}{5} \cdot \frac{e^x}{\sqrt{2}} \sin(x - \pi/4).$$

PROBLEMS 6.3

1. If  $I_n = \int x^n e^x dx$ , show that  $I_n + n I_{n-1} = x^n e^x$ . Hence find  $I_4$ . (Madras, 2000)
2. If  $u_n = \int_0^a x^n e^{-x} dx$ , prove that  $u_n - (n + a) u_{n-1} + a(n-1) u_{n-2} = 0$ . (Madras, 2003)
3. Obtain a reduction formula for  $\int x^m (\log x)^n dx$ . Hence evaluate  $\int_0^1 x^5 (\log x)^3 dx$ . (S.V.T.U., 2009; Bhillai, 2005)
4. If  $n$  is a positive integer, show that  $\int_0^1 x^m (\log x)^n dx = \frac{(-1)^n n!}{(m+1)^{n+1}}$ ,  $m > -1$ .
5. If  $I_n = \int_0^{\pi/2} x \sin^n x dx$  ( $n > 1$ ), prove that  $n^2 I_n = n(n-1) I_{n-2} + 1$ . Hence evaluate  $I_5$ .
6. If  $I_n = \int_0^{\pi/2} x \cos^n x dx$  ( $n > 1$ ), prove that  $I_n = \frac{n-1}{n} I_{n-2} - \frac{1}{n^2}$ . Hence evaluate  $I_4$ .
7. If  $u_n = \int_0^{\pi/2} x^n \sin x dx$ , ( $n > 1$ ), prove that  $u_n + n(n-1) u_{n-2} = n(\pi/2)^{n-1}$ . Hence evaluate  $u_2$ . (Madras, 2000 S)
8. If  $I_n = \int x^n \sin ax dx$ , show that  $a^2 I_n = -ax^n \cos ax + nx^{n-1} \sin ax - n(n-1) I_{n-2}$ . (Marathwada, 2008)
9. Prove that  $\int_0^{\pi/2} \cos^{n-2} x \sin nx dx = \frac{1}{n-1}$ ,  $n > 1$ .
10. If  $I_{m,n} = \int_0^{\pi/2} \cos^m x \cos nx dx$ , prove that  $I_{m,n} = \frac{m(m-1)}{m^2 - n^2} I_{m-2,n}$ .
11. Find a reduction formula for  $\int e^{ax} \cos^n x dx$ . Hence evaluate  $\int_0^{\pi/2} e^{2x} \cos^3 x dx$ .
12. Obtain a reduction formula for  $I_m = \int_0^\infty e^{-x} \sin^m x dx$  where  $m \geq 2$  in the form  $(1 + m^2) I_m = m(m-1) I_{m-2}$ . Hence evaluate  $I_4$ . (Gorakhpur, 1999)

6.8 DEFINITE INTEGRALS

**Property I.**  $\int_a^b f(x) dx = \int_a^b f(t) dt$

(i.e., the value of a definite integral depends on the limits and not on the variable of integration).

Let  $\int f(x) dx = \phi(x)$ ;  $\therefore \int_a^b f(x) dx = \phi(b) - \phi(a)$ .

Then  $\int f(t) dt = \phi(t)$ ;  $\therefore \int_a^b f(t) dt = \phi(b) - \phi(a)$ .

Hence the result.

**Property II.**  $\int_a^b f(x) dx = - \int_b^a f(x) dx$

(i.e., the interchange of limits changes the sign of the integral).

Let  $\int f(x) dx = \phi(x)$ ;  $\therefore \int_a^b f(x) dx = \phi(b) - \phi(a)$

and  $-\int_b^a f(x) dx = -[\phi(x)]_b^a = -[\phi(a) - \phi(b)] = \phi(b) - \phi(a)$ .

Hence the result.

**Property III.**  $\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$

Let  $\int f(x) dx = \phi(x)$ , so that  $\int_a^b f(x) dx = \phi(b) - \phi(a)$  ... (1)

Also  $\int_a^c f(x) dx + \int_c^b f(x) dx = [\phi(x)]_a^c + [\phi(x)]_c^b$   
 $= [\phi(c) - \phi(a)] + [\phi(b) - \phi(c)] = \phi(b) - \phi(a)$  ... (2)

Hence the result follows from (1) and (2).

**Property IV.**  $\int_0^a f(x) dx = \int_0^a f(a-x) dx$

Put  $x = a - t$ , so that  $dx = -dt$ . Also when  $x = 0$ ,  $t = a$ ; when  $x = a$ ,  $t = 0$ .

$\therefore \int_0^a f(x) dx = - \int_a^0 f(a-t) dt = \int_0^a f(a-t) dt = \int_0^a f(a-x) dx$  [Prop. II]

**Example 6.12.** Evaluate  $\int_0^{\pi/2} \frac{\sqrt{(\sin x)}}{\sqrt{(\sin x)} + \sqrt{(\cos x)}} dx$ .

**Solution.** Let  $I = \int_0^{\pi/2} \frac{\sqrt{(\sin x)}}{\sqrt{(\sin x)} + \sqrt{(\cos x)}} dx$

Then  $I = \int_0^{\pi/2} \frac{\sqrt{\left[\sin\left(\frac{1}{2}\pi - x\right)\right]}}{\sqrt{\left[\sin\left(\frac{1}{2}\pi - x\right)\right]} + \sqrt{\left[\cos\left(\frac{1}{2}\pi - x\right)\right]}} dx$  [Prop. IV]  
 $= \int_0^{\pi/2} \frac{\sqrt{(\cos x)}}{\sqrt{(\cos x)} + \sqrt{(\sin x)}} dx$

Adding  $2I = \int_0^{\pi/2} \frac{\sqrt{(\sin x)} + \sqrt{(\cos x)}}{\sqrt{(\sin x)} + \sqrt{(\cos x)}} dx = \int_0^{\pi/2} dx = \left| x \right|_0^{\pi/2} = \frac{\pi}{2}$ .

Hence  $I = \pi/4$ .

**Example 6.13.** Evaluate  $\int_0^1 \frac{\log(1+x)}{1+x^2} dx$ . (Cochin, 2005)

**Solution.** Let  $I = \int_0^1 \frac{\log(1+x)}{1+x^2} dx$  [ Put  $x = \tan \theta$  so that  $dx = \sec^2 \theta d\theta$   
When  $x = 0$ ,  $\theta = 0$ ; when  $x = 1$ ,  $\theta = \pi/4$  ]  
 $= \int_0^{\pi/4} \frac{\log(1+\tan \theta)}{1+\tan^2 \theta} \cdot \sec^2 \theta d\theta = \int_0^{\pi/4} \log(1+\tan \theta) d\theta$   
 $= \int_0^{\pi/4} \log\left[1+\tan\left(\frac{\pi}{4} - \theta\right)\right] d\theta = \int_0^{\pi/4} \log\left(1 + \frac{1-\tan \theta}{1+\tan \theta}\right) d\theta$  [Prop. IV]  
 $= \int_0^{\pi/4} \log\left(\frac{2}{1+\tan \theta}\right) d\theta = \log 2 \int_0^{\pi/4} d\theta - I$

Transposing,  $2I = \log 2 \cdot \left| \theta \right|_0^{\pi/4} = \frac{\pi}{4} \log 2$ . Hence  $I = \frac{\pi}{8} \log 2$ .

**Example 6.14.** Evaluate  $\int_0^{\pi} \frac{x \sin^3 x}{1+\cos^2 x} dx$ . (Madras, 2006)

**Solution.** Let  $I = \int_0^{\pi} \frac{x \sin^3 x}{1+\cos^2 x} dx$



Then 
$$I = \int_0^{\pi} \frac{(\pi - x) \sin^3 (\pi - x)}{1 + \cos^2 (\pi - x)} dx \quad [\text{Prop. IV}]$$

$$= \int_0^{\pi} \frac{(\pi - x) \sin^3 x}{1 + \cos^2 x} dx = \pi \int_0^{\pi} \frac{\sin^3 x}{1 + \cos^2 x} dx - I$$

Transposing, 
$$2I = \pi \int_0^{\pi} \frac{\sin^3 x}{1 + \cos^2 x} dx$$

$$= -\pi \int_1^{-1} (1 - t^2) \frac{dt}{1 + t^2} \quad \left| \begin{array}{l} \text{Put } \cos x = t \text{ so that } -\sin x dx = dt \\ \text{When } x = 0, t = 1; \text{ When } x = \pi, t = -1; \end{array} \right.$$

$$= \pi \int_1^{-1} \frac{-2 + (1 + t^2)}{1 + t^2} dt = -2\pi \int_1^{-1} \frac{dt}{1 + t^2} + \pi \int_1^{-1} dt$$

$$= -2\pi \left[ \tan^{-1} t \right]_1^{-1} + \pi \left[ t \right]_1^{-1} = -2\pi \left( -\frac{\pi}{4} - \frac{\pi}{4} \right) - 2\pi. \text{ Hence, } I = \pi^2/2 - \pi.$$

**Property V.** 
$$\int_{-a}^a f(x) dx = 2 \int_0^a f(x) dx, \text{ if } f(x) \text{ is an even function,}$$
  

$$= 0 \quad \text{if } f(x) \text{ is an odd function.}$$

(Bhopal, 2008)

$$\int_{-a}^a f(x) dx = \int_{-a}^0 f(x) dx + \int_0^a f(x) dx$$

...(1) [Prop. I]

In  $\int_{-a}^0 f(x) dx$ , put  $x = -t$ , so that  $dx = -dt$

$$\therefore \int_{-a}^0 f(x) dx = - \int_a^0 f(-t) dt = \int_0^a f(-t) dt = \int_0^a f(-x) dx$$

[Prop. II]

Substituting in (1), we get

$$\int_{-a}^a f(x) dx = \int_0^a f(-x) dx + \int_0^a f(x) dx$$

...(2)

(i) If  $f(x)$  is an even function,  $f(-x) = f(x)$ .

$$\therefore \text{ from (2), } \int_{-a}^a f(x) dx = \int_0^a f(x) dx + \int_0^a f(x) dx = 2 \int_0^a f(x) dx$$

(ii) If  $f(x)$  is an odd function,  $f(-x) = -f(x)$ .

$$\therefore \text{ from (2), } \int_{-a}^a f(x) dx = - \int_0^a f(x) dx + \int_0^a f(x) dx = 0.$$

**Property VI.** 
$$\int_0^{2a} f(x) dx = 2 \int_0^a f(x) dx, \text{ if } f(2a - x) = f(x)$$
  

$$= 0, \quad \text{if } f(2a - x) = -f(x)$$

$$\int_0^{2a} f(x) dx = \int_0^a f(x) dx + \int_a^{2a} f(x) dx$$

...(1) [Prop. III]

In  $\int_a^{2a} f(x) dx$ , put  $x = 2a - t$ , so that  $dx = -dt$

Also when  $x = a$ ,  $t = a$ ; when  $x = 2a$ ,  $t = 0$ .

$$\therefore \int_a^{2a} f(x) dx = - \int_a^0 f(2a - t) dt = \int_0^a f(2a - t) dt = \int_0^a f(2a - x) dx$$

[Prop. II]

Substituting in (1), we get

$$\int_0^{2a} f(x) dx = \int_0^a f(x) dx + \int_0^a f(2a - x) dx$$

...(2)

(i) If  $f(2a - x) = f(x)$ , then from (2)

$$\int_0^{2a} f(x) dx = \int_0^a f(x) dx + \int_0^a f(x) dx = 2 \int_0^a f(x) dx$$

(ii) If  $f(2a - x) = -f(x)$ , then from (2)

$$\int_0^{2a} f(x) dx = \int_0^a f(x) dx - \int_0^a f(x) dx = 0.$$

**Cor. 1.** If  $n$  is even,  $\int_0^\pi \sin^m x \cos^n x dx = 2 \int_0^{\pi/2} \sin^m x \cos^n x dx$  and if  $n$  is odd,  $\int_0^\pi \sin^m x \cos^n x dx = 0$ .

**Cor. 2.** If  $m$  is odd,  $\int_0^{2\pi} \sin^m x \cos^n x dx = 0$

$$\begin{aligned} \text{and if } m \text{ is even, } \int_0^{2\pi} \sin^m x \cos^n x dx &= 2 \int_0^\pi \sin^m x \cos^n x dx \\ &= 4 \int_0^{\pi/2} \sin^m x \cos^n x dx, \text{ if } n \text{ is even} = 0, \text{ if } n \text{ is odd.} \end{aligned}$$

**Example 6.15.** Evaluate  $\int_0^\pi \theta \sin^2 \theta \cos^4 \theta d\theta$ .

(V.T.U., 2009 S)

**Solution.** Let  $I = \int_0^\pi \theta \sin^2 \theta \cos^4 \theta d\theta$

$$\text{Then } I = \int_0^\pi (\pi - \theta) \sin^2 (\pi - \theta) \cos^4 (\pi - \theta) d\theta = \pi \int_0^\pi \sin^2 \theta \cos^4 \theta d\theta - I \quad [\text{Prop. IV}]$$

$$\text{or } 2I = \pi \int_0^\pi \sin^2 \theta \cos^4 \theta d\theta = 2\pi \int_0^{\pi/2} \sin^2 \theta \cos^4 \theta d\theta \quad [\text{Prop. VI Cor. 2}]$$

$$= 2\pi \cdot \frac{1 \cdot 3 \cdot 1}{6 \cdot 4 \cdot 2} = \frac{\pi}{2} = \frac{\pi^2}{16}$$

$$\text{Hence } I = \frac{\pi^2}{32}$$

**Example 6.16.** Evaluate  $\int_0^{\pi/2} \log \sin x dx$ .

(Anna, 2005 S)

**Solution.** Let  $I = \int_0^{\pi/2} \log \sin x dx \quad \dots(i)$

$$\text{then } I = \int_0^{\pi/2} \log \sin (\pi/2 - x) dx = \int_0^{\pi/2} \log \cos x dx \quad \dots(ii)$$

Adding (i) and (ii)

$$\begin{aligned} 2I &= \int_0^{\pi/2} (\log \sin x + \log \cos x) dx \\ &= \int_0^{\pi/2} \log (\sin x + \cos x) dx = \int_0^{\pi/2} \log \left( \frac{\sin 2x}{2} \right) dx \\ &= \int_0^{\pi/2} \log \sin 2x dx - \int_0^{\pi/2} \log 2 dx = \int_0^{\pi/2} \log \sin 2x dx - \log 2 \int_0^{\pi/2} dx \\ &= \int_0^{\pi/2} \log \sin 2x dx - \log 2 \cdot x \Big|_0^{\pi/2} = I' - \frac{\pi}{2} \log 2 \quad \dots(iii) \end{aligned}$$

$$\text{where } I' = \int_0^{\pi/2} \log \sin 2x dx \quad \left[ \begin{array}{l} \text{Put, } 2x = t, \text{ so that } 2dx = dt \\ \text{When } x = 0, t = 0; \text{ when } x = \pi/2, t = \pi \end{array} \right]$$

$$= \frac{1}{2} \int_0^\pi \log \sin t dt = \frac{1}{2} \int_0^\pi \log \sin x dx \quad [\because \log \sin (\pi - x) = \log \sin x, \text{ Prop. IV}]$$

$$= \frac{1}{2} \cdot 2 \int_0^{\pi/2} \log \sin x dx = I.$$

Thus from (iii),  $2I = I - (\pi/2) \log 2$ , i.e.,  $I = -(\pi/2) \log 2$ .

Obs. The following are its immediate deductions :

$$\int_0^{\pi/2} \log \sin x \, dx = \int_0^{\pi/2} \log \cos x \, dx = -\frac{\pi}{2} \log 2$$

and  $\int_0^{\pi} \log \sin x \, dx = -\pi \log 2.$

**Example 6.17.** Evaluate  $\int_0^1 \frac{\sin^{-1} x}{x} dx.$

**Solution.** Put  $\sin^{-1} x = \theta$  or  $x = \sin \theta$  so that  $dx = \cos \theta \, d\theta$

Also when  $x = 0, \theta = 0$  ; when  $x = 1, \theta = \pi/2.$

$$\begin{aligned} \therefore \int_0^1 \frac{\sin^{-1} x}{x} dx &= \int_0^{\pi/2} \theta \cdot \frac{\cos \theta}{\sin \theta} d\theta && \text{[Integrate by parts]} \\ &= [\theta \cdot \log \sin \theta]_0^{\pi/2} - \int_0^{\pi/2} 1 \cdot \log \sin \theta \, d\theta \\ &= - \int_0^{\pi/2} \log \sin \theta \, d\theta = - \left( -\frac{\pi}{2} \log 2 \right) = \frac{\pi}{2} \log 2 && \left[ \lim_{x \rightarrow 0} (x \log x) = 0 \right] \end{aligned}$$

#### PROBLEMS 6.4

Prove that :

1. (i)  $\int_0^{\pi/2} \log \tan x \, dx = 0$

(ii)  $\int_0^{\pi/2} \sin 2x \log \tan x \, dx = 0$

2. (i)  $\int_0^{\infty} \frac{x^7 (1-x^{12})}{(1+x)^{28}} dx = 0$

(ii)  $\int_0^{\pi/4} \log (1 + \tan \theta) d\theta = \frac{\pi}{8} \log_e 2$  (Madras, 2000)

3. (i)  $\int_0^{\pi/2} \frac{\sin x}{\sin x + \cos x} dx = \frac{\pi}{4}$

(ii)  $\int_0^a \frac{dx}{x + \sqrt{a^2 + x^2}} = \frac{\pi}{4}$

4. (i)  $\int_0^{\pi/2} \frac{dx}{1 + \sqrt{\cot x}} = \frac{\pi}{4}.$

(ii)  $\int_0^{\pi/2} \frac{\sin^n x}{\sin^n x + \cos^n x} dx = \frac{\pi}{4}.$

5. (i)  $\int_0^{\pi/2} \frac{x \tan x}{\sec x + \cos x} dx = \frac{\pi^2}{4}.$

(ii)  $\int_0^{\pi} \frac{x}{1 + \sin x} dx = \pi$  (Anna, 2002 S)

6. (i)  $\int_0^{\pi} \frac{x \tan x}{\sec x + \tan x} dx = \frac{1}{2} \pi (\pi - 2).$

(ii)  $\int_0^{\pi/2} \frac{x \, dx}{\sin x + \cos x} = \frac{\pi}{2\sqrt{2}} \log (\sqrt{2} + 1).$

Evaluate :

7. (i)  $\int_0^{\pi} \sin^4 x \, dx$

(ii)  $\int_0^{2\pi} \cos^6 x \, dx$

(iii)  $\int_0^{\pi} \sin^6 x \cos^4 x \, dx$  (V.T.U., 2001)

(iv)  $\int_0^{2\pi} \sin^4 x \cos^6 x \, dx$

8. (i)  $\int_0^{\pi} x \sin^7 x \, dx$  (V.T.U., 2009)

(ii)  $\int_0^{\pi} x \cos^4 x \sin^5 x \, dx$  (Marathwada, 2008)

Prove that :

9. (i)  $\int_0^{\pi} \frac{x \, dx}{a^2 \cos^2 x + b^2 \sin^2 x} = \frac{\pi^2}{2ab}$

(ii)  $\int_0^{\pi/2} \frac{x \, dx}{2 \sin^2 x + \cos^2 x} = \frac{\pi^2}{2\sqrt{2}}$

10. (i)  $\int_0^{\pi} \frac{x \, dx}{a^2 - \cos^2 x} = \frac{\pi^2}{2a\sqrt{a^2 - 1}}$  ( $a > 1$ )

(ii)  $\int_0^{\pi} \frac{x \, dx}{1 + \sin^2 x} = \frac{\pi^2}{2\sqrt{2}}.$



$$11. \int_0^{\pi} \log(1 + \cos \theta) d\theta = -\pi \log_e 2$$

(Madras, 2003)

$$12. (i) \int_0^{\infty} \frac{\log(1+x^2)}{1+x^2} dx = \pi \log_e 2$$

$$(ii) \int_0^{\infty} \frac{\log(x+1/x)}{1+x^2} dx = \pi \log_e 2.$$

## 6.9 (1) INTEGRAL AS THE LIMIT OF A SUM

We have so far considered integration as inverse of differentiation. We shall now define the definite integral as the limit of a sum :

**Def.** If  $f(x)$  is continuous and single valued in the interval  $[a, b]$ , then the definite integral of  $f(x)$  between the limits  $a$  and  $b$  is defined by the equation

$$\int_a^b f(x) dx = \lim_{h \rightarrow 0} h [f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)],$$

where  $nh = b - a$ .

...(1)

### (2) Evaluation of limits of series

The summation definition of a definite integral enables us to express the limits of sums of certain types of series as definite integrals which can be easily evaluated. We rewrite (1) as follows :

$$\int_a^b f(x) dx = \lim_{n \rightarrow \infty} h \sum_{r=0}^{n-1} f(a+rh), \text{ where } nh = b - a.$$

Putting  $a = 0$  and  $b = 1$ , so that  $h = 1/n$ , we get

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{r=0}^{n-1} f\left(\frac{r}{n}\right) = \int_0^1 f(x) dx$$

Thus to express a given series as definite integral:

(i) Write the general term ( $T_r$  or  $T_{r+1}$  whichever involves  $r$ )  
i.e.,  $f(r/n) \cdot 1/n$

(ii) Replace  $r/n$  by  $x$  and  $1/n$  by  $dx$ ,

(iii) Integrate the resulting expression, taking

the lower limit =  $\lim_{n \rightarrow \infty} (r/n)$  where  $r$  is as in the first term,

and the upper limit =  $\lim_{n \rightarrow \infty} (r/n)$  where  $r$  is as in the last term.

**Example 6.18.** Find the limit, when  $n \rightarrow \infty$ , of the series

$$\frac{n}{n^2} + \frac{n}{n^2 + 1^2} + \frac{n}{n^2 + 2^2} + \dots + \frac{n}{n^2 + (n-1)^2}$$

**Solution.** Here the general term ( $= T_{r+1}$ ) =  $\frac{n}{n^2 + r^2} = \frac{n}{1 + (r/n)^2} \cdot \frac{1}{n}$

$$= \frac{1}{1+x^2} dx \quad [\text{Putting } r/n = x \text{ and } 1/n = dx]$$

Now for the first term  $r = 0$  and for the last term  $r = n - 1$

$$\therefore \text{ the lower limit of integration } = \lim_{n \rightarrow \infty} \left(\frac{0}{n}\right) = 0$$

$$\text{and the upper limit of integration } = \lim_{n \rightarrow \infty} \left(\frac{n-1}{n}\right) = \lim_{n \rightarrow \infty} \left(1 - \frac{1}{n}\right) = 1.$$

$$\text{Hence, the required limit } = \int_0^1 \frac{dx}{1+x^2} = \left| \tan^{-1} x \right|_0^1 = \tan^{-1}(1) - \tan^{-1}(0) = \pi/4.$$

[To find limit of a product by integration :

Let  $P = \lim_{n \rightarrow \infty} \left( \text{given product} \right)$

Take logs of both sides, so that

$$\log P = \lim_{n \rightarrow \infty} (\text{a series}) = k \text{ (say). Then } P = e^k.]$$

**Example 6.19.** Evaluate  $\lim_{n \rightarrow \infty} \left\{ \left(1 + \frac{1}{n}\right) \left(1 + \frac{2}{n}\right) \dots \left(1 + \frac{n}{n}\right) \right\}^{1/n}$ . (Bhopal, 2008)

**Solution.** Let  $P = \lim_{n \rightarrow \infty} \left\{ \left(1 + \frac{1}{n}\right) \left(1 + \frac{2}{n}\right) \dots \left(1 + \frac{n}{n}\right) \right\}^{1/n}$ .

Taking logs of both sides,

$$\log P = \lim_{n \rightarrow \infty} \frac{1}{n} \left\{ \log \left(1 + \frac{1}{n}\right) + \log \left(1 + \frac{2}{n}\right) + \dots + \log \left(1 + \frac{n}{n}\right) \right\}$$

Its general term  $= \log \left(1 + \frac{r}{n}\right) \cdot \frac{1}{n} = \log (1+x) \cdot dx$  [Putting  $r/n = x$  and  $1/n = dx$ ]

Also for first term  $r = 1$  and for the last term  $r = n$ .

$\therefore$  The lower limit of integration  $= \lim_{n \rightarrow \infty} (1/n) = 0$  and the upper limit  $= \lim_{n \rightarrow \infty} (n/n) = 1$

Hence  $\log P = \int_0^1 \log (1+x) dx = \int_0^1 \log (1+x) \cdot 1 dx$  [Integrate by parts]

$$= \left[ \log (1+x) \cdot x \right]_0^1 - \int_0^1 \frac{1}{1+x} \cdot x dx$$

$$= \log 2 - \int_0^1 \frac{1+x-1}{1+x} dx = \log 2 - \int_0^1 dx + \int_0^1 \frac{dx}{1+x}$$

$$= \log 2 - \left[ x \right]_0^1 + \left[ \log (1+x) \right]_0^1 = \log 2 - 1 + \log 2$$

$$= \log 2^2 - \log_e e = \log (4/e). \text{ Hence, } P = 4/e.$$

### PROBLEMS 6.5

Find the limit, as  $n \rightarrow \infty$ , of the series :

$$1. \frac{1}{n+1} + \frac{1}{n+2} + \frac{1}{n+3} + \dots + \frac{1}{2n}. \quad (\text{Bhopal, 2009}) \quad 2. \frac{1}{n^3+1} + \frac{4}{n^3+8} + \frac{9}{n^3+27} + \dots + \frac{r^2}{n^3+r^3} + \dots + \frac{1}{2n}.$$

$$3. \lim_{n \rightarrow \infty} \frac{\sqrt{n}}{\sqrt{n^3}} + \frac{\sqrt{n}}{\sqrt{(n+3)^3}} + \frac{\sqrt{n}}{\sqrt{(n+6)^3}} + \dots + \frac{\sqrt{n}}{\sqrt{(n+3(n-1))^3}}.$$

Evaluate :

$$4. \lim_{n \rightarrow \infty} \sum_{r=1}^{n-1} \frac{1}{\sqrt{(n^2-r^2)}} \quad (\text{Bhopal, 2008})$$

$$5. \lim_{n \rightarrow \infty} \frac{[(n+1)(n+2) \dots (n+n)]^{1/n}}{n}.$$

$$6. \lim_{n \rightarrow \infty} \left\{ \left(1 + \frac{1}{n^2}\right) \left(1 + \frac{2^2}{n^2}\right) \left(1 + \frac{3^2}{n^2}\right) \dots \left(1 + \frac{n^2}{n^2}\right) \right\}^{1/n} \quad (\text{Bhopal, 2008})$$

### 6.10 AREAS OF CARTESIAN CURVES

(1) Area bounded by the curve  $y = f(x)$ , the  $x$ -axis and the ordinates  $x = a$ ,  $x = b$  is  $\int_a^b y \, dx$ .

Let  $AB$  be the curve  $y = f(x)$  between the ordinates  $LA$  ( $x = a$ ) and  $MB$  ( $x = b$ ). (Fig. 6.1)

Let  $P(x, y)$ ,  $P'(x + \delta x, y + \delta y)$  be two neighbouring points on the curve and  $NP$ ,  $N'P'$  be their respective ordinates.



Let the area  $ALNP$  be  $A$ , which depends on the position of  $P$  whose abscissa is  $x$ . Then the area  $PNN'P' = \delta A$ .

Complete the rectangles  $PN'$  and  $P'N'$ .

Then the area  $PNN'P'$  lies between the areas of the rectangles  $PN'$  and  $P'N'$ .

i.e.,  $\delta A$  lies between  $y\delta x$  and  $(y + \delta y)\delta x$

$\therefore \frac{\delta A}{\delta x}$  lies between  $y$  and  $y + \delta y$ .

Now taking limits as  $P' \rightarrow P$  i.e.,  $\delta x \rightarrow 0$  (and  $\therefore \delta y \rightarrow 0$ ),

$$dA/dx = y$$

Integrating both sides between the limits  $x = a$  and  $x = b$ , we have

$$A \Big|_a^b = \int_a^b y \, dx$$

or (value of  $A$  for  $x = b$ ) – (value of  $A$  for  $x = a$ ) =  $\int_a^b y \, dx$

Thus area  $ALMB = \int_a^b y \, dx$ .

(2) Interchanging  $x$  and  $y$  in the above formula, we see that the area bounded by the curve  $x = f(y)$ , the  $y$ -axis and the abscissae  $y = a, y = b$  is  $\int_a^b x \, dy$ . (Fig. 6.2)




Fig. 6.1




Fig. 6.2

**Obs. 1.** The area bounded by a curve, the  $x$ -axis and two ordinates is called the **area under the curve**. The process of finding the area of plane curves is often called **quadrature**.

**Obs. 2. Sign of an area.** An area whose boundary is described in the anti-clockwise direction is considered positive and an area whose boundary is described in the clockwise direction is taken as negative.

In Fig. 6.3, the area  $ALMB \left( = \int_a^b y \, dx \right)$  which is described in the anti-clockwise direction and lies above the  $x$ -axis, will give a positive result.

In Fig. 6.4, the area  $ALMB \left( = \int_a^b y \, dx \right)$  which is described in the clockwise direction and lies below the  $x$ -axis, will give a negative result.




Fig. 6.3




Fig. 6.4




Fig. 6.5

In Fig. 6.5, the area  $ALNB \left( = \int_a^b y \, dx \right)$  will not consist of the sum of the area  $ALN \left( = \int_a^c y \, dx \right)$  and the area  $NMB \left( = \int_c^b y \, dx \right)$ , but their difference.

Thus to find the total area in such cases the numerical value of the area of each portion must be evaluated separately and their results added afterwards.

**Example 6.20.** Find the area of the loop of the curve  $ay^2 = x^2(a - x)$ . (S.V.T.U., 2009 ; Osmania, 2000)

**Solution.** Let us trace the curve roughly to get the limits of integration.

(i) The curve is symmetrical about  $x$ -axis.



- (ii) It passes through the origin. The tangents at the origin are  $ay^2 = ax^2$  or  $y = \pm x$ .  $\therefore$  Origin is a node.
- (iii) The curve has no asymptotes.
- (iv) The curve meets the  $x$ -axis at  $(0, 0)$  and  $(a, 0)$ . It meets the  $y$ -axis at  $(0, 0)$  only.

From the equation of the curve, we have  $y = \frac{x}{\sqrt{a}} \sqrt{a-x}$

For  $x > a$ ,  $y$  is imaginary. Thus no portion of the curve lies to the right of the line  $x = a$ . Also  $x \rightarrow -\infty, y \rightarrow \infty$ .

Thus the curve is as shown in Fig. 6.6.

$\therefore$  Area of the loop = 2 (area of upper half of the loop)

$$\begin{aligned} &= 2 \int_0^a y \, dx = 2 \int_0^a x \sqrt{\frac{a-x}{a}} \, dx = \frac{2}{\sqrt{a}} \int_0^a [a - (a-x)] \sqrt{a-x} \, dx \\ &= \frac{2}{\sqrt{a}} \int_0^a [a(a-x)^{1/2} - (a-x)^{3/2}] \, dx = 2\sqrt{a} \left[ \frac{(a-x)^{3/2}}{-3/2} \right]_0^a - \frac{2}{\sqrt{a}} \left[ \frac{(a-x)^{5/2}}{-5/2} \right]_0^a \\ &= -\frac{4}{3} \sqrt{a} (0 - a^{3/2}) + \frac{4}{5\sqrt{a}} (0 - a^{5/2}) = \frac{4}{3} a^2 - \frac{4}{5} a^2 = \frac{8}{15} a^2. \end{aligned}$$




Fig. 6.6

**Example 6.21.** Find the area included between the curve  $y^2(2a-x) = x^3$  and its asymptote. (V.T.U., 2003)

**Solution.** The curve is as shown in Fig. 4.23.

Area between the curve and the asymptote

$$\begin{aligned} &= 2 \int_0^{2a} y \, dx = 2 \int_0^{2a} \sqrt{\frac{x^3}{2a-x}} \, dx \\ &= 2 \int_0^{\pi/2} \sqrt{\frac{(2a \sin^2 \theta)^3}{2a \cos^2 \theta}} \cdot 4a \sin \theta \cos \theta \, d\theta \\ &= 16a^2 \int_0^{\pi/2} \sin^4 \theta \, d\theta = 16a^2 \cdot \frac{3 \cdot 1}{4 \cdot 2} \cdot \frac{\pi}{2} = 3\pi a^2. \end{aligned}$$

Put  $x = 2a \sin^2 \theta$   
so that  $dx = 4a \sin \theta \cos \theta \, d\theta$

**Example 6.22.** Find the area enclosed by the curve  $a^2 x^2 = y^3(2a-y)$ .

**Solution.** Let us first find the limits of integration.

- (i) The curve is symmetrical about  $y$ -axis.
- (ii) It passes through the origin and the tangents at the origin are  $x^2 = 0$  or  $x = 0, x = 0$ .  
 $\therefore$  There is a cusp at the origin.
- (iii) The curve has no asymptote.
- (iv) The curve meets the  $x$ -axis at the origin only and meets the  $y$ -axis at  $(0, 2a)$ . From the equation of the curve, we have

$$x = \frac{y}{a} \sqrt{y(2a-y)}$$

For  $y < 0$  or  $y > 2a$ ,  $x$  is imaginary. Thus the curve entirely lies between  $y = 0$  ( $x$ -axis) and  $y = 2a$ , which is shown in Fig. 6.7.




Fig. 6.7

$$\begin{aligned} \therefore \text{Area of the curve} &= 2 \int_0^{2a} x \, dy = \frac{2}{a} \int_0^{2a} y \sqrt{y(2a-y)} \, dy \\ &= \frac{2}{a} \int_0^{\pi/2} 2a \sin^2 \theta \sqrt{[2a \sin^2 \theta (2a - 2a \sin^2 \theta)]} \times 4a \sin \theta \cos \theta \, d\theta \\ &= 32a^2 \int_0^{\pi/2} \sin^4 \theta \cos^2 \theta \, d\theta = 32a^2 \frac{3 \cdot 1 \times 1}{6 \cdot 4 \cdot 2} \cdot \frac{\pi}{2} = \pi a^2. \end{aligned}$$

Put  $y = 2a \sin^2 \theta$   
 $\therefore dy = 4a \sin \theta \cos \theta \, d\theta$

**Example 6.23.** Find the area enclosed between one arch of the cycloid  $x = a(\theta - \sin \theta)$ ,  $y = a(1 - \cos \theta)$ ; and its base. (V.T.U., 2000)

**Solution.** To describe its first arch,  $\theta$  varies from 0 to  $2\pi$  i.e.,  $x$  varies from 0 to  $2a\pi$  (Fig. 6.8).

$$\therefore \text{ Required area} = \int_{x=0}^{2\pi a} y \, dx$$

where  $y = a(1 - \cos \theta)$ ,  $dx = a(1 - \cos \theta) \, d\theta$ .

$$\begin{aligned} &= \int_{\theta=0}^{\pi/2} a(1 - \cos \theta) \cdot a(1 - \cos \theta) \, d\theta \\ &= 2a^2 \int_0^{\pi} (1 - \cos \theta)^2 \, d\theta = 8a^2 \int_0^{\pi} \sin^4 \frac{\theta}{2} \, d\theta \\ &= 16a^2 \int_0^{\pi/2} \sin^4 \phi \, d\phi, \text{ putting } \theta/2 = \phi \text{ so that } d\theta = 2d\phi. \\ &= 16a^2 \cdot \frac{3}{4} \cdot \frac{1}{2} \cdot \frac{\pi}{2} = 3\pi a^2. \end{aligned}$$




Fig. 6.8

**Example 6.24.** Find the area of the tangent cut off from the parabola  $x^2 = 8y$  by the line  $x - 2y + 8 = 0$ .

**Solution.** Given parabola is  $x^2 = 8y$  ... (i)

and the straight line is  $x - 2y + 8 = 0$  ... (ii)

Substituting the value of  $y$  from (ii) in (i), we get

$$x^2 = 4(x + 8) \text{ or } x^2 - 4x - 32 = 0$$

or  $(x - 8)(x + 4) = 0 \therefore x = 8, -4$ .

Thus (i) and (ii) intersect at  $P$  and  $Q$  where  $x = 8$  and  $x = -4$ . (Fig. 6.9)

$\therefore$  Required area  $POQ$  (i.e., dotted area) = area bounded by straight line (ii) and  $x$ -axis from  $x = -4$  to  $x = 8$  - area bounded by parabola (i) and  $x$ -axis from  $x = -4$  to  $x = 8$ .

$$\begin{aligned} &= \int_{-4}^8 y \, dx, \text{ from (ii)} - \int_{-4}^8 y \, dx, \text{ from (i)} \\ &= \int_{-4}^8 \frac{x+8}{2} \, dx - \int_{-4}^8 \frac{x^2}{8} \, dx = \frac{1}{2} \left[ \frac{x^2}{2} + 8x \right]_{-4}^8 - \frac{1}{8} \left[ \frac{x^3}{3} \right]_{-4}^8 \\ &= \frac{1}{2} \{ (32 + 64) - (-24) \} - \frac{1}{24} (512 + 64) = 36. \end{aligned}$$




Fig. 6.9

**Example 6.25.** Find the area common to the parabola  $y^2 = ax$  and the circle  $x^2 + y^2 = 4ax$ .

**Solution.** Given parabola is  $y^2 = ax$  ... (i)

and the circle is  $x^2 + y^2 = 4ax$  ... (ii)

Both these curves are symmetrical about  $x$ -axis. Solving (i) and (ii) for  $x$ , we have

$$x^2 + ax = 4ax \text{ or } x(x - 3a) = 0$$

or  $x = 0, 3a$ .

Thus the two curves intersect at the points where  $x = 0$  and  $x = 3a$ . (Fig. 6.10).

Also (ii) meets the  $x$ -axis at  $A(4a, 0)$ .

Area common to (i) and (ii) i.e., the shaded area

$$\begin{aligned} &= 2[\text{Area } ORP + \text{Area } PRA] \quad (\text{By symmetry}) \\ &= 2 \left[ \int_0^{3a} y \, dx, \text{ from (i)} + \int_{3a}^{4a} y \, dx, \text{ from (ii)} \right] \\ &= 2 \left[ \int_0^{3a} \sqrt{ax} \, dx + \int_{3a}^{4a} \sqrt{4ax - x^2} \, dx \right] \end{aligned}$$




Fig. 6.10



$$\begin{aligned}
 &= 2\sqrt{a} \left[ \frac{x^{3/2}}{3/2} \right]_{3a}^{4a} + 2 \int_{3a}^{4a} \sqrt{4a^2 - (x-2a)^2} dx \\
 &= \frac{4\sqrt{a}}{3} (3a)^{3/2} + 2 \left[ \frac{1}{2} (x-2a) \sqrt{4a^2 - (x-2a)^2} + \frac{4a^2}{2} \sin^{-1} \frac{x-2a}{2a} \right]_{3a}^{4a} \\
 &= 4\sqrt{3} a^2 + 2 \left[ \left( 0 - \frac{1}{2} a \sqrt{3} a \right) + 2a^2 \left( \frac{\pi}{2} - \frac{\pi}{6} \right) \right] \\
 &= 4\sqrt{3} a^2 - \sqrt{3} a^2 + \frac{4}{3} \pi a^2 = \left( 3\sqrt{3} + \frac{4}{3} \pi \right) a^2.
 \end{aligned}$$

### PROBLEMS 6.6

- (i) Find the area of the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ . (Kerala, 2005 ; V.T.U., 2003 S)

(ii) Find the area bounded by the parabola  $y^2 = 4ax$  and its latus-rectum.
- Find the area bounded by the curve  $y = x(x-3)(x-5)$  and the  $x$ -axis.
- Find the area included between the curve  $ay^2 = x^3$ , the  $x$ -axis and the ordinates  $x = a$ .
- Find the area of the loop of the curve :
 

(i)  $3ay^2 = x(x-a)^2$  (Rajasthan, 2005)
(ii)  $x(x^2 + y^2) = a(x^2 - y^2)$  (P.T.U., 2010)
- Find the whole area of the curve :
 

(i)  $a^2x^2 = y^3(2a - y)$  (Nagpur, 2009)
(ii)  $8a^2y^2 = x^2(a^2 - x^2)$  (V.T.U., 2006)
- Find the area included between the curve and its asymptotes in each case :
 

(i)  $xy^2 = a^2(a - x)$ . (V.T.U., 2003)
(ii)  $x^2y^2 = a^2(y^2 - x^2)$ . (V.T.U., 2007)
- Show that the area of the loop of the curve  $y^2(a+x) = x^2(3a-x)$  is equal to the area between the curve and its asymptote.
- Find the whole area of the *astroid*  $x^{2/3} + y^{2/3} = a^{2/3}$  or  $x = a \cos^3 \theta$ ,  $y = a \sin^3 \theta$ . (V.T.U., 2005)
- Find the area bounded by the curve  $\sqrt{x} + \sqrt{y} = \sqrt{a}$  and the coordinate axes.
- Find the area included between the *cycloid*  $x = a(\theta + \sin \theta)$ ,  $y = a(1 - \cos \theta)$  and its base. Also find the area between the curve and the  $x$ -axis. (Gorakhpur, 1999)
- Find the area common to the circles  $x^2 + y^2 = 4$  and  $x^2 + y^2 = 4x$ .
- Prove that the area common to the parabolas  $x^2 = 4ay$  and  $y^2 = 4ax$  is  $16a^2/3$ . (S.V.T.U., 2008 ; Kurukshetra, 2005)
- Find the area included between the circle  $x^2 + y^2 = 2ax$  and the parabola  $y^2 = ax$ .
- Find the area bounded by the parabola  $y^2 = 4ax$  and the line  $x + y = 3a$ .
- Find the area of the segment cut off from the parabola  $y = 4x - x^2$  by the straight line  $y = x$ . (V.T.U., 2010 ; S.V.T.U., 2008)

### (2) Areas of polar curves. Area bounded by the curve $r = f(\theta)$ and the radii vectors

$$\theta = \alpha, \theta = \beta \text{ is } \frac{1}{2} \int_{\alpha}^{\beta} r^2 d\theta.$$

Let  $AB$  be the curve  $r = f(\theta)$  between the radii vectors  $OA(\theta = \alpha)$  and  $OB(\theta = \beta)$ . Let  $P(r, \theta)$ ,  $P'(r + \delta r, \theta + \delta \theta)$  be any two neighbouring points on the curve. (Fig. 6.11)

Let the area  $OAP = A$  which is a function of  $\theta$ . Then the area  $OPP' = \delta A$ . Mark circular arcs  $PQ$  and  $P'Q'$  with centre  $O$  and radii  $OP$  and  $OP'$ .

Evidently area  $OPP'$  lies between the sectors  $OPQ$  and  $OP'Q'$  i.e.,  $\delta A$  lies between  $\frac{1}{2} r^2 \delta \theta$  and  $\frac{1}{2} (r + \delta r)^2 \delta \theta$ .

$$\therefore \frac{\delta A}{\delta \theta} \text{ lies between } \frac{1}{2} r^2 \text{ and } \frac{1}{2} (r + \delta r)^2.$$




Fig. 6.11



Now taking limits as  $\delta\theta \rightarrow 0$  ( $\therefore \delta r \rightarrow 0$ ),  $\frac{dA}{d\theta} = \frac{1}{2}r^2$

Integrating both sides from  $\theta = \alpha$  to  $\theta = \beta$ , we get  $A \Big|_{\alpha}^{\beta} = \int_{\alpha}^{\beta} \frac{1}{2}r^2 d\theta$

or (value of  $A$  for  $\theta = \beta$ ) – (value of  $A$  for  $\theta = \alpha$ ) =  $\frac{1}{2} \int_{\alpha}^{\beta} r^2 d\theta$

Hence the required area  $OAB = \frac{1}{2} \int_{\alpha}^{\beta} r^2 d\theta$ .

**Example 6.26.** Find the area of the cardioid  $r = a(1 - \cos \theta)$ .

(V.T.U., 2004)

**Solution.** The curve is as shown in Fig. 6.12. Its upper half is traced from  $\theta = 0$  to  $\theta = \pi$ .

$$\begin{aligned} \therefore \text{Area of the curve} &= 2 \cdot \frac{1}{2} \int_0^{\pi} r^2 d\theta = a^2 \int_0^{\pi} (1 - \cos \theta)^2 d\theta \\ &= a^2 \int_0^{\pi} (2 \sin^2 \theta/2) d\theta = 4a^2 \int_0^{\pi} \sin^4 \theta/2 \cdot d\theta \\ &= 8a^2 \int_0^{\pi/2} \sin^4 \phi d\phi, \text{ putting } \theta/2 = \phi \text{ and } d\theta = 2d\phi. \\ &= 8a^2 \cdot \frac{3 \cdot 1}{4 \cdot 2} \cdot \frac{\pi}{2} = \frac{3\pi a^2}{2}. \end{aligned}$$




Fig. 6.12

**Example 6.27.** Find the area of a loop of the curve  $r = a \sin 3\theta$ .

**Solution.** The curve is as shown in Fig. 4.35. It consists of three loops.

Putting  $r = 0$ ,  $\sin 3\theta = 0 \therefore 3\theta = 0$  or  $\pi$  i.e.,  $\theta = 0$  or  $\pi/3$  which are the limits for the first loop.

$$\begin{aligned} \therefore \text{Area of a loop} &= \frac{1}{2} \int_0^{\pi/3} r^2 d\theta = \frac{1}{2} a^2 \int_0^{\pi/3} \sin^2 3\theta d\theta = \frac{a^2}{4} \int_0^{\pi/3} (1 - \cos 6\theta) d\theta \\ &= \frac{a^2}{4} \left[ \theta - \frac{\sin 6\theta}{6} \right]_0^{\pi/3} = \frac{a^2}{4} \left( \frac{\pi}{3} - 0 \right) = \frac{\pi a^2}{12}. \end{aligned}$$

**Obs.** The limits of integration for a loop of  $r = a \sin n\theta$  or  $r = a \cos n\theta$  are the two consecutive values of  $\theta$  when  $r = 0$ .

**Example 6.28.** Prove that the area of a loop of the curve  $x^3 + y^3 = 3axy$  is  $3a^2/2$ .

**Solution.** Changing to polar form (by putting  $x = r \cos \theta$ ,  $y = r \sin \theta$ ),  $r = \frac{3a \sin \theta \cos \theta}{\cos^3 \theta + \sin^3 \theta}$

Putting  $r = 0$ ,  $\sin \theta \cos \theta = 0$ .

$\therefore \theta = 0, \pi/2$ , which are the limits of integration for its loop.

$\therefore$  Area of the loop

$$\begin{aligned} &= \frac{1}{2} \int_0^{\pi/2} r^2 d\theta = \frac{1}{2} \int_0^{\pi/2} \frac{9a^2 \sin^2 \theta \cos^2 \theta}{(\cos^3 \theta + \sin^3 \theta)^2} d\theta \\ &= \frac{9a^2}{2} \int_0^{\pi/2} \frac{\tan^2 \theta \sec^2 \theta}{(1 + \tan^3 \theta)^2} d\theta \quad [\text{Dividing num. and denom. by } \cos^6 \theta] \\ &= \frac{3a^2}{2} \int_1^{\infty} \frac{dt}{t^2}, \quad \text{putting } 1 + \tan^3 \theta = t \text{ and } 3 \tan^2 \theta \sec^2 \theta d\theta = dt. \\ &= \frac{3a^2}{2} \left[ \frac{t^{-1}}{-1} \right]_1^{\infty} = \frac{3a^2}{2} (-0 + 1) = \frac{3a^2}{2}. \end{aligned}$$

**Example 6.29.** Find the area common to the circles

$$r = a\sqrt{2} \text{ and } r = 2a \cos \theta$$

**Solution.** The equations of the circles are  $r = a\sqrt{2}$  ... (i) and  $r = 2a \cos \theta$  ... (ii)

(i) represents a circle with centre at (0, 0) and radius  $a\sqrt{2}$ . (ii) represents a circle symmetrical about OX, with centre at (a, 0) and radius a.

The circles are shown in Fig. 6.13. At their point of intersection P, eliminating r from (i) and (ii),

$$a\sqrt{2} = 2a \cos \theta \text{ i.e., } \cos \theta = 1/\sqrt{2}$$

or

$$\theta = \pi/4$$

$\therefore$  Required area = 2  $\times$  area OAPQ

(By symmetry)

$$= 2(\text{area OAP} + \text{area OPQ})$$

$$= 2 \left[ \frac{1}{2} \int_0^{\pi/4} r^2 d\theta, \text{ for (i)} + \frac{1}{2} \int_{\pi/4}^{\pi/2} r^2 d\theta, \text{ for (ii)} \right]$$

$$= \int_0^{\pi/4} (a\sqrt{2})^2 d\theta + \int_{\pi/4}^{\pi/2} (2a \cos \theta)^2 d\theta = 2a^2 \left| \theta \right|_0^{\pi/4} + 4a^2 \int_{\pi/4}^{\pi/2} \frac{1 + \cos 2\theta}{2} d\theta$$

$$= 2a^2 (\pi/4 - 0) + 2a^2 \left| \theta + \frac{\sin 2\theta}{2} \right|_{\pi/4}^{\pi/2} = \frac{\pi a^2}{2} + 2a^2 \left( \frac{\pi}{2} - \frac{\pi}{4} - \frac{1}{2} \right) = a^2 (\pi - 1).$$




Fig. 6.13

**Example 6.30.** Find the area common to the cardioids  $r = a(1 + \cos \theta)$  and  $r = a(1 - \cos \theta)$ .

(Kurukshetra, 2006 ; V.T.U., 2006)

**Solution.** The cardioid  $r = a(1 + \cos \theta)$  is ABCOB'A and the cardioid  $r = a(1 - \cos \theta)$  is OC'BA'B'O.

Both the cardioids are symmetrical about the initial line OX and intersect at B and B' (Fig. 6.14)

$\therefore$  Required area (shaded) = 2 area OC'BCO

$$= 2 [\text{area OC'BO} + \text{area OBCO}]$$

$$= 2 \left[ \left\{ \int_0^{\pi/2} \frac{1}{2} r^2 d\theta \right\}_{r=a(1-\cos\theta)} + \left\{ \int_{\pi/2}^{\pi} \frac{1}{2} r^2 d\theta \right\}_{r=a(1+\cos\theta)} \right]$$

$$= a^2 \int_0^{\pi/2} (1 - \cos \theta)^2 d\theta + a^2 \int_{\pi/2}^{\pi} (1 + \cos \theta)^2 d\theta$$

$$= a^2 \left\{ \int_0^{\pi/2} (1 - 2 \cos \theta + \cos^2 \theta) d\theta + \int_{\pi/2}^{\pi} [1 + 2 \cos \theta + \cos^2 \theta] d\theta \right\}$$

$$= a^2 \left\{ \int_0^{\pi} (1 + \cos^2 \theta) d\theta - 2 \int_0^{\pi/2} \cos \theta d\theta + 2 \int_{\pi/2}^{\pi} \cos \theta d\theta \right\}$$

$$= a^2 \left\{ \int_0^{\pi} \left( 1 + \frac{1 + \cos 2\theta}{2} \right) d\theta - 2 \left| \sin \theta \right|_0^{\pi/2} + 2 \left| \sin \theta \right|_{\pi/2}^{\pi} \right\}$$

$$= a^2 \left\{ \left[ \frac{3}{2} \theta + \frac{\sin 2\theta}{4} \right]_0^{\pi} - 2(1 - 0) + 2(0 - 1) \right\} = \left( \frac{3\pi}{2} - 4 \right) a^2.$$




Fig. 6.14

### PROBLEMS 6.7

- Find the whole area of  
(i) the cardioid  $r = a(1 + \cos \theta)$  (V.T.U., 2008) (ii) the lemniscate  $r^2 = a^2 \cos 2\theta$ ; (V.T.U., 2006)
- Find the area of one loop of the curve  
(i)  $r = a \sin 2\theta$ . (ii)  $r = a \cos 3\theta$ .
- Show that the area included between the folium  $x^3 + y^3 = 3axy$  and its asymptote is equal to the area of loop.
- Prove that the area of the loop of the curve  $x^3 + y^3 = 3axy$  is three times the area of the loop of the curve  $r^2 = a^2 \cos 2\theta$ .
- Find the area inside the circle  $r = a \sin \theta$  and lying outside the cardioid  $r = a(1 - \cos \theta)$ . (Anna, 2009)
- Find the area outside the circle  $r = 2a \cos \theta$  and inside the cardioid  $r = a(1 + \cos \theta)$ . (Kurukshetra, 2006)



### 6.11 LENGTHS OF CURVES

(1) The length of the arc of the curve  $y = f(x)$  between the points where  $x = a$  and  $x = b$  is

$$\int_a^b \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$

Let  $AB$  be the curve  $y = f(x)$  between the points  $A$  and  $B$  where  $x = a$  and  $x = b$  (Fig. 6.15)

Let  $P(x, y)$  be any point on the curve and arc  $AP = x$  so that it is a function of  $x$ .

Then 
$$\frac{ds}{dx} = \sqrt{1 + \left(\frac{dy}{dx}\right)^2} \quad [(1) \text{ of p. 164}]$$

$$\begin{aligned} \therefore \int_a^b \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx &= \int_a^b \frac{ds}{dx} \cdot dx = \left| s \right|_{x=a}^{x=b} \\ &= (\text{value of } s \text{ for } x = b) - (\text{value of } s \text{ for } x = a) = \text{arc } AB - 0 \end{aligned}$$

Hence, the arc  $AB = \int_a^b \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$ .

(2) The length of the arc of the curve  $x = f(y)$  between the points where  $y = a$  and  $y = b$ , is

$$\int_a^b \sqrt{1 + \left(\frac{dx}{dy}\right)^2} dy \quad [\text{Use (2) of p. 165}]$$

(3) The length of the arc of the curve  $x = f(t)$ ,  $y = \phi(t)$  between the points where  $t = a$  and  $t = b$ , is

$$\int_a^b \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt. \quad [\text{Use (3) p. 165}]$$

(4) The length of the arc of the curve  $r = f(\theta)$  between the points where  $\theta = \alpha$  and  $\theta = \beta$ , is

$$\int_\alpha^\beta \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2} d\theta. \quad [\text{Use (1) of p. 165}]$$

(5) The length of the arc of the curve  $\theta = f(r)$  between the points where  $r = a$  and  $r = b$ , is

$$\int_a^b \sqrt{1 + \left(r \frac{d\theta}{dr}\right)^2} dr \quad [\text{Use (2) of p. 166}]$$




Fig. 6.15

**Example 6.31.** Find the length of the arc of the parabola  $x^2 = 4ay$  measured from the vertex to one extremity of the latus-rectum. (Delhi, 2002)

**Solution.** Let  $A$  be the vertex and  $L$  an extremity of the latus-rectum so that at  $A$ ,  $x = 0$  and at  $L$ ,  $x = 2a$ . (Fig. 6.16).

Now  $y = x^2/4a$  so that  $\frac{dy}{dx} = \frac{1}{4a} \cdot 2x = \frac{x}{2a}$

$$\begin{aligned} \therefore \text{arc } AL &= \int_0^{2a} \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx \\ &= \int_0^{2a} \sqrt{1 + \left(\frac{x}{2a}\right)^2} dx = \frac{1}{2a} \int_0^{2a} \sqrt{(2a)^2 + x^2} dx \end{aligned}$$




Fig. 6.16



$$= \frac{1}{2a} \left[ \frac{x\sqrt{(2a)^2 + x^2}}{2} + \frac{(2a)^2}{2} \sinh^{-1} \frac{x}{2a} \right]_0^{2a} = \frac{1}{2a} \left[ \frac{2a\sqrt{(2a)^2 + (2a)^2}}{2} + 2a^2 \sinh^{-1} 1 \right]$$

$$= a[\sqrt{2} + \sinh^{-1} 1] = a[\sqrt{2} + \log(1 + \sqrt{2})] \quad [\because \sinh^{-1} x = \log(x + \sqrt{1+x^2})]$$

**Example 6.32.** Find the perimeter of the loop of the curve  $3ay^2 = x(x-a)^2$ .

**Solution.** The curve is symmetrical about the  $x$ -axis and the loop lies between the limits  $x = 0$  and  $x = a$ . (Fig. 6.17).

We have  $y = \frac{\sqrt{x(x-a)}}{\sqrt{3a}}$

$$\therefore \frac{dy}{dx} = \frac{1}{\sqrt{3a}} \left[ \frac{3}{2} x^{1/2} - \frac{a}{2} x^{-1/2} \right] = \frac{1}{2\sqrt{3a}} \frac{3x-a}{\sqrt{x}}$$

$$\therefore \text{Perimeter of the loop} = 2 \int_0^a \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx \quad (\text{By symmetry})$$

$$= 2 \int_0^a \sqrt{1 + \frac{(3x-a)^2}{12ax}} dx = 2 \int_0^a \frac{\sqrt{9x^2 + 6ax + a^2}}{\sqrt{12ax}} dx$$

$$= \frac{1}{\sqrt{3a}} \int_0^a \frac{3x+a}{\sqrt{x}} dx = \frac{1}{\sqrt{3a}} \int_0^a (3x^{1/2} + ax^{-1/2}) dx$$

$$= \frac{1}{\sqrt{3a}} \left[ \frac{3x^{3/2}}{3/2} + a \frac{x^{1/2}}{1/2} \right]_0^a = \frac{1}{\sqrt{3a}} (4a^{3/2}) = \frac{4a}{\sqrt{3}}.$$




Fig. 6.17

**Example 6.33.** Find the length of one arch of the cycloid

$$x = a(t - \sin t), y = a(1 - \cos t).$$

(P.T.U., 2009 ; V.T.U., 2004)

**Solution.** As a point moves from one end  $O$  to the other end of its first arch, the parameter  $t$  increases from  $0$  to  $2\pi$ . [see Fig. 6.8]

Also  $\frac{dx}{dt} = a(1 - \cos t), \frac{dy}{dt} = a \sin t.$

$$\therefore \text{Length of an arch} = \int_0^{2\pi} \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

$$= \int_0^{2\pi} \sqrt{[a(1 - \cos t)]^2 + (a \sin t)^2} dt = a \int_0^{2\pi} \sqrt{2(1 - \cos t)} dt$$

$$= 2a \int_0^{2\pi} \sin t/2 dt = 2a \left[ -\frac{\cos t/2}{1/2} \right]_0^{2\pi} = 4a[(-\cos \pi) - (-\cos 0)] = 8a.$$

**Example 6.34.** Find the entire length of the cardioid  $r = a(1 + \cos \theta)$ .

(P.T.U., 2010 ; Bhopal, 2008 ; Kurukshetra, 2005)

Also show that the upper half is bisected by  $\theta = \pi/3$ .

(Bhillai, 2005)

**Solution.** The cardioid is symmetrical about the initial line and for its upper half,  $\theta$  increases from  $0$  to  $\pi$  (Fig. 6.18)

Also  $\frac{dr}{d\theta} = -a \sin \theta.$

$$\therefore \text{Length of the curve} = 2 \int_0^{\pi} \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2} d\theta$$

$$\begin{aligned}
 &= 2 \int_0^{\pi} \sqrt{[a(1 + \cos \theta)]^2 + (-a \sin \theta)^2} d\theta = 2a \int_0^{\pi} \sqrt{2(1 + \cos \theta)} d\theta \\
 &= 4a \int_0^{\pi} \cos \theta/2 d\theta = 4a \left[ \frac{\sin \theta/2}{1/2} \right]_0^{\pi} = 8a (\sin \pi/2 - \sin 0) = 8a.
 \end{aligned}$$

$\therefore$  Length of upper half of the curve is  $4a$ . Also length of the arc  $AP$  from  $0$  to  $\pi/3$ .

$$\begin{aligned}
 &= a \int_0^{\pi/3} \sqrt{2(1 + \cos \theta)} d\theta = 2a \int_0^{\pi/3} \cos \theta/2 \cdot d\theta \\
 &= 4a \left[ \sin \theta/2 \right]_0^{\pi/3} = 2a = \text{half the length of upper half of the cardioid.}
 \end{aligned}$$




Fig. 6.18

### PROBLEMS 6.8

- Find the length of the arc of the *semi-cubical parabola*  $ay^2 = x^3$  from the vertex to the ordinate  $x = 5a$ .
- Find the length of the curve (i)  $y = \log \sec x$  from  $x = 0$  to  $x = \pi/3$ . (V.T.U., 2010 S ; P.T.U., 2007)  
(ii)  $y = \log [(e^x - 1)/(e^x + 1)]$  from  $x = 1$  to  $x = 2$ .
- Find the length of the arc of the parabola  $y^2 = 4ax$  (i) from the vertex to one end of the latus-rectum.  
(ii) cut off by the line  $3y = 8x$ . (V.T.U., 2008 S ; Mumbai, 2006)
- Find the perimeter of the loop of the following curves :  
(i)  $ay^2 = x^2(a - x)$  (ii)  $9y^2 = (x - 2)(x - 5)^2$ .
- Find the length of the curve  $y^2 = (2x - 1)^2$  cut off by the line  $x = 4$ . (V.T.U., 2000 S)
- Show that the whole length of the curve  $x^2(a^2 - x^2) = 8a^2y^2$  is  $\pi a\sqrt{2}$ .
- (a) Find the length of an arch of the cycloid  $x = a(\theta + \sin \theta)$ ,  $y = a(1 - \cos \theta)$ .  
(b) By finding the length of the curve show that the curve  $x = a(\theta - \sin \theta)$ ,  $y = a(1 - \cos \theta)$ , is divided in the ratio  $1 : 3$  at  $\theta = 2\pi/3$ . (S.V.T.U., 2009)
- Find the whole length of the curve  $x = a \cos^3 t$ ,  $y = a \sin^3 t$  i.e.,  $x^{2/3} + y^{2/3} = a^{2/3}$ . (V.T.U., 2010 ; Marathwada, 2008 ; Rajasthan, 2006)

Also show that the line  $\theta = \pi/3$  divides the length of this *astroid* in the first quadrant in the ratio  $1 : 3$ .

(Mumbai, 2001)

- Find the length of the loop of the curve  $x = t^2$ ,  $y = t - t^3/3$ . (Mumbai, 2001)
- For the curve  $r = ae^{\theta} \cot \alpha$ , prove that  $s/r = \text{constant}$ ,  $s$  being measured from the origin.
- Find the length of the curve  $\theta = \frac{1}{2} \left( r + \frac{1}{r} \right)$  from  $r = 1$  to  $r = 3$ . (Marathwada, 2008)
- Find the perimeter of the *cardioid*  $r = a(1 - \cos \theta)$ . Also show that the upper half of the curve is bisected by the line  $\theta = 2\pi/3$ .
- Find the whole length of the *lemniscate*  $r^2 = a^2 \cos 2\theta$ .
- Find the length of the parabola  $r(1 + \cos \theta) = 2a$  as cut off by the latus-rectum. (J.N.T.U., 2003)

## 6.12 (1) VOLUMES OF REVOLUTION

(a) **Revolution about x-axis.** The volume of the solid generated by the revolution about the  $x$ -axis, of the area bounded by the curve  $y = f(x)$ , the  $x$ -axis and the ordinates  $x = a$ ,  $x = b$  is

$$\int_a^b \pi y^2 dx.$$

Let  $AB$  be the curve  $y = f(x)$  between the ordinates  $LA(x = a)$  and  $MB(x = b)$ .

Let  $P(x, y)$ ,  $P'(x + \delta x, y + \delta y)$  be two neighbouring points on the curve and  $NP$ ,  $N'P'$  be their respective ordinates (Fig. 6.19).

Let the volume of the solid generated by the revolution about  $x$ -axis of the area  $ALNP$  be  $V$ , which is clearly a function of  $x$ . Then the volume of the solid generated by the revolution of the area  $PNN'P'$  is  $\delta V$ . Complete the rectangles  $PN'$  and  $P'N$ .




Fig. 6.19



The  $\delta V$  lies between the volumes of the right circular cylinders generated by the revolution of rectangles  $PN'$  and  $P'N$ ,

i.e.,  $\delta V$  lies between  $\pi y^2 \delta x$  and  $\pi(y + \delta y)^2 \delta x$ .

$\therefore \frac{\delta V}{\delta x}$  lies between  $\pi y^2$  and  $\pi(y + \delta y)^2$ .

Now taking limits as  $P' \rightarrow P$ , i.e.,  $\delta x \rightarrow 0$  (and  $\delta y \rightarrow 0$ ),  $\frac{dV}{dx} = \pi y^2$

$\therefore \int_a^b \frac{dV}{dx} dx = \int_a^b \pi y^2 dx$  or  $V \Big|_{x=a}^b = \int_a^b \pi y^2 dx$

or (value of  $V$  for  $x = b$ ) – (value of  $V$  for  $x = a$ )

i.e., volume of the solid obtained by the revolution of the area  $ALMB = \int_a^b \pi y^2 dx$ .

**Example 6.35.** Find the volume of a sphere of radius  $a$ .

(S.V.T.U., 2007)

**Solution.** Let the sphere be generated by the revolution of the semi-circle  $ABC$ , of radius  $a$  about its diameter  $CA$  (Fig. 6.20)

Taking  $CA$  as the  $x$ -axis and its mid-point  $O$  as the origin, the equation of the circle  $ABC$  is  $x^2 + y^2 = a^2$ .

$\therefore$  Volume of the sphere = 2 (volume of the solid generated by the revolution about  $x$ -axis of the quadrant  $OAB$ )

$$\begin{aligned} &= 2 \int_0^a \pi y^2 dx = 2\pi \int_0^a (a^2 - x^2) dx \\ &= 2\pi \left[ a^2 x - \frac{x^3}{3} \right]_0^a = 2\pi \left[ a^3 - \frac{a^3}{3} - (0 - 0) \right] = \frac{4}{3} \pi a^3. \end{aligned}$$




Fig. 6.20

**Example 6.36.** Find the volume formed by the revolution of loop of the curve  $y^2(a+x) = x^2(3a-x)$ , about the  $x$ -axis.

(Marathwada, 2008)

**Solution.** The curve is symmetrical about the  $x$ -axis, and for the upper half of its loop  $x$  varies from 0 to  $3a$  (Fig. 6.21)

$$\begin{aligned} \therefore \text{Volume of the loop} &= \int_0^{3a} \pi y^2 dx = \pi \int_0^{3a} \frac{x^2(3a-x)}{a+x} dx \\ &= \pi \int_0^{3a} \frac{-x^3 + 3ax^2}{x+a} dx \end{aligned}$$

[Divide the numerator by the denominator]

$$\begin{aligned} &= \pi \int_0^{3a} \left[ -x^2 + 4ax - 4a^2 + \frac{4a^3}{x+a} \right] dx = \pi \left[ -\frac{x^3}{3} + 4a \cdot \frac{x^2}{2} - 4a^2 x + 4a^3 \log(x+a) \right]_0^{3a} \\ &= \pi \left[ \frac{-27a^3}{3} + 2a \cdot 9a^2 - 4a^2 \cdot 3a + 4a^3 \log 4a - (4a^3 \log a) \right] \\ &= \pi a^3 (-3 + 4 \log 4) = \pi a^3 (8 \log 2 - 3). \end{aligned}$$




Fig. 6.21

**Example 6.37.** Prove that the volume of the reel formed by the revolution of the cycloid  $x = a(\theta + \sin \theta)$ ,  $y = a(1 - \cos \theta)$  about the tangent at the vertex is  $\pi^2 a^3$ .

(V.T.U., 2003)

**Solution.** The arch  $AOB$  of the cycloid is symmetrical about the  $y$ -axis and the tangent at the vertex is the  $x$ -axis. For half the cycloid  $OA$ ,  $\theta$  varies from 0 to  $\pi$ . (Fig. 4.31).

Hence the required volume

$$= 2 \int_{\theta=0}^{\theta=\pi} \pi y^2 dx = 2\pi \int_0^{\pi} a^2(1 - \cos \theta)^2 \cdot a(1 + \cos \theta) d\theta$$



$$\begin{aligned}
 &= 2\pi a^3 \int_0^\pi (2 \sin^2 \theta/2) \cdot (2 \cos^2 \theta/2) d\theta \\
 &= 16\pi a^3 \int_0^\pi \sin^4 \theta/2 \cdot \cos^2 \theta/2 \cdot d\theta \quad [\text{Put } \theta/2 = \phi, d\theta = 2d\phi] \\
 &= 32\pi a^3 \int_0^{\pi/2} \sin^4 \phi \cos^2 \phi d\phi = 32\pi a^3 \cdot \frac{3 \cdot 1 \times 1}{6 \cdot 4 \cdot 2} \cdot \frac{\pi}{2} = \pi^2 a^3.
 \end{aligned}$$

**Example 6.38.** Find the volume of the solid formed by revolving about  $x$ -axis, the area enclosed by the parabola  $y^2 = 4ax$ , its evolute  $27ay^2 = 4(x - 2a)^3$  and the  $x$ -axis.

**Solution.** The curve  $27ay^2 = 4(x - 2a)^3$  ... (i) is symmetrical about  $x$ -axis and is a semi-cubical parabola with vertex at  $A(2a, 0)$ . The parabola  $y^2 = 4ax$  and (i) intersect at  $B$  and  $C$  where  $27a(4ax) = 4(x - 2a)^3$  or  $x^3 - 6ax^2 - 15a^2x - 8a^3 = 0$  which gives  $x = -a, -a, 8a$ . Since  $x$  is not negative, therefore we have  $x = 8a$  (Fig. 6.22).

$\therefore$  Required volume = Volume obtained by revolving the shaded area  $OAB$  about  $x$ -axis = Vol. obtained by revolving area  $OMBO$  - Vol. obtained by revolving area  $ADBA$

$$\begin{aligned}
 &= \int_0^{8a} \pi y^2 (= 4ax) dx - \int_{2a}^{8a} \pi y^2 [\text{for (i)}] dx \\
 &= 4a\pi \left[ \frac{x^2}{2} \right]_0^{8a} - \frac{4\pi}{27a} \int_{2a}^{8a} (x - 2a)^3 dx \\
 &= 128\pi a^3 - \frac{4\pi}{27a} \left[ \frac{(x - 2a)^4}{4} \right]_{2a}^{8a} \\
 &= 128\pi a^3 - \frac{\pi}{27a} (6a)^4 = 80\pi a^3.
 \end{aligned}$$

(b) **Revolution about the  $y$ -axis.** Interchanging  $x$  and  $y$  in the above formula, we see that the volume of the solid generated by the revolution about  $y$ -axis, of the area, bounded by the curve  $x = f(y)$ , the  $y$ -axis and the abscissae  $y = a, y = b$  is

$$\int_a^b \pi x^2 dy.$$

**Example 6.39.** Find the volume of the reel-shaped solid formed by the revolution about the  $y$ -axis, of the part of the parabola  $y^2 = 4ax$  cut off by the latus-rectum. (Rohtak, 2003)

**Solution.** Given parabola is  $x = y^2/4a$ .

Let  $A$  be the vertex and  $L$  one extremity of the latus-rectum. For the arc  $AL$ ,  $y$  varies from 0 to  $2a$  (Fig. 6.23).

$\therefore$  required volume = 2 (volume generated by the revolution about the  $y$ -axis of the area  $ALC$ )

$$= 2 \int_0^{2a} \pi x^2 dy = 2\pi \int_0^{2a} \frac{y^4}{16a^2} \cdot dy = \frac{\pi}{8a^2} \left[ \frac{y^5}{5} \right]_0^{2a} = \frac{\pi}{40a^2} (32a^5 - 0) = \frac{4\pi a^3}{5}.$$

(c) **Revolution about any axis.** The volume of the solid generated by the revolution about any axis  $LM$  of the area bounded by the curve  $AB$ , the axis  $LM$  and the perpendiculars  $AL, BM$  on the axis, is

$$\int_{OL}^{OM} \pi (PN)^2 d(ON)$$

where  $O$  is a fixed point in  $LM$  and  $PN$  is perpendicular from any point  $P$  of the curve  $AB$  on  $LM$ .

With  $O$  as origin, take  $OLM$  as the  $x$ -axis and  $OY$ , perpendicular to it as the  $y$ -axis (Fig. 6.24).




Fig. 6.22




Fig. 6.23




Fig. 6.24

Let the coordinates of  $P$  be  $(x, y)$  so that  $x = ON, y = NP$

If  $OL = a, OM = b$ , then required volume  $= \int_a^b \pi y^2 dx = \int_{OL}^{OM} \pi(PN)^2 d(ON)$ .

**Example 6.40.** Find the volume of the solid obtained by revolving the cissoid  $y^2(2a - x) = x^3$  about its asymptote. (V.T.U., 2000)

**Solution.** Given curve is  $y = \frac{x^3}{2a - x}$  ... (i)

It is symmetrical about  $x$ -axis and the asymptote is  $x = 2a$ . (See Fig. 4.23). If  $P(x, y)$  be any point on it and  $PN$  is perpendicular on the asymptote  $AN$  then  $PN = 2a - x$  and

$$AN = y = \frac{x^{3/2}}{\sqrt{(2a - x)}} \quad \text{[From (i)]}$$

$$\therefore d(AN) = dy = \frac{\sqrt{(2a - x)}(3/2)\sqrt{x} - x^{3/2} \cdot \frac{1}{2}(2a - x)^{-1/2}(-1)}{2a - x} dx$$

$$= \frac{3\sqrt{x}(2a - x) + x^{3/2}}{2(2a - x)^{3/2}} dx = \frac{3ax^{1/2} - x^{3/2}}{(2a - x)^{3/2}} dx$$

$$\therefore \text{Required volume} = 2 \int_{x=0}^{x=2a} \pi(PN)^2 d(AN) = 2\pi \int_0^{2a} (2a - x)^2 \cdot \frac{3ax^{1/2} - x^{3/2}}{(2a - x)^{3/2}} \cdot dx$$

$$= 2\pi \int_0^{2a} \sqrt{(2a - x)}(3a - x)\sqrt{x} dx \quad \left[ \begin{array}{l} \text{Put } x = 2a \sin^2 \theta \\ \text{then } dx = 4a \sin \theta \cos \theta d\theta \end{array} \right]$$

$$= 2\pi \int_0^{\pi/2} \sqrt{(2a)} \cos \theta (3a - 2a \sin^2 \theta) \sqrt{(2a)} \sin \theta \cdot 4a \sin \theta \cos \theta d\theta$$

$$= 16\pi a^3 \left[ 3 \int_0^{\pi/2} \sin^2 \theta \cos^2 \theta d\theta - 2 \int_0^{\pi/2} \sin^4 \theta \cos^2 \theta d\theta \right]$$

$$= 16\pi a^3 \left[ 3 \cdot \frac{1 \times 1}{4 \cdot 2} \cdot \frac{\pi}{2} - 2 \cdot \frac{3 \cdot 1 \times 1}{6 \cdot 4 \cdot 2} \cdot \frac{\pi}{2} \right] = 2\pi^2 a^3.$$

**(2) Volumes of revolution (polar curves).** The volume of the solid generated by the revolution of the area bounded by the curve  $r = f(\theta)$  and the radii vectors  $\theta = \alpha, \theta = \beta$  (Fig. 6.25)

$$(a) \text{ about the initial line } OX (\theta = 0) = \int_{\alpha}^{\beta} \frac{2\pi}{3} r^3 \sin \theta d\theta$$

$$(b) \text{ about the line } OY (\theta = \pi/2) = \int_{\alpha}^{\beta} \frac{2\pi}{3} r^3 \cos \theta d\theta.$$




Fig. 6.25

**Example 6.41.** Find the volume of the solid generated by the revolution of the cardioid  $r = a(1 + \cos \theta)$  about the initial line. (V.T.U., 2010 ; Kurukshetra, 2009 S)

**Solution.** The cardioid is symmetrical about the initial line and for its upper half  $\theta$  varies from 0 to  $\pi$ . [Fig. 6.18].

$$\therefore \text{required volume} = \int_0^{\pi} \frac{2}{3} \pi r^3 \sin \theta d\theta = \frac{2\pi}{3} \int_0^{\pi} a^3 (1 + \cos \theta)^3 \sin \theta d\theta$$

$$= -\frac{2\pi a^3}{3} \int_0^{\pi} (1 + \cos \theta)^3 \cdot (-\sin \theta) d\theta = -\frac{2\pi a^3}{3} \left[ \frac{(1 + \cos \theta)^4}{4} \right]_0^{\pi} = -\frac{\pi a^3}{6} [0 - 16] = \frac{8}{3} \pi a^3.$$

**Example 6.42.** Find the volume of the solid generated by revolving the lemniscate  $r^2 = a^2 \cos 2\theta$  about the line  $\theta = \pi/2$ . (V.T.U., 2006)

**Solution.** The curve is symmetrical about the pole. For the upper half of the R.H.S. loop,  $\theta$  varies from 0 to  $\pi/4$ . (Fig. 4.34).



∴ required volume = 2(volume generated by the half loop in the first quadrant)

$$\begin{aligned} &= 2 \int_0^{\pi/4} \frac{2}{3} \pi r^3 \cos \theta \, d\theta = \frac{4\pi}{3} \int_0^{\pi/4} a^3 (\cos 2\theta)^{3/2} \cos \theta \, d\theta \quad [\because r = a (\cos 2\theta)^{1/2}] \\ &= \frac{4\pi a^3}{3} \int_0^{\pi/4} (1 - 2 \sin^2 \theta)^{3/2} \cos \theta \, d\theta \quad \left[ \begin{array}{l} \text{Put } \sqrt{2} \sin \theta = \sin \phi \\ \therefore \sqrt{2} \cos \theta \, d\theta = \cos \phi \, d\phi \end{array} \right] \\ &= \frac{4\pi a^3}{3} \int_0^{\pi/2} (1 - \sin^2 \phi)^{3/2} \cdot \frac{1}{\sqrt{2}} \cos \phi \, d\phi = \frac{4\pi a^3}{3\sqrt{2}} \int_0^{\pi/2} \cos^4 \phi \, d\phi = \frac{4\pi}{3\sqrt{2}} a^3 \frac{3 \cdot 1}{4 \cdot 2} \cdot \frac{\pi}{2} = \frac{\pi a^3}{4\sqrt{2}} \end{aligned}$$

### PROBLEMS 6.9

- Find the volume generated by the revolution of the area bounded by  $x$ -axis, the catenary  $y = c \cosh x/c$  and the ordinates  $x = \pm c$ , about the axis of  $x$ .
- Find the volume of a spherical segment of height  $h$  cut off from a sphere of radius  $a$ .
- Find the volume generated by revolving the portion of the parabola  $y^2 = 4ax$  cut off by its latus-rectum about the axis of the parabola. (V.T.U., 2009)
- Find the volume generated by revolving the area bounded by the curve  $\sqrt{x} + \sqrt{y} = \sqrt{a}$ ,  $x = 0$ ,  $y = 0$  about the  $x$ -axis.
- Find the volume of the solid generated by revolving the ellipse  $x^2/a^2 + y^2/b^2 = 1$ .  
(i) about the major axis. (Bhopal, 2002 S) (ii) about the minor axis. (Bhillai, 2005)
- Obtain the volume of the frustrum of a right circular cone whose lower base has radius  $R$ , upper base is of radius  $r$  and altitude is  $h$ .
- Find the volume generated by the revolution of the curve  $27ay^2 = 4(x - 2a)^3$  about the  $x$ -axis.
- Find the volume of the solid formed by the revolution, about the  $x$ -axis, of the loop of the curve :  
(i)  $y^2(a - x) = x^2(a + x)$  (ii)  $2ay^2 = x(x - a)^2$  (iii)  $y^2 = x(2x - 1)^2$ .
- Find the volume obtained by revolving one arch of the cycloid  
(i)  $x = a(t - \sin t)$ ,  $y = a(1 - \cos t)$ , about its base. (Kurukshetra, 2006 ; V.T.U., 2005)  
(ii)  $x = a(\theta + \sin \theta)$ ,  $y = a(1 + \cos \theta)$ , about the  $x$ -axis.
- Find the volume of the spindle-shaped solid generated by the revolution of the astroid  $x^{2/3} + y^{2/3} = a^{2/3}$  about the  $x$ -axis. (P.T.U., 2010 ; S.V.T.U., 2008)
- Find the volume of the solid formed by the revolution, about the  $y$ -axis, of the area enclosed by the curve  $xy^2 = 4a^2(2a - x)$  and its asymptote. (V.T.U., 2006)
- Prove that the volume of the solid formed by the revolution of the curve  $(a^2 + x^2) = a^3$ , about its asymptote is  $\frac{1}{2} \pi^2 a^3$ .
- Find the volume generated by the revolution about the initial line of  
(i)  $r = 2a \cos \theta$  (ii)  $r = a(1 - \cos \theta)$ . (P.T.U., 2006)
- Determine the volume of the solid obtained by revolving the lemniscate  $r = a + b \cos \theta$  ( $a > b$ ) about the initial line. (Gorakhpur, 1999)
- Find the volume of the solid formed by revolving a loop of the lemniscate  $r^2 = a^2 \cos 2\theta$  about the initial line. (J.N.T.U., 2003 ; Delhi, 2002)

### 6.13 SURFACE AREAS OF REVOLUTION

(a) **Revolution about  $x$ -axis.** The surface area of the solid generated by the revolution about  $x$ -axis, of the arc of the curve  $y = f(x)$  from  $x = a$  to  $x = b$ , is

$$\int_{x=a}^{x=b} 2\pi y \, ds.$$

Let  $AB$  be the curve  $y = f(x)$  between the ordinates  $LA$  ( $x = a$ ) and  $MB$  ( $x = b$ ). Let  $P(x, y)$ ,  $P'(x + \delta x, y + \delta y)$  be two neighbouring points on the curve and  $NP$ ,  $N'P'$  be their respective ordinates (Fig. 6.19).

Let the arc  $AP = s$  so that arc  $PP' = \delta s$ . Let the surface-area generated by the revolution about  $x$ -axis of the arc  $AP$  be  $S$  and that generated by the revolution of the arc  $PP'$  be  $\delta S$ .

Since  $\delta s$  is small, the surface area  $\delta S$  may be regarded as lying between the curved surfaces of the right cylinders of radii  $PN$  and  $P'N'$  and of same thickness  $\delta s$ .

Thus  $\delta S$  lies between  $2\pi y \delta s$  and  $2\pi (y + \delta y) \delta s$

$\therefore \frac{\delta S}{\delta s}$  lies between  $2\pi y$  and  $2\pi (y + \delta y)$

Taking limits as  $P' \rightarrow P$ , i.e.,  $\delta s \rightarrow 0$  and  $\delta y \rightarrow 0$ ,  $dS/dx = 2\pi y$

$$\therefore \int_{x=a}^{x=b} \frac{dS}{ds} ds = \int_{x=a}^{x=b} 2\pi y ds \quad \text{or} \quad S \Big|_{x=a}^{x=b} = \int_{x=a}^{x=b} 2\pi y ds$$

or (value of  $S$  for  $x = b$ ) – (value of  $S$  for  $x = a$ ) =  $\int_{x=a}^{x=b} 2\pi y dx$

or surface area generated by the revolution of the arc  $AB$  – 0 =  $\int_{x=a}^{x=b} 2\pi y ds$ .

Hence, the required surface area =  $\int_{x=a}^{x=b} 2\pi y ds$ .

**Obs. Practical forms of the formula  $S = \int 2\pi y ds$ .**

(i) Cartesian form [for the curve  $y = f(x)$ ]

$$S = \int 2\pi y \frac{ds}{dx} dx, \text{ where } \frac{ds}{dx} = \sqrt{1 + \left(\frac{dy}{dx}\right)^2}$$

(ii) Parametric form [for the curve  $x = f(t)$ ,  $y = \phi(t)$ ]

$$S = \int 2\pi y \frac{ds}{dt} dt, \text{ where } \frac{ds}{dt} = \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2}$$

(iii) Polar form [for the curve  $r = f(\theta)$ ]

$$S = \int 2\pi y \frac{ds}{d\theta} d\theta, \text{ where } y = r \sin \theta, \frac{ds}{d\theta} = \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2}.$$

**Example 6.43.** Find the surface of the solid formed by revolving the cardioid  $r = a(1 + \cos \theta)$  about the initial line.  
(V.T.U., 2009 ; Rajasthan, 2006 ; J.N.T.U., 2003)

**Solution.** The cardioid is symmetrical about the initial line and for its upper half,  $\theta$  varies from 0 to  $\pi$  (Fig. 6.18).

Also 
$$\frac{ds}{d\theta} = \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2} = \sqrt{a^2(1 + \cos \theta)^2 + a^2 \sin^2 \theta}$$

$$= a \sqrt{2(1 + \cos \theta)} = a \sqrt{2 \cdot 2 \cos^2 \theta / 2} = 2a \cos \theta / 2$$

$\therefore$  required surface =  $\int_0^\pi 2\pi y \frac{ds}{d\theta} d\theta = 2\pi \int_0^\pi r \sin \theta \cdot 2a \cos \theta / 2 d\theta$

$$= 4\pi a \int_0^\pi a(1 + \cos \theta) \sin \theta \cdot \cos \theta / 2 d\theta = 4\pi a^2 \int_0^\pi 2 \cos^2 \frac{\theta}{2} \cdot 2 \sin \frac{\theta}{2} \cos \frac{\theta}{2} \cdot \cos \frac{\theta}{2} d\theta$$

$$= 16\pi a^2 \int_0^\pi \cos^4 \frac{\theta}{2} \sin \frac{\theta}{2} d\theta = 16\pi a^2 (-2) \int_0^\pi \cos^4 \frac{\theta}{2} \left(-\sin \frac{\theta}{2} \cdot \frac{1}{2}\right) d\theta$$

$$= -32\pi a^2 \left| \frac{\cos^5 \theta / 2}{5} \right|_0^\pi = \frac{-32\pi a^2}{5} (0 - 1) = \frac{32\pi a^2}{5}.$$

(b) **Revolution about y-axis.** Interchanging  $x$  and  $y$  in the above formula, we see that the surface of the solid generated by the revolution about y-axis, of the arc of the curve  $x = f(y)$  from  $y = a$  to  $y = b$  is

$$\int_{y=a}^{y=b} 2\pi x ds.$$

**Example 6.44.** Find the surface area of the solid generated by the revolution of the astroid  $x = a \cos^3 t$ ,  $y = a \sin^3 t$ , about the y-axis.



**Solution.** The astroid is symmetrical about the  $x$ -axis, and for its portion in the first quadrant  $t$  varies from 0 to  $\pi/2$ . (Fig. 4.29).

Also  $\frac{dx}{dt} = -3a \cos^2 t \sin t$ ,  $\frac{dy}{dt} = 3a \sin^2 t \cos t$ .

$$\frac{ds}{dt} = \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} = \sqrt{[9a^2 \cos^4 t \sin^2 t + 9a^2 \sin^4 t \cos^2 t]}$$

$$= 3a \sin t \cos t \sqrt{(\cos^2 t + \sin^2 t)} = 3a \sin t \cos t$$

$$\therefore \text{required surface} = 2 \int_0^{\pi/2} 2\pi x \frac{ds}{dt} \cdot dt = 4\pi \int_0^{\pi/2} a \cos^3 t \cdot 3a \sin t \cos t \, dt$$

$$= 12\pi a^2 \int_0^{\pi/2} \sin t \cos^4 t \, dt = 12\pi a^2 \frac{3 \cdot 1}{5 \cdot 3 \cdot 1} = \frac{12\pi a^2}{5}.$$

### PROBLEMS 6.10

- Find the area of the surface generated by revolving the arc of the catenary  $y = c \cosh x/c$  from  $x = 0$  to  $x = c$  about the  $x$ -axis.
- Find the area of the surface formed by the revolution of  $y^2 = 4ax$  about its axis, by the arc from the vertex to one end of the latus-rectum.
- Find the surface of the solid generated by the revolution of the ellipse  $x^2/a^2 + y^2/b^2 = 1$  about the  $x$ -axis.  
(Raipur, 2005 ; Bhopal, 2002 S)
- Find the volume and surface of the *right circular cone* formed by the revolution of a right-angled triangle about a side which contains the right angle.
- Obtain the surface area of a *sphere* of radius  $a$ .
- Show that the surface area of the solid generated by the revolution of the curve  $x = a \cos^3 t$ ,  $y = a \sin^3 t$  about the  $x$ -axis, is  $12\pi^2/5$ .
- The arc of the curve  $x^{2/3} + y^{2/3} = a^{2/3}$  in the first quadrant revolves about  $x$ -axis. Show that the area of the surface generated is  $6\pi a^2/5$ .
- Find the surface area of the solid generated by revolving the *cycloid*  $x = a(t - \sin t)$ ,  $y = a(1 - \cos t)$  about the base.  
(Marathwada, 2008 ; Cochin, 2005 ; Kurukshetra, 2005)
- Find the surface area of the solid got by revolving the arch of the *cycloid*  
 $x = a(\theta + \sin \theta)$ ,  $y = a(1 + \cos \theta)$  about the base.  
(V.T.U., 2010 S)
- Prove that the surface and volume of the solid generated by the revolution about the  $x$ -axis, of the loop of the curve  $x = t^2$ ,  $y = t - t^3/3$ , [or  $9y^2 = x(x - 3)^2$ ], are respectively  $3\pi$  and  $3\pi/4$ .
- Prove that the surface of the solid generated by the revolution of the *tractrix*  $x = a \cos t + \frac{a}{2} \log \tan^2 t/2$ ,  $y = a \sin t$ , about  $x$ -axis is  $4\pi a^2$ .
- Find the surface area of the solid of revolution of the curve  $r = 2a \cos \theta$  about the initial line.  
(V.T.U., 2009)
- Find the surface of the solid generated by the revolution of the *cardioid*  $r = a(1 - \cos \theta)$  about the initial line.
- Find the surface of the solid generated by the revolution of the *lemniscate*  $r^2 = a^2 \cos 2\theta$  about the initial line.  
(V.T.U., 2005)
- The part of *parabola*  $y^2 = 4ax$  cut off by the latus-rectum revolves about the tangent at the vertex. Find the curved surface of the reel thus formed.

### 6.14 OBJECTIVE TYPE OF QUESTIONS

### PROBLEMS 6.11

Choose the correct answer or fill up the blanks in the following problems :

- If  $f(x) = f(2a - x)$ , then  $\int_0^{2a} f(x) \, dx$  is equal to



- (a)  $\int_a^0 f(2a-x) dx$  (b)  $2 \int_0^a f(x) dx$  (c)  $-2 \int_0^a f(x) dx$  (d) 0.
2.  $\int_0^{\pi/2} \frac{\sqrt{\sin x}}{\sqrt{\sin x} + \sqrt{\cos x}} dx$  is equal to  
 (a) 0 (b) 1 (c)  $\frac{\pi}{4}$  (d)  $\frac{\pi}{2}$ .
3. The value of definite integral  $\int_{-a}^a |x| dx$  is equal to  
 (a)  $a$  (b)  $a^2$  (c) 0 (d)  $2a$ .
4.  $\lim_{n \rightarrow \infty} \left[ \frac{n}{n^2} + \frac{n}{n^2+1^2} + \frac{n}{n^2+2^2} + \dots + \frac{n}{n^2+(n-1)^2} \right]$  is equal to  
 (a)  $-\frac{\pi}{4}$  (b) 0 (c)  $\frac{\pi}{4}$  (d)  $\frac{\pi}{3}$ .
5.  $\int_0^{\pi/2} \frac{\cos 2x}{\cos x + \sin x} dx$  equals  
 (a) -1 (b) 0 (c) 1 (d) 2.
6.  $\lim_{n \rightarrow \infty} \left( \frac{1}{n} + \frac{1}{n+1} + \frac{1}{n+2} + \dots + \frac{1}{3n} \right)$  equals  
 (a)  $\log_e 2$  (b)  $2 \log_e 2$  (c)  $\log_e 3$  (d)  $2 \log_e 3$ .
7.  $\int_0^{\pi} \sin^5 \left( \frac{x}{2} \right) dx$  is equal to  
 (a)  $\frac{16}{15}$  (b)  $\frac{15}{16} \pi$  (c)  $\frac{16}{15} \pi^2$  (d)  $\frac{15}{16}$ .
8.  $\int_0^{\pi/2} \sin^{99} x \cos x dx$  is equal to  
 (a)  $\frac{1}{99}$  (b)  $\frac{\pi}{100}$  (c)  $\frac{99}{100}$  (d) None of these. (V.T.U., 2009)
9. The value of  $\int_{-\pi/2}^{\pi/2} \cos^7 x dx$  is  
 (a)  $\frac{32\pi}{35}$  (b)  $\frac{32}{35}$  (c) zero.
10. The length of the arc of the equiangular spiral  $r = ae^{\theta \cot \alpha}$  between the points for which the radii vectors are  $r_1$  and  $r_2$  is  
 (a)  $(r_2 - r_1) \operatorname{cosec} \alpha$  (b)  $(r_2 - r_1) \cos \alpha$  (c)  $(r_2 - r_1) \sin \alpha$  (d)  $(r_2 - r_1) \sec \alpha$ .
11. The area of the region in the first quadrant bounded by the y-axis and the curves  $y = \sin x$  and  $y = \cos x$  is  
 (a)  $\sqrt{2}$  (b)  $\sqrt{2} + 1$  (c)  $\sqrt{2} - 1$  (d)  $2\sqrt{2} - 1$ .
12. The value of  $\int_0^1 x^{3/2} (1-x)^{3/2} dx$  is  
 (a)  $\pi/32$  (b)  $-\pi/32$  (c)  $3\pi/128$  (d)  $-3\pi/128$ . (V.T.U., 2010)
13. The entire length of the cardioid  $r = 5(1 + \cos \theta)$  is  
 (a) 40 (b) 30 (c) 20 (d) 5. (V.T.U., 2009)
14. The area of the cardioid  $r = a(1 - \cos \theta)$  is .....
15. If  $S_1$  and  $S_2$  are surface areas of the solids generated by revolving the ellipses  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  and  $\frac{x^2}{b^2} + \frac{y^2}{a^2} = 1$  about the y-axis, then  
 (a)  $S_1 > S_2$  (b)  $S_1 < S_2$  (c)  $S_1 = S_2$  (d) can't predict.
16. The area of the loop of the curve  $r = a \sin 3\theta$  is .....
17. If  $I_n = \int_0^{\pi/4} \tan^n \theta d\theta$ , then  $n(I_{n-1} + I_{n+1}) = \dots\dots\dots$  18.  $\int_0^2 x^3 \sqrt{2x-x^2} dx = \dots\dots\dots$



19.  $\int_0^{\pi/2} \sin 2\theta \log \tan \theta d\theta$  is equal to  
 (a) 1 (b) -1 (c) 0 (d)  $\pi/2$ .
20. The area of the loop of the folium of Descartes  $x^3 + y^3 - 3xy = 0$  is  
 (a)  $\pi$  (b)  $\pi/2$  (c) 1.5 (d) 3.
21. The volume of the frustrum of a right circular cone whose lower base has radius  $r_1$  and upper base has radius  $r_2$  and altitude is  $h = \dots\dots$
22. The length of the arc of the curve  $y = \log \sec x$  from  $x = 0$  to  $x = \pi/4$  is  
 (a)  $\log_e 2$  (b)  $\log_e 3$  (c)  $\log_e (1 + \sqrt{2})$  (d)  $\log_e (1 + \sqrt{3})$ . (Bhopal, 2008)
23. If  $v_1 =$  volume of the solid generated by revolving the area included between  $x$ -axis and  $x^2 + y^2 = a^2$  about  $x$ -axis ;  
 $v_2 =$  volume of the solid generated by revolving the entire area of the circle  $x^2 + y^2 = a^2$  about  $x$ -axis, then  
 (a)  $v_1 = v_2$  (b)  $v_2 = 2v_1$  (c)  $v_2 = 4v_1$  (d)  $v_2 = 16v_1$ .
24. If  $f(r, \theta) = f(-r, \theta)$ , then the curve is symmetrical about the .....  
 (a) initial line (b) pole (c) origin (d) tangential line. (V.T.U., 2010)
25. The volume generated by the revolution of the curve  $y = a^3 (a^2 + x^2)^{-1}$  about its asymptote is  
 (a)  $\pi^2 a^3/2$  (b)  $\pi a^3/2$  (c)  $\pi a^2/2$  (d)  $\pi a/2$ . (V.T.U., 2010)

# Multiple Integrals and Beta, Gamma Functions

1. Double integrals. 2. Change of order of integration. 3. Double integrals in Polar coordinates. 4. Areas enclosed by plane curves. 5. Triple integrals. 6. Volume of solids. 7. Change of variables. 8. Area of a curved surface. 9. Calculation of mass. 10. Centre of gravity. 11. Centre of pressure. 12. Moment of inertia. 13. Product of inertia ; Principal axes. 14. Beta function. 15. Gamma function. 16. Relation between beta and gamma functions. 17. Elliptic integrals. 18. Error function or Probability integral. 19. Objective Type of Questions.

## 7.1 DOUBLE INTEGRALS

The definite integral  $\int_a^b f(x) dx$  is defined as the limit of the sum

$$f(x_1) \delta x_1 + f(x_2) \delta x_2 + \dots + f(x_n) \delta x_n,$$

where  $n \rightarrow \infty$  and each of the lengths  $\delta x_1, \delta x_2, \dots$  tends to zero. A double integral is its counterpart in two dimensions.

Consider a function  $f(x, y)$  of the independent variables  $x, y$  defined at each point in the finite region  $R$  of the  $xy$ -plane. Divide  $R$  into  $n$  elementary areas  $\delta A_1, \delta A_2, \dots, \delta A_n$ . Let  $(x_r, y_r)$  be any point within the  $r$ th elementary area  $\delta A_r$ . Consider the sum

$$f(x_1, y_1) \delta A_1 + f(x_2, y_2) \delta A_2 + \dots + f(x_n, y_n) \delta A_n, \text{ i.e., } \sum_{r=1}^n f(x_r, y_r) \delta A_r$$

The limit of this sum, if it exists, as the number of sub-divisions increases indefinitely and area of each sub-division decreases to zero, is defined as the *double integral of  $f(x, y)$  over the region  $R$*  and is written as

$$\iint_R f(x, y) dA.$$

Thus

$$\iint_R f(x, y) dA = \lim_{\substack{n \rightarrow \infty \\ \delta A \rightarrow 0}} \sum_{r=1}^n f(x_r, y_r) \delta A_r \quad \dots(1)$$

The utility of double integrals would be limited if it were required to take limit of sums to evaluate them. However, there is another method of evaluating double integrals by successive single integrations.

For purpose of evaluation, (1) is expressed as the repeated integral  $\int_{x_1}^{x_2} \int_{y_1}^{y_2} f(x, y) dx dy$ .

Its value is found as follows :

(i) When  $y_1, y_2$  are functions of  $x$  and  $x_1, x_2$  are constants,  $f(x, y)$  is first integrated w.r.t.  $y$  keeping  $x$  fixed between limits  $y_1, y_2$  and then resulting expression is integrated w.r.t.  $x$  within the limits  $x_1, x_2$  i.e.,

$$I_1 = \int_{x_1}^{x_2} \left[ \int_{y_1}^{y_2} f(x, y) dy \right] dx$$

where integration is carried from the inner to the outer rectangle.



Figure 7.1 illustrates this process. Here  $AB$  and  $CD$  are the two curves whose equations are  $y_1 = f_1(x)$  and  $y_2 = f_2(x)$ .  $PQ$  is a vertical strip of width  $dx$ .

Then the inner rectangle integral means that the integration is along one edge of the strip  $PQ$  from  $P$  to  $Q$  ( $x$  remaining constant), while the outer rectangle integral corresponds to the sliding of the edge from  $AC$  to  $BD$ .

Thus the whole region of integration is the area  $ABDC$ .




Fig. 7.1




Fig. 7.2

(ii) When  $x_1, x_2$  are functions of  $y$  and  $y_1, y_2$  are constants,  $f(x, y)$  is first integrated w.r.t.  $x$  keeping  $y$  fixed, within the limits  $x_1, x_2$  and the resulting expression is integrated w.r.t.  $y$  between the limits  $y_1, y_2$ , i.e.,

$$I_2 = \int_{y_1}^{y_2} \left[ \int_{x_1}^{x_2} f(x, y) dx \right] dy \quad \text{which is geometrically illustrated by Fig. 7.2.}$$

Here  $AB$  and  $CD$  are the curves  $x_1 = f_1(y)$  and  $x_2 = f_2(y)$ .  $PQ$  is a horizontal strip of width  $dy$ .

Then inner rectangle indicates that the integration is along one edge of this strip from  $P$  to  $Q$  while the outer rectangle corresponds to the sliding of this edge from  $AC$  to  $BD$ .

Thus the whole region of integration is the area  $ABDC$ .

(iii) When both pairs of limits are constants, the region of integration is the rectangle  $ABDC$  (Fig. 7.3).

In  $I_1$ , we integrate along the vertical strip  $PQ$  and then slide it from  $AC$  to  $BD$ .

In  $I_2$ , we integrate along the horizontal strip  $P'Q'$  and then slide it from  $AB$  to  $CD$ .

Here obviously  $I_1 = I_2$ .

Thus for constant limits, it hardly matters whether we first integrate w.r.t.  $x$  and then w.r.t.  $y$  or vice versa.




Fig. 7.3

**Example 7.1.** Evaluate  $\int_0^5 \int_0^{x^2} x(x^2 + y^2) dx dy$ .

$$\begin{aligned} \text{Solution.} \quad I &= \int_0^5 dx \int_0^{x^2} (x^3 + xy^3) dy = \int_0^5 \left[ x^3 y + x \cdot \frac{y^3}{3} \right]_0^{x^2} dx = \int_0^5 \left[ x^3 \cdot x^2 + x \cdot \frac{y^6}{3} \right] dx \\ &= \int_0^5 \left( x^5 + \frac{x^7}{3} \right) dx = \left[ \frac{x^6}{6} + \frac{x^8}{24} \right]_0^5 = 5^6 \left[ \frac{1}{6} + \frac{5^2}{24} \right] = 18880.2 \text{ nearly.} \end{aligned}$$

**Example 7.2.** Evaluate  $\iint_A xy dx dy$ , where  $A$  is the domain bounded by  $x$ -axis, ordinate  $x = 2a$  and the curve  $x^2 = 4ay$ .

**Solution.** The line  $x = 2a$  and the parabola  $x^2 = 4ay$  intersect at  $L(2a, a)$ . Figure 7.4 shows the domain  $A$  which is the area  $OML$ .

Integrating first over a vertical strip  $PQ$ , i.e., w.r.t.  $y$  from  $P(y = 0)$  to  $Q(y = x^2/4a)$  on the parabola and then w.r.t.  $x$  from  $x = 0$  to  $x = 2a$ , we have

$$\iint_A xy dx dy = \int_0^{2a} dx \int_0^{x^2/4a} xy dy = \int_0^{2a} x \left[ \frac{y^2}{2} \right]_0^{x^2/4a} dx$$




Fig. 7.4

$$= \frac{1}{32a^2} \int_0^{2a} x^5 dx = \frac{1}{32a^2} \left[ \frac{x^6}{6} \right]_0^{2a} = \frac{a^4}{3}.$$

**Otherwise** integrating first over a horizontal strip  $RS$ , i.e., w.r.t.  $x$  from,  $R(x = 2\sqrt{ay})$  on the parabola to  $S(x = 2a)$  and then w.r.t.  $y$  from  $y = 0$  to  $y = a$ , we get

$$\begin{aligned} \iint_A xy \, dx \, dy &= \int_0^a dy \int_{2\sqrt{ay}}^{2a} xy \, dx = \int_0^a y \left[ \frac{x^2}{2} \right]_{2\sqrt{ay}}^{2a} dy \\ &= 2a \int_0^a (ay - y^2) dy = 2a \left[ \frac{ay^2}{2} - \frac{y^3}{3} \right]_0^a = \frac{a^4}{3}. \end{aligned}$$

**Example 7.3.** Evaluate  $\iint_R x^2 \, dx \, dy$  where  $R$  is the region in the first quadrant bounded by the lines  $x = y$ ,  $y = 0$ ,  $x = 8$  and the curve  $xy = 16$ .

**Solution.** The line  $AL$  ( $x = 8$ ) intersects the hyperbola  $xy = 16$  at  $A(8, 2)$  while the line  $y = x$  intersects this hyperbola at  $B(4, 4)$ . Figure 7.5 shows the region  $R$  of integration which is the area  $OLAB$ . To evaluate the given integral, we divide this area into two parts  $OMB$  and  $MLAB$ .




Fig. 7.5

$$\begin{aligned} \therefore \iint_R x^2 \, dx \, dy &= \int_{x \text{ at } 0}^{x \text{ at } M} \int_{y \text{ at } P}^{y \text{ at } Q} x^2 \, dx \, dy + \int_{x \text{ at } M}^{x \text{ at } L} \int_{y \text{ at } P'}^{y \text{ at } Q'} x^2 \, dx \, dy \\ &= \int_0^4 \int_0^x x^2 \, dx \, dy + \int_4^8 \int_0^{16/x} x^2 \, dx \, dy \\ &= \int_0^4 x^2 \, dx \left[ y \right]_0^x + \int_4^8 x^2 \, dx \left[ y \right]_0^{16/x} \\ &= \int_0^4 x^3 \, dx + \int_4^8 16x \, dx = \left[ \frac{x^4}{4} \right]_0^4 + 16 \left[ \frac{x^2}{2} \right]_4^8 = 448 \end{aligned}$$

## 7.2 CHANGE OF ORDER OF INTEGRATION

In a double integral with variable limits, the change of order of integration changes the limit of integration. While doing so, sometimes it is required to split up the region of integration and the given integral is expressed as the sum of a number of double integrals with changed limits. To fix up the new limits, it is always advisable to draw a rough sketch of the region of integration.

The change of order of integration quite often facilitates the evaluation of a double integral. The following examples will make these ideas clear.

**Example 7.4.** By changing the order of integration of  $\int_0^\infty \int_0^\infty e^{-xy} \sin px \, dx \, dy$ , show that

$$\int_0^\infty \frac{\sin px}{x} \, dx = \frac{\pi}{2}.$$

(U.P.T.U., 2004)

**Solution.** 
$$\int_0^\infty \int_0^\infty e^{-xy} \sin px \, dx \, dy = \int_0^\infty \left( \int_0^\infty e^{-xy} \sin px \, dx \right) dy$$



$$\begin{aligned}
 &= \int_0^\infty \left| -\frac{e^{-xy}}{p^2 + y^2} (p \cos px + y \sin px) \right|_0^\infty dy \\
 &= \int_0^\infty \frac{p}{p^2 + y^2} dy = \left| \tan^{-1} \left( \frac{y}{p} \right) \right|_0^\infty = \frac{\pi}{2} \quad \dots(i)
 \end{aligned}$$

On changing the order of integration, we have

$$\begin{aligned}
 \int_0^\infty \int_0^\infty e^{-xy} \sin px \, dx \, dy &= \int_0^\infty \sin px \left\{ \int_0^\infty e^{-xy} \, dy \right\} dx \\
 &= \int_0^\infty \sin px \left| \frac{e^{-xy}}{-x} \right|_0^\infty dx = \int_0^\infty \frac{\sin px}{x} dx \quad \dots(ii)
 \end{aligned}$$

Thus from (i) and (ii), we have  $\int_0^\infty \frac{\sin px}{x} dx = \frac{\pi}{2}$ .

**Example 7.5.** Change the order of integration in the integral

$$I = \int_{-a}^a \int_0^{\sqrt{a^2 - y^2}} f(x, y) \, dx \, dy.$$

**Solution.** Here the elementary strip is parallel to  $x$ -axis (such as  $PQ$ ) and extends from  $x = 0$  to  $x = \sqrt{a^2 - y^2}$  (i.e., to the circle  $x^2 + y^2 = a^2$ ) and this strip slides from  $y = -a$  to  $y = a$ . This shaded semi-circular area is, therefore, the region of integration (Fig. 7.6).

On changing the order of integration, we first integrate w.r.t.  $y$  along a vertical strip  $RS$  which extends from  $R [y = -\sqrt{a^2 - x^2}]$  to  $S [y = \sqrt{a^2 - x^2}]$ . To cover the given region, we then integrate w.r.t.  $x$  from  $x = 0$  to  $x = a$ .




Fig. 7.6

$$\begin{aligned}
 \text{Thus} \quad I &= \int_0^a dx \int_{-\sqrt{a^2 - x^2}}^{\sqrt{a^2 - x^2}} f(x, y) \, dy \\
 \text{or} \quad &= \int_0^a \int_{-\sqrt{a^2 - x^2}}^{\sqrt{a^2 - x^2}} f(x, y) \, dy \, dx.
 \end{aligned}$$

**Example 7.6.** Evaluate  $\int_0^1 \int_{e^x}^e dy \, dx / \log y$  by changing the order of integration.

**Solution.** Here the integration is first w.r.t.  $y$  from  $P$  on  $y = e^x$  to  $Q$  on the line  $y = e$ . Then the integration is w.r.t.  $x$  from  $x = 0$  to  $x = 1$ , giving the shaded region  $ABC$  (Fig. 7.7).

On changing the order of integration, we first integrate w.r.t.  $x$  from  $R$  on  $x = 0$  to  $S$  on  $x = \log y$  and then w.r.t.  $y$  from  $y = 1$  to  $y = e$ .

$$\begin{aligned}
 \text{Thus} \quad \int_0^1 \int_{e^x}^e \frac{dy \, dx}{\log y} &= \int_1^e \int_0^{\log y} \frac{dx \, dy}{\log y} \\
 &= \int_1^e \frac{dy}{\log y} \left| x \right|_0^{\log y} = \int_1^e dy = \left| y \right|_1^e = e - 1.
 \end{aligned}$$




Fig. 7.7

**Example 7.7.** Change the order of integration in  $I = \int_0^{4a} \int_{x^2/4a}^{2\sqrt{ax}} dy \, dx$  and hence evaluate.

(Nagpur, 2009 ; P.T.U., 2009 S)

**Solution.** Here integration is first w.r.t.  $y$  and  $P$  on the parabola  $x^2 = 4ay$  to  $Q$  on the parabola  $y^2 = 4ax$  and then w.r.t.  $x$  from  $x = 0$  to  $x = 4a$  giving the shaded region of integration (Fig. 7.8).

On changing the order of integration, we first integrate w.r.t.  $x$  from  $R$  to  $S$ , then w.r.t.  $y$  from  $y = 0$  to  $y = 4a$

$$\begin{aligned} \therefore I &= \int_0^{4a} \int_{y^2/4a}^{2\sqrt{ay}} dx dy = \int_0^{4a} dy \left[ x \right]_{y^2/4a}^{2\sqrt{ay}} = \int_0^{4a} (2\sqrt{ay} - y^2/4a) dy \\ &= \left[ 2\sqrt{a} \frac{y^{3/2}}{3/2} - \frac{y^3}{12a} \right]_0^{4a} = \frac{32a^2}{3} - \frac{16a^2}{3} = \frac{16a^2}{3}. \end{aligned}$$




Fig. 7.8

**Example 7.8.** Change the order of integration and hence evaluate

$$I = \int_0^a \int_{\sqrt{ax}}^a \frac{y^2 dx dy}{\sqrt{(y^4 - a^2 x^2)}}$$

(S.V.T.U., 2006 S)

**Solution.** Here integration is first w.r.t.  $y$  from  $P$  on the parabola  $y^2 = ax$  to  $Q$  on the line  $y = a$ , then w.r.t.  $x$  from  $x = 0$  to  $x = a$ , giving the shaded region  $OAB$  of integration (Fig. 7.9).

On changing the order of integration, we first integrate w.r.t.  $x$  from  $R$  to  $S$ , then w.r.t.  $y$  from  $y = 0$  to  $y = a$ .

$$\begin{aligned} \therefore I &= \int_0^a \int_0^{y^2/a} \frac{y^2 dy}{\sqrt{(y^4 - a^2 x^2)}} dx = \frac{1}{a} \int_0^a \int_0^{y^2/a} y^2 dy \frac{dx}{\sqrt{[(y^2/a)^2 - x^2]}} dx \\ &= \frac{1}{a} \int_0^a y^2 dy \left[ \sin^{-1} \left( \frac{xa}{y^2} \right) \right]_0^{y^2/a} = \frac{1}{a} \int_0^a y^2 dy [\sin^{-1}(1) - \sin^{-1}(0)] \\ &= \frac{\pi}{2a} \int_0^a y^2 dy = \frac{\pi}{2a} \left[ \frac{y^3}{3} \right]_0^a = \frac{\pi a^2}{6}. \end{aligned}$$




Fig. 7.9

**Example 7.9.** Change the order of integration in  $I = \int_0^1 \int_{x^2}^{2-x} xy dx dy$  and hence evaluate the same.

(Bhopal, 2008 ; V.T.U., 2008 ; S.V.T.U., 2007 ; P.T.U., 2005 ; U.P.T.U., 2005)

**Solution.** Here the integration is first w.r.t.  $y$  along a vertical strip  $PQ$  which extends from  $P$  on the parabola  $y = x^2$  to  $Q$  on the line  $y = 2 - x$ . Such a strip slides from  $x = 0$  to  $x = 1$ , giving the region of integration as the curvilinear triangle  $OAB$  (shaded) in Fig. 7.10.

On changing the order of integration, we first integrate w.r.t.  $x$  along a horizontal strip  $P'Q'$  and that requires the splitting up of the region  $OAB$  into two parts by the line  $AC$  ( $y = 1$ ), i.e., the curvilinear triangle  $OAC$  and the triangle  $ABC$ .

For the region  $OAC$ , the limits of integration for  $x$  are from  $x = 0$  to  $x = \sqrt{y}$  and those for  $y$  are from  $y = 0$  to  $y = 1$ . So the contribution to  $I$  from the region  $OAC$  is

$$I_1 = \int_0^1 dy \int_0^{\sqrt{y}} xy dx$$

For the region  $ABC$ , the limits of integration for  $x$  are from  $x = 0$  to  $x = 2 - y$  and those for  $y$  are from  $y = 1$  to  $y = 2$ . So the contribution to  $I$  from the region  $ABC$  is

$$I_2 = \int_1^2 dy \int_0^{2-y} xy dx.$$




Fig. 7.10



Hence, on reversing the order of integration,

$$\begin{aligned} I &= \int_0^1 dy \int_0^{\sqrt{y}} xy \, dx + \int_1^2 dy \int_0^{2-y} xy \, dx \\ &= \int_0^1 dy \left[ \frac{x^2}{2} \cdot y \right]_0^{\sqrt{y}} + \int_1^2 dy \left[ \frac{x^2}{2} \cdot y \right]_0^{2-y} = \frac{1}{2} \int_0^1 y^2 dy + \frac{1}{2} \int_1^2 y(2-y)^2 dy = \frac{1}{6} + \frac{5}{24} = \frac{3}{8}. \end{aligned}$$

**Example 7.10.** Change the order of integration in  $I = \int_0^1 \int_x^{\sqrt{2-x^2}} \frac{x}{\sqrt{x^2+y^2}} dx$  and hence evaluate it. (J.N.T.U., 2005 ; Rohtak, 2003)

**Solution.** Here the integration is first w.r.t.  $y$  along  $PQ$  which extends from  $P$  on the line  $y = x$  to  $Q$  on the circle  $y = \sqrt{2-x^2}$ . Then  $PQ$  slides from  $y = 0$  to  $y = 1$ , giving the region of integration  $OAB$  as in Fig. 7.11.

On changing the order of integration, we first integrate w.r.t.  $x$  from  $P'$  to  $Q'$  and that requires splitting the region  $OAB$  into two parts  $OAC$  and  $ABC$ .

For the region  $OAC$ , the limits of integration for  $x$  are from  $x = 0$  to  $x = 1$  and those for  $y$  are from  $y = 0$  to  $y = 1$ . So the contribution to  $I$  from the region  $OAC$  is

$$I_1 = \int_0^1 dy \int_0^y \frac{x}{\sqrt{x^2+y^2}} dx.$$

For the region  $ABC$ , the limits of integration for  $x$  are 0 to  $\sqrt{2-y^2}$  and these for  $y$  are from 1 to  $\sqrt{2}$ . So the contribution to  $I$  from the region  $ABC$  is

$$I_2 = \int_1^{\sqrt{2}} dy \int_0^{\sqrt{2-y^2}} \frac{x}{\sqrt{x^2+y^2}} dx$$

Hence

$$\begin{aligned} I &= \int_0^1 \left[ (x^2+y^2)^{1/2} \right]_0^y dy + \int_1^{\sqrt{2}} \left[ (x^2+y^2)^{1/2} \right]_0^{\sqrt{2-y^2}} dy \\ &= \int_0^1 (\sqrt{2}-1)y \, dy + \int_1^{\sqrt{2}} \sqrt{2-y} \, dy = \frac{1}{2}(\sqrt{2}-1) + \sqrt{2}\sqrt{2-1} - \frac{1}{2} = 1 - 1/\sqrt{2}. \end{aligned}$$




Fig. 7.11

### 7.3 DOUBLE INTEGRALS IN POLAR COORDINATES

To evaluate  $\int_{\theta_1}^{\theta_2} \int_{r_1}^{r_2} f(r, \theta) dr d\theta$ , we first integrate w.r.t.  $r$  between limits  $r = r_1$  and  $r = r_2$  keeping  $\theta$  fixed and the resulting expression is integrated w.r.t.  $\theta$  from  $\theta_1$  to  $\theta_2$ . In this integral,  $r_1, r_2$  are functions of  $\theta$  and  $\theta_1, \theta_2$  are constants. Figure 7.12 illustrates the process geometrically.

Here  $AB$  and  $CD$  are the curves  $r_1 = f_1(\theta)$  and  $r_2 = f_2(\theta)$  bounded by the lines  $\theta = \theta_1$  and  $\theta = \theta_2$ .  $PQ$  is a wedge of angular thickness  $\delta\theta$ .

Then  $\int_{r_1}^{r_2} f(r, \theta) dr$  indicates that the integration is along  $PQ$  from  $P$  to  $Q$  while the integration w.r.t.  $\theta$  corresponds to the turning of  $PQ$  from  $AC$  to  $BD$ .

Thus the whole region of integration is the area  $ACDB$ . The order of integration may be changed with appropriate changes in the limits.




Fig. 7.12

**Example 7.11.** Evaluate  $\iint_R r \sin \theta \, dr \, d\theta$  over the cardioid  $r = a(1 - \cos \theta)$  above the initial line.

(Kerala, 2005)

**Solution.** To integrate first w.r.t.  $r$ , the limits are from 0 ( $r = 0$ ) to  $P$  [ $r = a(1 - \cos \theta)$ ] and to cover the region of integration  $R$ ,  $\theta$  varies from 0 to  $\pi$  (Fig. 7.13).

$$\begin{aligned} \therefore \iint_R r \sin \theta \, dr \, d\theta &= \int_0^\pi \sin \theta \left[ \int_0^{a(1-\cos \theta)} r \, dr \right] d\theta \\ &= \int_0^\pi \sin \theta \, d\theta \left[ \frac{r^2}{2} \right]_0^{a(1-\cos \theta)} = \frac{a^2}{2} \int_0^\pi (1 - \cos \theta)^2 \cdot \sin \theta \, d\theta \\ &= \frac{a^2}{2} \left[ \frac{(1 - \cos \theta)^3}{3} \right]_0^\pi = \frac{a^2}{2} \cdot \frac{8}{3} = \frac{4a^2}{3}. \end{aligned}$$




Fig. 7.13

**Example 7.12.** Calculate  $\iint r^3 \, dr \, d\theta$  over the area included between the circles  $r = 2 \sin \theta$  and  $r = 4 \sin \theta$ .

**Solution.** Given circles  $r = 2 \sin \theta$  ... (i)  
and  $r = 4 \sin \theta$  ... (ii)

are shown in Fig. 7.14. The shaded area between these circles is the region of integration.

If we integrate first w.r.t.  $r$ , then its limits are from  $P(r = 2 \sin \theta)$  to  $Q(r = 4 \sin \theta)$  and to cover the whole region  $\theta$  varies from 0 to  $\pi$ . Thus the required integral is

$$\begin{aligned} I &= \int_0^\pi d\theta \int_{2 \sin \theta}^{4 \sin \theta} r^3 \, dr = \int_0^\pi d\theta \left[ \frac{r^4}{4} \right]_{2 \sin \theta}^{4 \sin \theta} \\ &= 60 \int_0^\pi \sin^4 \theta \, d\theta = 60 \times 2 \int_0^{\pi/2} \sin^4 \theta \, d\theta = 120 \times \frac{3}{4} \cdot \frac{1}{2} \cdot \frac{\pi}{2} = 22.5 \pi. \end{aligned}$$




Fig. 7.14

### PROBLEMS 7.1

Evaluate the following integrals (1–7) :

- $\int_1^2 \int_1^3 xy^2 \, dx \, dy$ .
- $\int_0^1 \int_x^{\sqrt{x}} (x^2 + y^2) \, dx \, dy$ . (V.T.U., 2000)
- $\int_0^1 \int_0^x e^{x/y} \, dx \, dy$ . (P.T.U., 2005)
- $\int_0^1 \int_0^{\sqrt{1+x^2}} \frac{dy \, dx}{1+x^2+y^2}$ . (Rajasthan, 2005)
- $\iint xy \, dx \, dy$  over the positive quadrant of the circle  $x^2 + y^2 = a^2$ . (Rajasthan, 2006)
- $\iint (x+y)^2 \, dx \, dy$  over the area bounded by the ellipse  $x^2/a^2 + y^2/b^2 = 1$ . (Kurukshetra, 2009 S ; U.P.T.U., 2004 S)
- $\iint xy(x+y) \, dx \, dy$  over the area between  $y = x^2$  and  $y = x$ . (V.T.U., 2010)

Evaluate the following integrals by changing the order of integration (8–15) :

- $\int_0^a \int_y^a \frac{x \, dx \, dy}{x^2 + y^2}$ . (Bhopal, 2008)
- $\int_0^3 \int_1^{\sqrt{4-y}} (x+y) \, dx \, dy$ . (V.T.U., 2005 ; Anna, 2003 S ; Delhi, 2002)



10.  $\int_0^1 \int_x^{\sqrt{2-x^2}} \frac{x \, dy \, dx}{\sqrt{(x^2 + y^2)}} \quad (P.T.U., 2010 ; Marathwada, 2008 ; U.P.T.U., 2006)$
11.  $\int_0^{a/\sqrt{2}} \int_y^{\sqrt{a^2 - y^2}} \log(x^2 + y^2) \, dx \, dy \quad (a > 0).$
12.  $\int_0^1 \int_x^{\sqrt{x}} xy \, dy \, dx. \quad (V.T.U., 2010)$
13.  $\int_0^a \int_{a-\sqrt{a^2 - y^2}}^{a+\sqrt{a^2 - y^2}} xy \, dx \, dy. \quad (Anna, 2009)$
14.  $\int_0^\infty \int_x^\infty \frac{e^{-y}}{y} \, dy \, dx. \quad (Bhopal, 2009 ; S.V.T.U., 2009 ; V.T.U., 2007)$
15.  $\int_0^\infty \int_0^x xe^{-x^2/y} \, dy \, dx. \quad (S.V.T.U., 2006 ; U.P.T.U., 2005 ; V.T.U., 2004)$
16. Sketch the region of integration of the following integrals and change the order of integrations,  
 (i)  $\int_0^{2a} \int_{\sqrt{2ax-x^2}}^{\sqrt{2ax}} f(x) \, dx \, dy$  (Rajasthan, 2006) (ii)  $\int_0^{ae^{\pi/4}} \int_{2 \log(r/a)}^{\pi/2} f(r, \theta) r \, dr \, d\theta.$
17. Show that  $\iint_R r^2 \sin \theta \, dr \, d\theta = 2a^2/3$ , where  $R$  is the semi-circle  $r = 2a \cos \theta$  above the initial line.
18. Evaluate  $\iint \frac{r \, dr \, d\theta}{\sqrt{a^2 + r^2}}$  over one loop of the lemniscate  $r^2 = a^2 \cos 2\theta$ . (Rohtak, 2006 S ; P.T.U., 2005)
19. Evaluate  $\iint r^3 \, dr \, d\theta$  over the area bounded between the circles  $r = 2 \cos \theta$  and  $r = 4 \cos \theta$ .  
 (Anna, 2009 ; Madras, 2006)

## 7.4 AREA ENCLOSED BY PLANE CURVES

### (1) Cartesian coordinates.

Consider the area enclosed by the curves  $y = f_1(x)$  and  $y = f_2(x)$  and the ordinates  $x = x_1, x = x_2$  [Fig. 7.15 (a)].

Divide this area into vertical strips of width  $\delta x$ . If  $P(x, y), Q(x + \delta x, y + \delta y)$  be two neighbouring points, then the area of the small rectangle  $PQ = \delta x \delta y$ .

$$\therefore \text{area of strip } KL = \lim_{\delta y \rightarrow 0} \sum \delta x \delta y.$$

Since for all rectangles in this strip  $\delta x$  is the same and  $y$  varies from  $y = f_1(x)$  to  $y = f_2(x)$ .

$$\therefore \text{area of the strip } KL = \delta x \lim_{\delta y \rightarrow 0} \sum_{f_1(x)}^{f_2(x)} dy = \delta x \int_{f_1(x)}^{f_2(x)} dy.$$

Now adding up all such strips from  $x = x_1$  to  $x = x_2$ , we get the area  $ABCD$

$$= \lim_{\delta x \rightarrow 0} \sum_{x_1}^{x_2} \delta x \cdot \int_{f_1(x)}^{f_2(x)} dy = \int_{x_1}^{x_2} dx \int_{f_1(x)}^{f_2(x)} dy = \int_{x_1}^{x_2} \int_{f_1(x)}^{f_2(x)} dx \, dy$$

Similarly, dividing the area  $A'B'C'D'$  [Fig. 7.15(b)] into horizontal strips of width  $\delta y$ , we get the area  $A'B'C'D'$ .

$$= \int_{y_1}^{y_2} \int_{f_1(y)}^{f_2(y)} dx \, dy$$

### (2) Polar coordinates.

Consider an area  $A$  enclosed by a curve whose equation is in polar coordinates.

Let  $P(r, \theta), Q(r + \delta r, \theta + \delta \theta)$  be two neighbouring points. Mark circular areas of radii  $r$  and  $r + \delta r$  meeting  $OQ$  in  $R$  and  $OP$  (produced) in  $S$  (Fig. 7.16).




Fig. 7.15(a)




Fig. 7.15 (b)

Since arc  $PR = r\delta\theta$  and  $PS = \delta r$ .

$\therefore$  area of the curvilinear rectangle  $PRQS$  is approximately  
 $= PR \cdot PS = r\delta\theta \cdot \delta r$ .

If the whole area is divided into such curvilinear rectangles, the sum  $\sum r\delta\theta\delta r$  taken for all these rectangles, gives in the limit the area  $A$ .

$$\text{Hence } A = \lim_{\substack{\delta r \rightarrow 0 \\ \delta\theta \rightarrow 0}} \sum r\delta\theta\delta r = \iint r dr d\theta$$

where the limits are to be so chosen as to cover the entire area.




Fig. 7.16

**Example 7.13.** Find the area of a plate in the form of a quadrant of the ellipse

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1.$$

(V.T.U., 2001 ; Osmania, 2000 S)

**Solution.** Dividing the area into vertical strips of width

$\delta x$ ,  $y$  varies from  $K(y = 0)$  to  $L[y = b\sqrt{1 - x^2/b^2}]$  and then  $x$  varies from 0 to  $a$  (Fig. 7.17).

$\therefore$  required area

$$\begin{aligned} &= \int_0^a dx \int_0^{b\sqrt{1-x^2/a^2}} dy = \int_0^a dx [y]_0^{b\sqrt{1-x^2/a^2}} \\ &= \frac{b}{a} \int_0^a \sqrt{a^2 - x^2} dx = \pi ab/4. \end{aligned}$$




Fig. 7.17

**Otherwise,** dividing this area into horizontal strips of width  $\delta y$ ,  $x$  varies from  $M(x = 0)$  to

$N[x = a\sqrt{1 - y^2/b^2}]$  and then  $y$  varies from 0 to  $b$ .

$$\begin{aligned} \therefore \text{required area} &= \int_0^b dy \int_0^{a\sqrt{1-y^2/b^2}} dx = \int_0^b dy [x]_0^{a\sqrt{1-y^2/b^2}} \\ &= \frac{a}{b} \int_0^b \sqrt{b^2 - y^2} dy = \pi ab/4. \end{aligned}$$

**Obs.** The change of the order of integration does not in any way affect the value of the area.

**Example 7.14.** Show that the area between the parabolas  $y^2 = 4ax$  and  $x^2 = 4ay$  is  $\frac{16}{3} a^2$ .

(Kerala, 2005 ; Rohtak, 2003)

**Solution.** Solving the equations  $y^2 = 4ax$  and  $x^2 = 4ay$ , it is seen that the parabolas intersect at  $O(0, 0)$  and  $A(4a, 4a)$ . As such for the shaded area between these parabolas (Fig. 7.18)  $x$  varies from 0 to  $4a$  and  $y$  varies from  $P$  to  $Q$  i.e., from  $y = x^2/4a$  to  $y = 2\sqrt{ax}$ . Hence the required area

$$\begin{aligned} &= \int_0^{4a} \int_{x^2/4a}^{2\sqrt{ax}} dy dx = \int_0^{4a} (2\sqrt{ax} - x^2/4a) dx \\ &= \left[ 2\sqrt{a} \cdot \frac{2}{3} x^{3/2} - \frac{1}{4a} \cdot \frac{x^3}{3} \right]_0^{4a} = \frac{32}{3} a^2 - \frac{16}{3} a^2 = \frac{16}{3} a^2. \end{aligned}$$




Fig. 7.18



**Example 7.15.** Calculate the area included between the curve  $r = a(\sec \theta + \cos \theta)$  and its asymptote.

**Solution.** The curve is symmetrical about the initial line and has an asymptote  $r = a \sec \theta$  (Fig. 7.19).

Draw any line  $OP$  cutting the curve at  $P$  and its asymptote at  $P'$ . Along this line,  $\theta$  is constant and  $r$  varies from  $a \sec \theta$  at  $P'$  to  $a(\sec \theta + \cos \theta)$  at  $P$ . Then to get the upper half of the area,  $\theta$  varies from 0 to  $\pi/2$ .

$$\begin{aligned} \therefore \text{required area} &= 2 \int_0^{\pi/2} \int_{a \sec \theta}^{a(\sec \theta + \cos \theta)} r \, dr \, d\theta \\ &= 2 \int_0^{\pi/2} \left[ \frac{r^2}{2} \right]_{a \sec \theta}^{a(\sec \theta + \cos \theta)} d\theta \\ &= a^2 \int_0^{\pi/2} (2 + \cos^2 \theta) d\theta = 5\pi a^2/4. \end{aligned}$$




Fig. 7.19

**Example 7.16.** Find the area lying inside the cardioid  $r = a(1 + \cos \theta)$  and outside the circle  $r = a$ .

**Solution.** In Fig. 7.20,  $ABODA$  represents the cardioid  $r = a(1 + \cos \theta)$  and  $CBA'DC$  is the circle  $r = a$ .

Required area (shaded) = 2 (area  $ABCA$ )

$$\begin{aligned} &= 2 \int_0^{\pi/2} \int_{r=OP'}^{r=OP} r \, d\theta \, dr = 2 \int_0^{\pi/2} \int_a^{a(1+\cos \theta)} (r \, dr) d\theta \\ &= 2 \int_0^{\pi/2} \left[ \frac{r^2}{2} \right]_a^{a(1+\cos \theta)} d\theta = a^2 \int_0^{\pi/2} [(1 + \cos \theta)^2 - 1] d\theta \\ &= a^2 \int_0^{\pi/2} (\cos^2 \theta + 2 \cos \theta) d\theta = a^2 \left( \frac{1}{2} \cdot \frac{\pi}{2} + 2 \right) = \frac{a^2}{4} (\pi + 8). \end{aligned}$$




Fig. 7.20

## PROBLEMS 7.2

- Find, by double integration, the area lying between the parabola  $y = 4x - x^2$  and the line  $y = x$ . (Anna, 2009)
- Find the area lying between the parabola  $y = x^2$  and the line  $x + y - z = 0$ . (U.P.T.U., 2001)
- By double integration, find the whole area of the curve  $a^2x^2 = y^3(2a - y)$ . (J.N.T.U., 2005)
- Find, by double integration, the area enclosed by the curves  $y = 3x/(x^2 + 2)$  and  $4y = x^2$ . (Madras, 2000 S)
- Find, by double integration, the area of the lemniscate  $r^2 = a^2 \cos 2\theta$ . (Anna 2009 ; Mumbai, 2006)
- Find, by double integration, the area lying inside the circle  $r = a \sin \theta$  and outside the cardioid  $r = a(1 - \cos \theta)$ . (Mumbai, 2007)
- Find the area lying inside the cardioid  $r = 1 + \cos \theta$  and outside the parabola  $r(1 + \cos \theta) = 1$ .
- Find the area common to the circles  $r = a \cos \theta$ ,  $r = a \sin \theta$  by double integration.

## 7.5 TRIPLE INTEGRALS

Consider a function  $f(x, y, z)$  defined at every point of the 3-dimensional finite region  $V$ . Divide  $V$  into  $n$  elementary volumes  $\delta V_1, \delta V_2, \dots, \delta V_n$ . Let  $(x_r, y_r, z_r)$  be any point within the  $r$ th sub-division  $\delta V_r$ . Consider the sum

$$\sum_{r=1}^n f(x_r, y_r, z_r) \delta V_r.$$

The limit of this sum, if it exists, as  $n \rightarrow \infty$  and  $\delta V_r \rightarrow 0$  is called the *triple integral* of  $f(x, y, z)$  over the region  $V$  and is denoted by

$$\iiint_V f(x, y, z) dV.$$

For purposes of evaluation, it can also be expressed as the repeated integral

$$\int_{x_1}^{x_2} \int_{y_1}^{y_2} \int_{z_1}^{z_2} f(x, y, z) dx dy dz.$$

If  $x_1, x_2$  are constants ;  $y_1, y_2$  are either constants or functions of  $x$  and  $z_1, z_2$  are either constants or functions of  $x$  and  $y$ , then this integral is evaluated as follows :

First  $f(x, y, z)$  is integrated w.r.t.  $z$  between the limits  $z_1$  and  $z_2$  keeping  $x$  and  $y$  fixed. The resulting expression is integrated w.r.t.  $y$  between the limits  $y_1$  and  $y_2$  keeping  $x$  constant. The result just obtained is finally integrated w.r.t.  $x$  from  $x_1$  to  $x_2$ .

$$\text{Thus} \quad I = \int_{x_1}^{x_2} \int_{y_1(x)}^{y_2(x)} \int_{z_1(x,y)}^{z_2(x,y)} f(x, y, z) dz dy dx$$

where the integration is carried out from the innermost rectangle to the outermost rectangle.

The order of integration may be different for different types of limits.

**Example 7.17.** Evaluate  $\int_{-1}^1 \int_0^z \int_{x-z}^{x+z} (x + y + z) dx dy dz$ . (J.N.T.U., 2006 ; Cochin, 2005)

**Solution.** Integrating first w.r.t.  $y$  keeping  $x$  and  $z$  constant, we have

$$\begin{aligned} I &= \int_{-1}^1 \int_0^z \left[ xy + \frac{y^2}{2} + yz \right]_{x-z}^{x+z} dx dz = \int_{-1}^1 \int_0^z \left[ (x+z)(2z) + \frac{1}{2}4xz \right] dx dz \\ &= 2 \int_{-1}^1 \left[ \frac{x^2 z}{2} + z^2 x + \frac{x^2}{2} z \right]_0^z dz = 2 \int_{-1}^1 \left( \frac{z^3}{2} + z^3 + \frac{z^3}{2} \right) dz = 4 \left[ \frac{z^4}{4} \right]_{-1}^1 = 0. \end{aligned}$$

**Example 7.18.** Evaluate  $\int_0^1 \int_0^{\sqrt{1-x^2}} \int_0^{\sqrt{1-x^2-y^2}} xyz dx dy dz$ . (V.T.U., 2003 S)

**Solution.** We have

$$\begin{aligned} I &= \int_0^1 x \left[ \int_0^{\sqrt{1-x^2}} y \left\{ \int_0^{\sqrt{1-x^2-y^2}} z dz \right\} dy \right] dx = \int_0^1 x \left\{ \int_0^{\sqrt{1-x^2}} y \cdot \left[ \frac{z^2}{2} \right]_0^{\sqrt{1-x^2-y^2}} dy \right\} dx \\ &= \int_0^1 x \left\{ \int_0^{\sqrt{1-x^2}} y \cdot \frac{1}{2}(1-x^2-y^2) dy \right\} dx = \frac{1}{2} \int_0^1 x \left[ (1-x^2) \frac{y^2}{2} - \frac{y^4}{4} \right]_0^{\sqrt{1-x^2}} dx \\ &= \frac{1}{8} \int_0^1 [(1-x^2)^2 \cdot 2x - (1-x^2)^4 \cdot x] dx = \frac{1}{8} \int_0^1 (x - 2x^3 + x^5) dx \\ &= \frac{1}{8} \left[ \frac{x^2}{2} - \frac{2x^4}{4} + \frac{x^6}{6} \right]_0^1 = \frac{1}{8} \left( \frac{1}{2} - \frac{1}{2} + \frac{1}{6} \right) = \frac{1}{48}. \end{aligned}$$

### PROBLEMS 7.3

Evaluate the following integrals :

- $\int_0^a \int_0^b \int_0^c (x^2 + y^2 + z^2) dx dy dz$ . (Anna, 2009)
- $\int_c^e \int_b^d \int_a^a (x^2 + y^2 + z^2) dx dy dz$  (S.V.T.U., 2009 ; V.T.U. 2000)
- $\int_0^1 \int_{y^2}^1 \int_0^{1-x} x dz dx dy$  (Nagpur, 2009)
- $\int_0^a \int_0^x \int_0^{x+y} e^{x+y+z} dz dy dx$ . (V.T.U., 2010 ; Kurukshetra, 2009 S ; J.N.T.U., 2005)
- $\int_0^{\log 2} \int_0^x \int_0^{x+\log y} e^{x+y+z} dx dy dz$ . (Bhopal, 2008)
- $\int_1^e \int_1^{\log y} \int_1^{e^x} \log z dz dx dy$ . (S.V.T.U., 2008 ; Rohtak, 2005)
- $\int_0^{\pi/2} \int_0^{\sin \theta} \int_0^{\frac{a^2-r^2}{a}} rdz dr d\theta$ . (V.T.U., 2009)



## 7.6 VOLUMES OF SOLIDS

**(1) Volumes as double integrals.** Consider a surface  $z = f(x, y)$ . Let the orthogonal projection on  $XY$ -plane of its portion  $S'$  be the area  $S$  (Fig. 7.21).

Divide  $S$  into elementary rectangles of area  $\delta x \delta y$  by drawing lines parallel to  $X$  and  $Y$ -axes. With each of these rectangles as base, erect a prism having its length parallel to  $OZ$ .

$\therefore$  volume of this prism between  $S$  and the given surface  $z = f(x, y)$  is  $z \delta x \delta y$ .

Hence the volume of the solid cylinder on  $S$  as base, bounded by the given surface with generators parallel to the  $Z$ -axis.

$$\begin{aligned} &= \lim_{\substack{\delta x \rightarrow 0 \\ \delta y \rightarrow 0}} \sum \sum z \delta x \delta y \\ &= \iint z \, dx \, dy \quad \text{or} \quad \iint f(x, y) \, dx \, dy \end{aligned}$$

where the integration is carried over the area  $S$ .

**Obs.** While using polar coordinates, divide  $S$  into elements of area  $r \delta \theta \delta r$ .

$\therefore$  replacing  $dx dy$  by  $r \delta \theta \delta r$ , we get the required volume =  $\iint z r \, d\theta \, dr$ .

**Example 7.19.** Find the volume bounded by the cylinder  $x^2 + y^2 = 4$  and the planes  $y + z = 4$  and  $z = 0$ .  
(S.V.T.U., 2007 ; Cochin, 2005 ; Madras, 2000 S)

**Solution.** From Fig. 7.22, it is self-evident that  $z = 4 - y$  is to be integrated over the circle  $x^2 + y^2 = 4$  in the  $XY$ -plane. To cover the shaded half of this circle,  $x$  varies from 0 to  $\sqrt{4 - y^2}$  and  $y$  varies from  $-2$  to  $2$ .

$\therefore$  Required volume

$$\begin{aligned} &= 2 \int_{-2}^2 \int_0^{\sqrt{4-y^2}} z \, dx \, dy = 2 \int_{-2}^2 \int_0^{\sqrt{4-y^2}} (4 - y) \, dx \, dy \\ &= 2 \int_{-2}^2 (4 - y) [x]_0^{\sqrt{4-y^2}} \, dy = 2 \int_{-2}^2 (4 - y) \sqrt{4 - y^2} \, dy \\ &= 2 \int_{-2}^2 4\sqrt{4 - y^2} \, dy - 2 \int_{-2}^2 y\sqrt{4 - y^2} \, dy \\ &= 8 \int_{-2}^2 \sqrt{4 - y^2} \, dy \quad \text{[The second term vanishes as the integrand is an odd function.]} \\ &= 8 \left[ \frac{y\sqrt{4 - y^2}}{2} + \frac{4}{2} \sin^{-1} \frac{y}{2} \right]_{-2}^2 = 16\pi. \end{aligned}$$

### (2) Volume as triple integral

Divide the given solid by planes parallel to the coordinate planes into rectangular parallelepipeds of volume  $\delta x \delta y \delta z$  (Fig. 7.23).

$$\begin{aligned} \therefore \text{the total volume} &= \lim_{\substack{\delta x \rightarrow 0 \\ \delta y \rightarrow 0 \\ \delta z \rightarrow 0}} \sum \sum \sum \delta x \delta y \delta z \\ &= \iiint dx \, dy \, dz \end{aligned}$$

with appropriate limits of integration.




Fig. 7.21




Fig. 7.22




Fig. 7.23

**Example 7.20.** Calculate the volume of the solid bounded by the planes  $x = 0$ ,  $y = 0$ ,  $x + y + z = a$  and  $z = 0$ .  
(P.T.U., 2009)

**Solution.** Volume required =  $\int_0^a \int_0^{a-x} \int_0^{a-x-y} dz dy dx$

$$= \int_0^a \int_0^{a-x} (a-x-y) dy dx = \int_0^a \left[ (a-x)y - \frac{y^2}{2} \right]_0^{a-x} dx$$

$$= \int_0^a \left\{ (a-x)^2 - \frac{(a-x)^2}{2} \right\} dx = \frac{1}{2} \int_0^a (a-x)^2 dx = \frac{1}{2} \left[ -\frac{(a-x)^3}{3} \right]_0^a = \frac{a^3}{6}.$$

**Example 7.21.** Find the volume of the ellipsoid  $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$ .

(Anna, 2009 ; P.T.U., 2006 ; Kottayam, 2005)

**Solution.** Let  $OABC$  be the positive octant of the given ellipsoid which is bounded by the planes  $OAB$  ( $z = 0$ ),  $OBC$  ( $x = 0$ ),  $OCA$  ( $y = 0$ ) and the surface  $ABC$ , i.e.,

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1.$$

Divide this region  $R$  into rectangular parallelopipeds of volume  $\delta x \delta y \delta z$ . Consider such an element at  $P(x, y, z)$ . (Fig. 7.24)

$\therefore$  the required volume =  $8 \iiint_R dx dy dz$ .

In this region  $R$ ,

(i)  $z$  varies from 0 to  $MN$  where

$$MN = c \sqrt{1 - x^2/a^2 - y^2/b^2}.$$

(ii)  $y$  varies from 0 to  $EF$ , where  $EF = b \sqrt{1 - x^2/a^2}$  from the equation of the ellipse  $OAB$ , i.e.,

$$x^2/a^2 + y^2/b^2 = 1.$$

(iii)  $x$  varies from 0 to  $OA = a$ .

Hence the volume of the whole ellipsoid

$$\begin{aligned} &= 8 \int_0^a \int_0^{b\sqrt{1-x^2/a^2}} \int_0^{c\sqrt{1-x^2/a^2-y^2/b^2}} dz dy dx = 8 \int_0^a dx \int_0^{b\sqrt{1-x^2/a^2}} dy \left[ z \right]_0^{c\sqrt{1-x^2/a^2-y^2/b^2}} \\ &= 8c \int_0^a dx \int_0^{b\sqrt{1-x^2/a^2}} \sqrt{1-x^2/a^2-y^2/b^2} dy \\ &= \frac{8c}{b} \int_0^a dx \int_0^p \sqrt{\rho^2 - y^2} dy \quad \text{when } \rho = b \sqrt{1-x^2/a^2}. \\ &= \frac{8c}{b} \int_0^a dx \left[ \frac{y\sqrt{\rho^2 - y^2}}{2} + \frac{\rho^2}{2} \sin^{-1} \frac{y}{\rho} \right]_0^p = \frac{8c}{b} \int_0^a \frac{b^2}{2} \left( 1 - \frac{x^2}{a^2} \right) \frac{\pi}{2} dx \\ &= 2\pi bc \int_0^a \left( 1 - \frac{x^2}{a^2} \right) dx = 2\pi bc \left[ x - \frac{x^3}{3a^2} \right]_0^a = \frac{4\pi abc}{3}. \end{aligned}$$

**Otherwise.** See Problem 27 page 292.

### (3) Volumes of solids of revolution

Consider an elementary area  $\delta x \delta y$  at the point  $P(x, y)$  of a plane area  $A$ . (Fig. 7.25)

As this elementary area revolves about  $x$ -axis, we get a ring of volume

$$= \pi[(y + \delta y)^2 - y^2] \delta x = 2\pi y \delta x \delta y,$$

nearly to the first powers of  $\delta y$ .




Fig. 7.24



Hence the total volume of the solid formed by the revolution of the area  $A$  about  $x$ -axis.

$$= \iint_A 2\pi y \, dx dy.$$

In polar coordinates, the above formula for the volume becomes

$$\iint_A 2\pi r \sin \theta \cdot r d\theta dr, \text{ i.e. } \iint_A 2\pi r^2 \sin \theta \, d\theta dr$$

Similarly, the volume of the solid formed by the revolution of the area  $A$  about  $y$ -axis =  $\iint_A 2\pi x \, dx dy$ .




Fig. 7.25

**Example 7.22.** Calculate by double integration, the volume generated by the revolution of the cardioid  $r = a(1 - \cos \theta)$  about its axis.

**Solution.** Required volume

$$\begin{aligned} &= \int_0^\pi \int_0^{a(1-\cos \theta)} 2\pi r^2 \sin \theta \, dr \, d\theta \\ &= 2\pi \int_0^\pi \left[ \frac{r^3}{3} \right]_0^{a(1-\cos \theta)} \sin \theta \, d\theta \\ &= \frac{2\pi a^3}{3} \int_0^\pi (1 - \cos \theta)^3 \cdot \sin \theta \, d\theta = \frac{2\pi a^3}{3} \left[ \frac{(1 - \cos \theta)^4}{4} \right]_0^\pi = \frac{8\pi a^3}{3}. \end{aligned}$$




Fig. 7.26

## 7.7 CHANGE OF VARIABLES

An appropriate choice of co-ordinates quite often facilitates the evaluation of a double or a triple integral. By changing the variables, a given integral can be transformed into a simpler integral involving the new variables.

(1) In a double integral, let the variables  $x, y$  be changed to the new variables  $u, v$  by the transformation.

$$x = \phi(u, v), y = \psi(u, v)$$

where  $\phi(u, v)$  and  $\psi(u, v)$  are continuous and have continuous first order derivatives in some region  $R'_{uv}$  in the  $uv$ -plane which corresponds to the region  $R_{xy}$  in the  $xy$ -plane. Then

$$\iint_{R_{xy}} f(x, y) \, dx dy = \iint_{R'_{uv}} f[\phi(u, v), \psi(u, v)] |J| \, du dv \quad \dots(1)$$

where

$$J = \frac{\partial(x, y)}{\partial(u, v)} (\neq 0)$$

is the Jacobian of transformation\* from  $(x, y)$  to  $(u, v)$  coordinates.

(2) For triple integrals, the formula corresponding to (1) is

$$\iiint_{R_{xyz}} f(x, y, z) \, dx dy dz = \iiint_{R'_{uvw}} f[x(u, v, w), y(u, v, w), z(u, v, w)] |J| \, du dv dw$$

where

$$J = \frac{\partial(x, y, z)}{\partial(u, v, w)} (\neq 0)$$

is the Jacobian of transformation from  $(x, y, z)$  to  $(u, v, w)$  coordinates.

**Particular cases :**

(i) To change cartesian coordinates  $(x, y)$  to polar coordinates  $(r, \theta)$ , we have  $x = r \cos \theta, y = r \sin \theta$  and

$$J = \frac{\partial(x, y)}{\partial(r, \theta)} = r \quad [\text{Ex. 5.25, p. 216}]$$

$$\therefore \iint_{R_{xy}} f(x, y) \, dx dy = \iint_{R'_{r\theta}} f(r \cos \theta, r \sin \theta) \cdot r \, dr \, d\theta.$$

\* See footnote page 215.

(ii) To change rectangular coordinates  $(x, y, z)$  to cylindrical coordinates  $(\rho, \phi, z)$  — Fig. 8.27, we have

$$x = \rho \cos \phi, y = \rho \sin \phi, z = z$$

and

$$J = \frac{\partial(x, y, z)}{\partial(\rho, \phi, z)} = \rho \quad [\text{Ex. 5.25}]$$

$$\text{Then } \iiint_{R_{xyz}} f(x, y, z) dx dy dz = \iiint_{R'_{\rho\phi z}} f(\rho \cos \phi, \rho \sin \phi, z) \cdot \rho d\rho d\phi dz.$$

(iii) To change rectangular coordinates  $(x, y, z)$  to spherical polar coordinates  $(r, \theta, \phi)$ —Fig. 8.28, we have

$$x = r \sin \theta \cos \phi, y = r \sin \theta \sin \phi, z = r \cos \theta$$

and

$$J = \frac{\partial(x, y, z)}{\partial(r, \theta, \phi)} = r^2 \sin \theta \quad [\text{Ex. 5.25}]$$

$$\text{Then } \iiint_{R_{xyz}} f(x, y, z) dx dy dz = \iiint_{R'_{r\theta\phi}} f(r \sin \theta \cos \phi, r \sin \theta \sin \phi, r \cos \theta) \cdot r^2 \sin \theta dr d\theta d\phi$$

**Example 7.23.** Evaluate  $\iint_R (x+y)^2 dx dy$ , where  $R$  is the parallelogram in the  $xy$ -plane with vertices  $(1, 0), (3, 1), (2, 2), (0, 1)$  using the transformation  $u = x + y$  and  $v = x - 2y$ . (U.P.T.U., 2004)

**Solution.** The region  $R$ , i.e., parallelogram  $ABCD$  in the  $xy$ -plane becomes the region  $R'$ , i.e., rectangle  $A'B'C'D'$  in the  $uv$ -plane as shown in Fig. 7.27, by taking

$$u = x + y \quad \text{and} \quad v = x - 2y \quad \dots(i)$$




Fig. 7.27

From (i), we have

$$x = \frac{1}{3}(2u + v), y = \frac{1}{3}(u - v)$$

$$\therefore \frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = -\frac{1}{3}$$

Hence, the given integral

$$= \iint_{R'} u^2 |J| du dv = \int_1^4 \int_{-2}^1 u^2 \cdot \frac{1}{3} \cdot du dv = \frac{1}{3} \left[ \frac{u^3}{3} \right]_1^4 \cdot v \Big|_{-2}^1 = 21.$$

**Example 7.24.** Evaluate  $\iint_D xy\sqrt{1-x-y} dx dy$  where  $D$  is the region bounded by  $x = 0, y = 0$  and  $x + y = 1$  using the transformation  $x + y = u, y = uv$ . (Marathwada, 2008)



**Solution.** We have  $x = u - uv$ ,  $y = uv$

$$\therefore J = \frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} \partial x / \partial u & \partial y / \partial u \\ \partial x / \partial v & \partial y / \partial v \end{vmatrix} = \begin{vmatrix} 1-v & v \\ -u & u \end{vmatrix} = u.$$

Also when  $x = 0$ ,  $u = 0$ ,  $v = 1$ ; when  $y = 0$ ,  $u = 0$ ,  $v = 0$  and when  $x + y = 1$ ,  $u = 1$

$\therefore$  the limits of  $u$  are from 0 to 1 and limits of  $v$  are from 0 to 1.

$$\begin{aligned} \text{Thus } \iint_D xy \sqrt{1-x-y} \, dx dy &= \int_0^1 \int_0^1 u(1-v) uv (1-u)^{1/2} |J| \, du dv \\ &= \int_0^1 \int_0^1 u^3 (1-u)^{1/2} v(1-v) \, du dv \\ &= \int_0^1 u^3 (1-u)^{1/2} \, du \times \int_0^1 v(1-v) \, dv \\ &= \int_0^{\pi/2} \sin^6 \theta \cos \theta \cdot 2 \sin \theta \cos \theta \, d\theta \times \left[ \frac{v^2}{2} - \frac{v^3}{3} \right]_0^1 \\ &= 2 \int_0^{\pi/2} \sin^7 \theta \cos^2 \theta \, d\theta \left( \frac{1}{6} \right) = \frac{1}{3} \cdot \frac{6 \cdot 1}{9 \cdot 7 \cdot 5 \cdot 3} = \frac{2}{945}. \end{aligned}$$

$$\begin{aligned} \text{where } u &= \sin^2 \theta \\ du &= 2 \sin \theta \cos \theta \, d\theta \\ u = 0, \theta &= 0 \\ u = 1, \theta &= \pi/2 \end{aligned}$$

**Example 7.25.** Evaluate  $\int_0^\infty \int_0^\infty e^{-(x^2+y^2)} \, dx dy$  by changing to polar coordinates.

(Anna, 2003)

Hence show that  $\int_0^\infty e^{-x^2} \, dx = \sqrt{\pi/2}$ .

(Madras, 2003; U.P.T.U., 2003; J.N.T.U., 2000)

**Solution.** The region of integration being the first quadrant of the  $xy$ -plane,  $r$  varies from 0 to  $\infty$  and  $\theta$  varies from 0 to  $\pi/2$ . Hence,

$$\begin{aligned} I &= \int_0^\infty \int_0^\infty e^{-(x^2+y^2)} \, dx dy = \int_{\theta=0}^{\pi/2} \int_{r=0}^\infty e^{-r^2} r \, dr d\theta \\ &= -\frac{1}{2} \int_0^{\pi/2} \left\{ \int_0^\infty e^{-r^2} (-2r) \, dr \right\} d\theta = -\frac{1}{2} \int_0^{\pi/2} \left[ e^{-r^2} \right]_0^\infty d\theta = \frac{1}{2} \int_0^{\pi/2} d\theta = \frac{\pi}{4}. \end{aligned} \quad \dots(i)$$

$$\text{Also } I = \int_0^\infty e^{-x^2} \, dx \times \int_0^\infty e^{-y^2} \, dy = \left\{ \int_0^\infty e^{-x^2} \, dx \right\}^2 \quad \dots(ii)$$

$$\text{Thus, from (i) and (ii), we have } \int_0^\infty e^{-x^2} \, dx = \sqrt{\pi/2}. \quad \dots(iii)$$

**Example 7.26.** Find the volume bounded by the paraboloid  $x^2 + y^2 = az$ , the cylinder  $x^2 + y^2 = 2ay$  and the plane  $z = 0$ .

**Solution.** The required volume is found by integrating  $z = (x^2 + y^2)/a$  over the circle  $x^2 + y^2 = 2ay$ .

Changing to polar coordinates in the  $xy$ -plane, we have  $x = r \cos \theta$ ,  $y = r \sin \theta$  so that  $z = r^2/a$  and the polar equation of the circle is  $r = 2a \sin \theta$ .

To cover this circle,  $r$  varies from 0 to  $2a \sin \theta$  and  $\theta$  varies from 0 to  $\pi$ . (Fig. 7.28)

Hence the required volume

$$\begin{aligned} &= \int_0^\pi \int_0^{2a \sin \theta} z \cdot r \, d\theta dr = \frac{1}{a} \int_0^\pi d\theta \int_0^{2a \sin \theta} r^3 \, dr \\ &= \frac{1}{a} \int_0^\pi d\theta \left[ \frac{r^4}{4} \right]_0^{2a \sin \theta} = 4a^3 \int_0^\pi \sin^4 \theta \, d\theta = \frac{3\pi a^3}{2}. \end{aligned}$$




Fig. 7.28

**Example 7.27.** Find, by triple integration, the volume of the sphere  $x^2 + y^2 + z^2 = a^2$ .

(Bhopal, 2009 ; Madras, 2006 ; V.T.U., 2003 S)

**Solution.** Changing to polar spherical coordinates by putting

$$x = r \sin \theta \cos \phi, y = r \sin \theta \sin \phi, z = r \cos \theta$$

we have  $dx dy dz = r^2 \sin \theta dr d\theta d\phi$ .

Also the volume of the sphere is 8 times the volume of its portion in the positive octant for which  $r$  varies from 0 to  $a$ ,  $\theta$  varies from 0 to  $\pi/2$  and  $\phi$  varies from 0 to  $\pi/2$ .

$\therefore$  volume of the sphere

$$\begin{aligned} &= 8 \int_0^a \int_0^{\pi/2} \int_0^{\pi/2} r^2 \sin \theta dr d\theta d\phi = 8 \int_0^a r^2 dr \cdot \int_0^{\pi/2} \sin \theta d\theta \cdot \int_0^{\pi/2} d\phi \\ &= 8 \cdot \left[ \frac{r^3}{3} \right]_0^a \cdot \left[ -\cos \theta \right]_0^{\pi/2} \cdot \frac{\pi}{2} = 4\pi \cdot \frac{a^3}{3} \cdot (-0 + 1) = \frac{4}{3} \pi a^3. \end{aligned}$$

**Example 7.28.** Find the volume of the portion of the sphere  $x^2 + y^2 + z^2 = a^2$  lying inside the cylinder  $x^2 + y^2 = ay$ . (Rohtak, 2003)

**Solution.** The required volume is easily found by changing to cylindrical coordinates  $(\rho, \phi, z)$ . We therefore, have

$$x = \rho \cos \phi, y = \rho \sin \phi, z = z$$

and

$$J = \frac{\partial(x, y, z)}{\partial(\rho, \phi, z)} = \rho.$$

Then the equation of the sphere becomes  $\rho^2 + z^2 = a^2$  and that of cylinder becomes  $\rho = a \sin \phi$ .

The volume inside the cylinder bounded by the sphere is twice the volume shown shaded in the Fig. 7.29 for which  $z$  varies from 0 to  $\sqrt{(a^2 - \rho^2)}$ ,  $\rho$  varies from 0 to  $a \sin \phi$  and  $\phi$  varies from 0 to  $\pi$ .




Fig. 7.29

$$\begin{aligned} \text{Hence the required volume} &= 2 \int_0^\pi \int_0^{a \sin \phi} \int_0^{\sqrt{(a^2 - \rho^2)}} \rho dz d\rho d\phi \\ &= 2 \int_0^\pi \int_0^{a \sin \phi} \rho \sqrt{(a^2 - \rho^2)} d\rho d\phi = 2 \int_0^\pi \left[ -\frac{1}{3} (a^2 - \rho^2)^{3/2} \right]_0^{a \sin \phi} d\phi \\ &= \frac{2a^3}{3} \int_0^\pi (1 - \cos^3 \phi) d\phi = \frac{2a^3}{9} (3\pi - 4). \end{aligned}$$

**Example 7.29.** Evaluate  $\int_0^1 \int_0^{\sqrt{(1-x^2)}} \int_{\sqrt{(x^2+y^2)}}^1 \frac{dz dy dx}{\sqrt{(x^2+y^2+z^2)}}$ .

(V.T.U., 2008)

**Solution.** We change to spherical polar coordinates  $(r, \theta, \phi)$ , so that

$$x = r \sin \theta \cos \phi, y = r \sin \theta \sin \phi, z = r \cos \theta$$

and

$$J = r^2 \sin \theta, x^2 + y^2 + z^2 = r^2.$$

The region of integration is common to the cone  $z^2 = x^2 + y^2$  and the cylinder  $x^2 + y^2 = 1$  bounded by the plane  $z = 1$  in the positive octant (Fig. 7.30). Hence  $\theta$  varies from 0 to  $\pi/4$ ,  $r$  varies from 0 to  $\sec \theta$  and  $\phi$  varies from 0 to  $\pi/2$ .

$\therefore$  given integral becomes

$$\begin{aligned} &\int_0^{\pi/2} \int_0^{\pi/4} \int_0^{\sec \theta} \frac{1}{r} \cdot r^2 \sin \theta dr d\theta d\phi = \int_0^{\pi/2} d\phi \int_0^{\pi/4} \left[ \frac{r^2}{2} \right]_0^{\sec \theta} \sin \theta d\theta \\ &= \frac{\pi}{2} \int_0^{\pi/4} \frac{\sec^2 \theta}{2} \sin \theta d\theta = \frac{\pi}{4} \int_0^{\pi/4} \sec \theta \tan \theta d\theta = \frac{\pi}{4} \left[ \sec \theta \right]_0^{\pi/4} = \frac{(\sqrt{2} - 1) \pi}{4}. \end{aligned}$$




Fig. 7.30



**Example 7.30.** Find the volume of the solid surrounded by the surface

$$(x/a)^{2/3} + (y/b)^{2/3} + (z/c)^{2/3} = 1.$$

(Hissar, 2005 S)

**Solution.** Changing the variables,  $x, y, z$  to  $X, Y, Z$  where,  $(x/a)^{1/3} = X, (y/b)^{1/3} = Y, (z/c)^{1/3} = Z$   
i.e.,  $x = aX^3, y = bY^3, z = cZ^3$  so that  $J = \partial(x, y, z)/\partial(X, Y, Z) = 27 abc X^2 Y^2 Z^2$ .

$$\therefore \text{required volume} = \iiint dx dy dz = 27 abc \iiint X^2 Y^2 Z^2 dX dY dZ$$

taken throughout the sphere  $X^2 + Y^2 + Z^2 = 1$ . ... (i)

Now change  $X, Y, Z$  to spherical polar coordinates  $r, \theta, \phi$  so that  $X = r \sin \theta \cos \phi, Y = r \sin \theta \sin \phi, Z = r \cos \theta$ , and  $\partial(X, Y, Z)/\partial(r, \theta, \phi) = r^2 \sin \theta$ . To describe the positive octant of the sphere (i),  $r$  varies from 0 to 1,  $\theta$  from 0 to  $\pi/2$  and  $\phi$  from 0 to  $\pi/2$ .

$$\begin{aligned} \therefore \text{required volume} &= 27abc \times 8 \int_0^1 \int_0^{\pi/2} \int_0^{\pi/2} r^2 \sin^2 \theta \cos^2 \phi \times r^2 \sin^2 \theta \sin^2 \phi \cdot r^2 \cos^2 \theta \cdot r^2 \sin \theta dr d\theta d\phi \\ &= 216 abc \int_0^1 r^8 dr \int_0^{\pi/2} \sin^4 \theta \cos^2 \theta d\theta \int_0^{\pi/2} \sin^2 \phi \cos^2 \phi d\phi = 4\pi abc/35. \end{aligned}$$

#### PROBLEMS 7.4

Evaluate the following integrals by changing to polar co-ordinates :

- $\int_0^1 \int_0^{\sqrt{1-y^2}} (x^2 + y^2) dy dx$ . (P.T.U., 2010)
- $\int_0^2 \int_0^{\sqrt{2x-x^2}} \frac{x dx dy}{x^2 + y^2}$  (Anna, 2009)
- $\int_0^{4a} \int_{y^2/4a}^y \frac{x^2 - y^2}{x^2 + y^2} dx dy$  (Mumbai, 2006)
- $\iint xy(x^2 + y^2)^{n/2} dx dy$  over the positive quadrant of  $x^2 + y^2 = 4$ , supposing  $n + 3 > 0$ . (S.V.T.U., 2007)
- $\iint \frac{dx dy}{(1 + x^2 + y^2)^2}$  over one loop of the lemniscate  $(x^2 + y^2) = x^2 - y^2$ . (Mumbai, 2007)
- Transform the following to cartesian form and hence evaluate  $\int_0^\pi \int_0^a r^3 \sin \theta \cos \theta dr d\theta$ . (P.T.U., 2005)
- $\iint y^2 dx dy$  over the area outside  $x^2 + y^2 - ax = 0$  and inside  $x^2 + y^2 - 2ax = 0$ . (Mumbai, 2006)
- By using the transformation  $x + y = u, y = uv$ , show that  $\int_0^1 \int_0^{1-x} e^{y/(x+y)} dy dx = \frac{1}{2}(e - 1)$ . (P.T.U., 2003)
- Transform  $\int_0^{\pi/2} \int_0^{\pi/2} \frac{\sin \phi}{\sin \theta} d\phi d\theta$  by the substitution  $x = \sin \phi \cos \theta, y = \sin \phi \sin \theta$  and show that its value is  $\pi$ . (U.P.T.U., 2001)

Evaluate the following integrals by changing to spherical coordinates :

- $\int_0^1 \int_0^{\sqrt{1-x^2}} \int_0^{\sqrt{1-x^2-y^2}} \frac{dx dy dz}{\sqrt{(1-x^2-y^2-z^2)}}$ . (V.T.U., 2006 ; Kottayam, 2005)
- $\iiint_V \frac{dx dy dz}{x^2 + y^2 + z^2}$  where  $V$  is the volume of the sphere  $x^2 + y^2 + z^2 = a^2$ . (Anna, 2009)
- Evaluate  $\iiint \frac{dx dy dz}{(1 + x + y + z)^3}$  over the volume of the tetrahedron  $x = 0, y = 0, z = 0, x + y + z = 1$ . (Mumbai, 2007)
- Show that  $\iiint \frac{dx dy dz}{\sqrt{(a^2 - x^2 - y^2 - z^2)}} = \frac{\pi^2 a^3}{8}$ , the integral being extended for all the values of the variables for which the expression is real. (U.T.U., 2010)
- $\iiint z^2 dx dy dz$ , taken over the volume bounded by the surfaces  $x^2 + y^2 = a^2, x^2 + y^2 = z$  and  $z = 0$ .



15. Find the volume bounded by the  $xy$ -plane, the cylinder  $x^2 + y^2 = 1$  and the plane  $x + y + z = 3$ . (I.S.M., 2001)
16. Find the volume bounded by the  $xy$ -plane, the paraboloid  $2z = x^2 + y^2$  and the cylinder  $x^2 + y^2 = 4$ . (Raipur, 2005)
17. Find the volume cut from the sphere  $x^2 + y^2 + z^2 = a^2$  by the cone  $x^2 + y^2 = z^2$ .
18. Find the volume common to the cylinders  $x^2 + y^2 = a^2$  and  $x^2 + z^2 = a^2$ . (S.V.T.U., 2006)
19. Find the volume cut off from the cylinder  $x^2 + y^2 = ax$  by the planes  $z = 0$  and  $z = x$ . (U.P.T.U., 2006)
20. Find the volume enclosed by the cylinders  $x^2 + y^2 = 2ax$  and  $z^2 = 2ax$ . (Marathwada, 2008)
21. Find the volume of the cylinder  $x^2 + y^2 - 2ax = 0$ , intercepted between the paraboloid  $x^2 + y^2 = 2az$  and the  $xy$ -plane.
22. Find the volume bounded by the cylinder  $x^2 + y^2 = 4$  and the hyperboloid  $x^2 + y^2 - z^2 = 1$ .
23. Find the volume of the region bounded by  $z = x^2 + y^2$ ,  $z = 0$ ,  $x = -a$ ,  $x = a$  and  $y = -a$ ,  $y = a$ .
24. Prove, by using a double integral that the volume generated by the revolution of the cardioid  $r = a(1 + \cos \theta)$  about its axis is  $8\pi a^3/3$ . (V.T.U., 2000)
25. Evaluate  $\iiint (x + y + z) dx dy dz$  over the tetrahedron bounded by the planes  $x = 0$ ,  $y = 0$ ,  $z = 0$  and  $x + y + z = 1$ . [See Fig. 7.34]
26. Find the volume of the tetrahedron bounded by the coordinate planes and the plane  $\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$ . (Burdwan, 2003)
27. Work out example 7.21 by changing the variables.

## 7.8 AREA OF A CURVED SURFACE

Consider a point  $P$  of the surface  $S : z = f(x, y)$ . Let its projection on the  $xy$ -plane be the region  $A$ . Divide it into area elements by drawing lines parallel to the axes of  $X$  and  $Y$ . (Fig. 7.31).

On the element  $\delta x \delta y$  as base, erect a cylinder having generators parallel to  $OZ$  and meeting the surface  $S$  in an element of area  $\delta S$ .

As  $\delta x \delta y$  is the projection of  $\delta S$  on the  $xy$ -plane,

$\therefore \delta x \delta y = \delta S \cdot \cos \gamma$ , where  $\gamma$  is the angle between the  $xy$ -plane and the tangent plane to  $S$  at  $P$ , i.e., it is the angle between the  $Z$ -axis and the normal to  $S$  at  $P (= \angle Z'PN)$ .

Now since the direction cosines of the normal to the surface  $F(x, y, z) = 0$  proportional to

$$\frac{\partial F}{\partial x}, \frac{\partial F}{\partial y}, \frac{\partial F}{\partial z}.$$

$\therefore$  the direction cosines of the normal to  $S [F = f(x, y) - z]$  are proportional to  $-\frac{\partial z}{\partial x}, -\frac{\partial z}{\partial y}, 1$  and those of the  $z$ -axis are  $0, 0, 1$ .




Fig. 7.31

$$\text{Hence } \cos \gamma = \frac{1}{\sqrt{\left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2 + 1}} \quad \therefore \delta S = \frac{\delta x \delta y}{\cos \gamma} = \sqrt{\left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2 + 1} \delta x \delta y$$

$$\text{Hence } S = \lim_{\delta S \rightarrow 0} \sum \delta S = \iint_A \sqrt{\left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2 + 1} dx dy$$

Similarly, if  $B$  and  $C$  be the projections of  $S$  on the  $yz$ - and  $zx$ -planes respectively, then

$$S = \iint_B \sqrt{\left(\frac{\partial x}{\partial y}\right)^2 + \left(\frac{\partial x}{\partial z}\right)^2 + 1} dy dz$$

and

$$S = \iint_C \sqrt{\left(\frac{\partial y}{\partial z}\right)^2 + \left(\frac{\partial y}{\partial x}\right)^2 + 1} dz dx.$$



**Example 7.31.** Find the area of the portion of the cylinder  $x^2 + z^2 = 4$  lying inside the cylinder  $x^2 + y^2 = 4$ .

**Solution.** Figure 7.32 shows one-eighth of the required area. Its projection on the  $xy$ -plane is a quadrant circle  $x^2 + y^2 = 4$ .

For the cylinder  $x^2 + z^2 = 4$ , ... (i)

we have  $\frac{\partial z}{\partial x} = -\frac{x}{z}$ ,  $\frac{\partial z}{\partial y} = 0$ .

so that  $\left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2 + 1 = \frac{x^2 + z^2}{z^2} = \frac{4}{4 - x^2}$ .

Hence the required surface area = 8 (surface area of the upper portion of (i) lying within the cylinder  $x^2 + y^2 = 4$  in the positive octant)

$$= 8 \int_0^2 \int_0^{\sqrt{4-x^2}} \frac{2}{\sqrt{4-x^2}} dx dy = 16 \int_0^2 dx = 32 \text{ sq. units.}$$




Fig. 7.32

**Example 7.32.** Find the area of the portion of the sphere  $x^2 + y^2 + z^2 = 9$  lying inside the cylinder  $x^2 + y^2 = 3y$ .

**Solution.** Figure 7.33 shows one-fourth of the required area. Its projection on the  $xy$ -plane is the semi-circle  $x^2 + y^2 = 3y$  bounded by the  $Y$ -axis.

For the sphere

$$x^2 + y^2 + z^2 = 9, \quad \frac{\partial z}{\partial x} = -\frac{x}{z} \quad \text{and} \quad \frac{\partial z}{\partial y} = -\frac{y}{z}$$

$$\therefore \left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2 + 1 = (x^2 + y^2 + z^2)/z^2$$

$$= \frac{9}{9 - x^2 - y^2} = \frac{9}{9 - r^2} \quad \text{when } x = r \cos \theta, y = r \sin \theta.$$

Using polar coordinates, the required area is found by integrating  $3/\sqrt{9-r^2}$  over the semi-circle  $r = 3 \sin \theta$ , for which  $r$  varies from 0 to  $3 \sin \theta$  and  $\theta$  varies from 0 to  $\pi/2$ .

Hence the required surface area

$$\begin{aligned} &= 4 \int_0^{\pi/2} \int_0^{3 \sin \theta} \frac{3}{\sqrt{9-r^2}} r d\theta dr = -6 \int_0^{\pi/2} \left| \frac{\sqrt{9-r^2}}{1/2} \right|_0^{3 \sin \theta} d\theta \\ &= 36 \int_0^{\pi/2} (1 - \cos \theta) d\theta = 36 \left| \theta - \sin \theta \right|_0^{\pi/2} = 18 (\pi - 2) \text{ sq. units.} \end{aligned}$$




Fig. 7.33

### PROBLEMS 7.5

1. Show that the surface area of the sphere  $x^2 + y^2 + z^2 = a^2$  is  $4\pi a^2$ .
2. Find the area of the portion of the cylinder  $x^2 + y^2 = 4y$  lying inside the sphere  $x^2 + y^2 + z^2 = 16$ .
3. Find the area of the portion of the sphere  $x^2 + y^2 + z^2 = a^2$  lying inside the cylinder  $x^2 + y^2 = ax$ .
4. Find the area of the surface of the cone  $x^2 + y^2 = z^2$  cut off by the surface of the cylinder  $x^2 + y^2 = a^2$  above the  $xy$ -plane.
5. Compute the area of that part of the plane  $x + y + z = 2a$  which lies in the first octant and is bounded by the cylinder  $x^2 + y^2 = a^2$ .

(Burdwan, 2003)

### 7.9 CALCULATION OF MASS

(a) **For a plane lamina**, if the surface density at the point  $P(x, y)$  be  $\rho = f(x, y)$  then the elementary mass at  $P = \rho \delta x \delta y$ .

$$\therefore \text{total mass of the lamina} = \iint \rho \, dx \, dy \quad \dots(i)$$

with integrals embracing the whole area of the lamina.

In polar coordinates, taking  $\rho = \phi(r, \theta)$  at the point  $P(r, \theta)$ ,

$$\text{total mass of the lamina} = \iint \rho r \, d\theta \, dr \quad \dots(ii)$$

(b) **For a solid**, if the density at the point  $P(x, y, z)$  be  $\rho = f(x, y, z)$ , then

$$\text{total mass of the solid} = \iiint \rho \, dx \, dy \, dz \text{ with appropriate limits of integration.}$$

**Example 7.33.** Find the mass of the tetrahedron bounded by the coordinates planes and the plane

$$\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1, \text{ the variable density } \rho = \mu xyz. \quad (\text{Rohtak, 2003 ; U.P.T.U., 2003})$$

**Solution.** Elementary mass at  $P = \mu xyz \cdot \delta x \delta y \delta z$ .

$$\therefore \text{the whole mass} = \iiint \mu xyz \, dx \, dy \, dz,$$

the integrals embracing the whole volume  $OABC$  (Fig. 7.34). The limits for  $z$  are from 0 to  $z = c(1 - x/a - y/b)$ .

The limits for  $y$  are from 0 to  $y = b(1 - x/a)$  and limits for  $x$  are from 0 to  $a$ .

Hence the required mass

$$\begin{aligned} &= \int_0^a \int_0^{b(1-x/a)} \int_0^{c(1-x/a-y/b)} \mu xyz \, dz \, dy \, dx \\ &= \mu \int_0^a \int_0^{b(1-x/a)} xy \left[ \frac{z^2}{2} \right]_0^{c(1-x/a-y/b)} dy \, dx \\ &= \mu \int_0^a \int_0^{b(1-x/a)} xy \cdot \frac{c^2}{2} \left( 1 - \frac{x}{a} - \frac{y}{b} \right)^2 dy \, dx \\ &= \frac{\mu c^2}{2} \int_0^a \int_0^{b(1-x/a)} x \cdot \left[ \left( 1 - \frac{x}{a} \right)^2 y - 2 \left( 1 - \frac{x}{a} \right) \frac{y^2}{b} + \frac{y^3}{b^2} \right] dy \, dx \\ &= \frac{\mu c^2}{2} \int_0^a x \left[ \left( 1 - \frac{x}{a} \right)^2 \frac{y^2}{2} - 2 \left( 1 - \frac{x}{a} \right) \frac{y^3}{3b} + \frac{y^4}{4b^2} \right]_0^{b(1-x/a)} dx \\ &= \frac{\mu c^2}{2} \int_0^a x \left[ \frac{b^2}{2} \left( 1 - \frac{x}{a} \right)^4 - \frac{2b^2}{3} \left( 1 - \frac{x}{a} \right)^4 + \frac{b^2}{4} \left( 1 - \frac{x}{a} \right)^4 \right] dx = \frac{\mu b^2 c^2}{24} \int_0^a x \left( 1 - \frac{x}{a} \right)^4 dx = \frac{\mu a^2 b^2 c^2}{720}. \end{aligned}$$




Fig. 7.34

### 7.10 CENTRE OF GRAVITY

(a) **To find the C.G.  $(\bar{x}, \bar{y})$  of a plane lamina**, take the element of mass  $\rho \delta x \delta y$  at the point  $P(x, y)$ . Then

$$\bar{x} = \frac{\iint x \rho \, dx \, dy}{\iint \rho \, dx \, dy}, \bar{y} = \frac{\iint y \rho \, dx \, dy}{\iint \rho \, dx \, dy} \text{ with integrals embracing the whole lamina.}$$

While using polar coordinates, take the elementary mass as  $\rho r \delta \theta \delta r$  at the point  $P(r, \theta)$  so that  $x = r \cos \theta$ ,  $y = r \sin \theta$ .

$$\therefore \bar{x} = \frac{\iint r \cos \theta \rho r \, d\theta \, dr}{\iint \rho r \, d\theta \, dr}, \bar{y} = \frac{\iint r \sin \theta \rho r \, d\theta \, dr}{\iint \rho r \, d\theta \, dr}$$



(b) To find the C.G.  $(\bar{x}, \bar{y}, \bar{z})$  of a solid, take an element of mass  $\rho \delta x \delta y \delta z$  enclosing the point  $P(x, y, z)$ .

Then

$$\bar{x} = \frac{\iiint x \rho \, dx dy dz}{\iiint \rho \, dx dy dz}, \quad \bar{y} = \frac{\iiint y \rho \, dx dy dz}{\iiint \rho \, dx dy dz} \quad \text{and} \quad \bar{z} = \frac{\iiint z \rho \, dx dy dz}{\iiint \rho \, dx dy dz}.$$

**Example 7.34.** Find by double integration, the centre of gravity of the area of the cardioid  $r = a(1 + \cos \theta)$ .

**Solution.** The cardioid being symmetrical about the initial line, its C.G. lies on OX, i.e.,  $\bar{y} = 0$  (Fig. 7.35).




Fig. 7.35

$$\begin{aligned} \therefore \bar{x} &= \frac{\iint \rho r \cos \theta \cdot r d\theta dr}{\iint \rho r d\theta dr} = \frac{\int_{-\pi}^{\pi} \int_0^{a(1+\cos \theta)} \cos \theta \cdot r^2 dr \cdot d\theta}{\int_{-\pi}^{\pi} \int_0^{a(1+\cos \theta)} r dr \cdot d\theta} \\ &= \frac{\int_{-\pi}^{\pi} \cos \theta \left[ \frac{r^3}{3} \right]_0^{a(1+\cos \theta)} d\theta}{\int_{-\pi}^{\pi} \left[ \frac{r^2}{2} \right]_0^{a(1+\cos \theta)} d\theta} = \frac{2a}{3} \cdot \frac{\int_{-\pi}^{\pi} \cos \theta (1 + \cos \theta)^3 d\theta}{\int_{-\pi}^{\pi} (1 + \cos \theta)^2 d\theta} \\ &= \frac{2a}{3} \cdot \frac{2 \cdot \int_0^{\pi} (3 \cos^2 \theta + \cos^4 \theta) d\theta}{2 \cdot \int_0^{\pi} (1 + \cos^2 \theta) d\theta} \quad \left\{ \because \int_{-\pi}^{\pi} \cos^n \theta d\theta = 2 \int_0^{\pi} \cos^n \theta d\theta \text{ or } 0 \right. \\ &\quad \left. \text{according as } n \text{ is even or odd.} \right\} \\ &= \frac{2a}{3} \cdot \frac{2 \cdot \int_0^{\pi/2} (3 \cos^2 \theta + \cos^4 \theta) d\theta}{2 \cdot \int_0^{\pi/2} (1 + \cos^2 \theta) d\theta} \quad \left( \begin{array}{l} \text{as the powers of} \\ \cos \theta \text{ are even} \end{array} \right) = \frac{2a}{3} \cdot \frac{3 \cdot \frac{1}{2} \cdot \frac{\pi}{2} + \frac{3}{4} \cdot \frac{1}{2} \cdot \frac{\pi}{2}}{\frac{\pi}{2} + \frac{1}{2} \cdot \frac{\pi}{2}} = \frac{5a}{6} \end{aligned}$$

Hence the C.G. of the cardioid is at  $G(5a/6, 0)$ .

**Example 7.35.** Using double integration, find the C.G. of a lamina in the shape of a quadrant of the curve  $(x/a)^{2/3} + (y/b)^{2/3} = 1$ , the density being  $\rho = kxy$ , where  $k$  is a constant.

**Solution.** Let  $G(\bar{x}, \bar{y})$  be the C.G. of the lamina OAB (Fig. 7.36), so that

$$\bar{x} = \frac{\iint kxy \cdot x dx dy}{\iint kxy \cdot dx dy} = \frac{\iint x^2 y dx dy}{\iint xy dx dy}$$

where the integrals are taken over the area OAB so that  $y$  varies from 0 to  $y$  (to be found from the equation of the curve in terms of  $x$ ) and then  $x$  varies from 0 to  $a$ .

Thus

$$\bar{x} = \frac{\int_0^a \int_0^y x^2 y dy dx}{\int_0^a \int_0^y xy dy dx} = \frac{\int_0^a x^2 \cdot \left[ y^2/2 \right]_0^y dx}{\int_0^a x \cdot \left[ y^2/2 \right]_0^y dx} = \frac{\int_0^a x^2 y^2 dx}{\int_0^a xy^2 dx}$$

For any point on the curve, we have

$$x = a \cos^3 \theta, y = b \sin^3 \theta \text{ so that} \\ dx = -3a \cos^2 \theta \sin \theta d\theta.$$

Also when  $x = 0, \theta = \pi/2$ ; when  $x = a, \theta = 0$ .




Fig. 7.36

Hence

$$\bar{x} = \frac{\int_{-\pi/2}^0 a^2 \cos^6 \theta \cdot b^2 \sin^6 \theta \cdot (-3a \cos^2 \theta \sin \theta) d\theta}{\int_{-\pi/2}^0 a \cos^3 \theta \cdot b^2 \sin^6 \theta \cdot (-3a \cos^2 \theta \sin \theta) d\theta}$$

$$= a \frac{\int_0^{\pi/2} \sin^7 \theta \cos^8 \theta d\theta}{\int_0^{\pi/2} \sin^7 \theta \cos^5 \theta d\theta} = \frac{128}{429} a$$

Similarly,

$$\bar{y} = \frac{\int_0^a \int_0^y kxy \cdot y \, dx dy}{\int_0^a \int_0^y kxy \cdot dx dy} = \frac{128}{429} b.$$

Hence the required C.G. is  $G \left( \frac{128}{429} a, \frac{128}{429} b \right)$ .

### 7.11 CENTRE OF PRESSURE

Consider plane area  $A$  immersed vertically in a homogeneous liquid. Take the line of intersection of the given plane with the free surface of the liquid as the  $x$ -axis and any line lying in this plane and perpendicular to it downwards as the  $y$ -axis (Fig. 7.37).

If  $p$  be the pressure at the point  $P(x, y)$  of the area  $A$ , then the pressure on an elementary area  $\delta x \delta y$  at  $P$  is  $p \delta x \delta y$  which is normal to the plane.

$\therefore$  the resultant pressure on  $A = \iint p \, dx dy$ .

If this resultant pressure acting at  $C(h, k)$  is equivalent to pressure at various points such as  $p \delta x \delta y$  distributed over the whole area  $A$ , then  $C$  is called the *centre of pressure*.

$\therefore$  taking the moment of the resultant pressure at  $C$  and the sum of the moments of the individual pressures such as  $p \delta x \delta y$  at  $P(x, y)$  about the  $y$ -axis, we get

$$h \iint p \, dx dy = \iint x \cdot p \, dx dy, \quad \text{i.e., } h = \frac{\iint x \cdot dx dy}{\iint p \, dx dy}$$

Similarly, taking moments about  $x$ -axis, we have

$$k = \frac{\iint y \cdot p \, dx dy}{\iint p \, dx dy} \quad \text{with integrals embracing the whole of the area } A.$$

While using polar coordinates, replace  $x$  by  $r \cos \theta$ ,  $y$  by  $r \sin \theta$  and  $dx dy$  by  $r \, d\theta dr$  in the above formulae.

**Example 7.36.** A horizontal boiler has a flat bottom and its ends are plane and semi-circular. If it is just full of water, show that the depth of the centre of pressure of either end is  $0.7 \times$  total depth approximately.

**Solution.** Let the semi-circle  $x^2 + y^2 = a^2$  represent an end of the given boiler (Fig. 7.38). By symmetry, its centre of pressure lies on  $OY$ .

If  $w$  be the weight of water per unit volume, then the pressure  $p$  at the point  $P(x, y) = w(a - y)$ .

$\therefore$  the height  $k$  of the C.P. above  $OX$ , is given by

$$k = \frac{\iint y \cdot p \, dx dy}{\iint p \, dx dy} = \frac{\int_{-a}^a \int_0^{\sqrt{a^2 - x^2}} w(a - y) y \, dy \cdot dx}{\int_{-a}^a \int_0^{\sqrt{a^2 - x^2}} w(a - y) \, dy \cdot dx}$$

$$= \frac{\int_{-a}^a \left[ ay^2/2 - y^3/3 \right]_0^{\sqrt{a^2 - x^2}} dx}{\int_{-a}^a \left[ ay - y^2/2 \right]_0^{\sqrt{a^2 - x^2}} dx} = \frac{\int_{-a}^a \left[ \frac{a}{2} (a^2 - x^2) - \frac{1}{3} (a^2 - x^2)^{3/2} \right] dx}{\int_{-a}^a \left[ a(a^2 - x^2)^{1/2} - \frac{1}{2} (a^2 - x^2) \right] dx}$$




Fig. 7.37




Fig. 7.38



Now put  $x = a \sin \theta$ , so that  $dx = a \cos \theta d\theta$ .

Also when  $x = -a$ ,  $\theta = -\pi/2$ , and when  $x = a$ ,  $\theta = \pi/2$ .

$$\begin{aligned} \therefore k &= \frac{\int_{-\pi/2}^{\pi/2} \left[ \frac{a^3}{2} \cos^2 \theta - \frac{a^3}{3} \cos^3 \theta \right] a \cos \theta d\theta}{\int_{-\pi/2}^{\pi/2} \left[ a^2 \cos \theta - \frac{a^2}{2} \cos^2 \theta \right] a \cos \theta d\theta} \\ &= \frac{\frac{a}{3} \cdot \frac{2}{2} \int_0^{\pi/2} (3 \cos^3 \theta - 2 \cos^4 \theta) d\theta}{\frac{2}{2} \int_0^{\pi/2} (2 \cos^2 \theta - \cos^3 \theta) d\theta} = \frac{a}{4} \left( \frac{16 - 3\pi}{3\pi - 4} \right) = 0.3a \text{ nearly.} \end{aligned}$$

Hence, the depth of the C.P.  $= a - k = 0.7a$  approximately.

### PROBLEMS 7.6

1. A lamina is bounded by the curves  $y = x^2 - 3x$  and  $y = 2x$ . If the density at any point is given by  $\lambda xy$ , find by double integration, the mass of the lamina.
2. Find the mass of a lamina in the form of cardioid  $r = a(1 + \cos \theta)$  whose density at any point varies as the square of its distance from the initial line.
3. Find the mass of a solid in the form of the positive octant of the sphere  $x^2 + y^2 + z^2 = 9$ , if the density at any point is  $2xyz$ .
4. Find the centroid of the area enclosed by the parabola  $y^2 = 4ax$ , the axis of  $x$  and its latus-rectum.
5. The density at any point  $(x, y)$  of a lamina is  $\sigma(x + y)/a$  where  $\sigma$  and  $a$  are constants. The lamina is bounded by the lines  $x = 0$ ,  $y = 0$ ,  $x = a$ ,  $y = b$ . Find the position of its centre of gravity.
6. Find the centroid of a loop of the lemniscate  $r^2 = a^2 \cos 2\theta$ .
7. A plane in the form of a quadrant of the ellipse  $(x/a)^2 + (y/b)^2 = 1$  is of small but varying thickness, the thickness at any point being proportional to the product of the distances of that point from the axes; show that the coordinates of the centroid are  $(8a/15, 8b/15)$ . (Nagpur, 2009)
8. In a semi-circular disc bounded by a diameter  $OA$ , the density at any point varies as the distance from  $O$ ; find the position of the centre of gravity.
9. Find the centroid of the tetrahedron bounded by the coordinate planes and the plane  $x + y + z = 1$ , the density at any point varying as its distance from the face  $z = 0$ .
10. Find  $\bar{x}$  where  $(\bar{x}, \bar{y}, \bar{z})$  is the centroid of the region  $R$  bounded by the parabolic cylinder  $z = 4 - x^2$  and the planes  $x = 0$ ,  $y = 0$ ,  $y = 6$ ,  $z = 0$ . (Assume that the density is constant).
11. If the density at any point of the solid octant of the ellipsoid  $(x/a)^2 + (y/b)^2 + (z/c)^2 = 1$  varies as  $xyz$ , find the coordinates of the C.G. of the solid. (P.T.U., 2005)
12. A horizontal boiler has a flat bottom and its ends consist of a square 1 metre wide surmounted by an isosceles triangle of height 0.5 metre. Determine the depth of the centre of pressure of either end when the boiler is just full.
13. A quadrant of a circle is just, immersed vertically in a heavy homogeneous liquid with one edge in the surface. Find the centre of pressure.
14. Find the depth of the centre of pressure of a square lamina immersed in the liquid, with one vertex in the surface and the diagonal vertical.
15. Find the centre of pressure of a triangular lamina immersed in a homogeneous liquid with one side in the free surface. (P.T.U., 2003)
16. A uniform semi-circular is lamina immersed in a fluid with its plane vertical and its boundary diameter on the free surface. If the density at any point of the fluid varies as the depth of the point below the free surface, find the position of the centre of pressure of the lamina.

### 7.12 (1) MOMENT OF INERTIA

If a particle of mass  $m$  of a body be at a distance  $r$  from a given line, then  $mr^2$  is called the *moment of inertia of the particle about the given line* and the sum of similar expressions taken for all the particles of the body, i.e.,  $\sum mr^2$  is called the *moment of inertia of the body about the given line* (Fig. 7.39).



If  $M$  be the total mass of the body and we write its moment of inertia  $= Mk^2$ , then  $k$  is called the *radius of gyration* of the body about the axis.

(2) **M.I. of plane lamina.** Consider the elementary mass  $\rho \delta x \delta y$  at the point  $P(x, y)$  of a plane area  $A$  so that its M.I. about  $x$ -axis  $= \rho \delta x \delta y y^2$ .

$$\therefore \text{M.I. of the lamina about } x\text{-axis, i.e. } I_x = \iint_A \rho y^2 dx dy.$$

$$\text{Similarly, M.I. of the lamina about } y\text{-axis, i.e. } I_y = \iint_A \rho x^2 dx dy.$$

Also M.I. of the lamina about an axis perpendicular to the  $xy$ -plane, i.e.,

$$I_z = \iint_A \rho (x^2 + y^2) dx dy.$$

(3) **M.I. of a solid.** Consider an elementary mass  $\rho \delta x \delta y \delta z$  enclosing a point  $P(x, y, z)$  of a solid of volume  $V$ .

$$\text{Distance of } P \text{ from the } x\text{-axis} = \sqrt{(y^2 + z^2)}.$$

$$\therefore \text{M.I. of this element about the } x\text{-axis} = \rho \delta x \delta y \delta z (y^2 + z^2).$$

$$\text{Thus M.I. of this solid about } x\text{-axis, i.e., } I_x = \iiint_V \rho (y^2 + z^2) dx dy dz.$$

$$\text{Similarly, its M.I. about } y\text{-axis, i.e., } I_y = \iiint_V \rho (z^2 + x^2) dx dy dz$$

$$\text{and M.I. about } z\text{-axis, i.e., } I_z = \iiint_V \rho (x^2 + y^2) dx dy dz.$$

(4) Sometimes we require the moment of inertia of a body about axes other than the principal axes. The following theorems prove useful for this purpose :

**I. Theorem of perpendicular axis.** If the moment of inertia of a lamina about two perpendicular axes  $OX, OY$  in its plane are  $I_x$  and  $I_y$ , then the moment of inertia of the lamina about an axis  $OZ$ , perpendicular to it is given by  $I_z = I_x + I_y$ .

Its proof follows from the relations giving  $I_x, I_y$  and  $I_z$  for a plane lamina [(2) above].

**II. Steiner's theorem\*.** If the moment of inertia of a body of mass  $M$  about an axis through its centre of gravity is  $I$ , then  $I'$ , moment of inertia about a parallel axis at a distance  $d$  from the first axis, is given by  $I' = I + Md^2$ .

Its proof will be found in any text book on Dynamics of a Rigid Body.

**Example 7.37.** Find the M.I. of the area bounded by the curve  $r^2 = a^2 \cos 2\theta$  about its axis.

**Solution.** Given curve is symmetrical about the pole and for half of the loop in the first quadrant  $\theta$  varies from 0 to  $\pi/4$  (Fig. 7.40).

Elementary area at  $P(r, \theta) = r d\theta dr$ .

If  $\rho$  be the surface density, then elementary mass

$$= \rho r d\theta dr \quad \dots(i)$$

$$\therefore \text{its total mass } M = 4 \int_0^{\pi/4} \int_0^{a\sqrt{\cos 2\theta}} \rho r dr d\theta$$

$$= 2\rho a^2 \int_0^{\pi/4} \cos 2\theta d\theta = \rho a^2 \quad \dots(ii)$$

Now M.I. of the elementary mass (i) about the  $x$ -axis.

$$= \rho r d\theta dr \cdot y^2 = \rho r d\theta dr (r \sin \theta)^2 = \rho r^3 \sin^2 \theta dr d\theta$$

Hence the M.I. of the whole area

$$= 4 \int_0^{\pi/4} \int_0^{a\sqrt{\cos 2\theta}} \rho r^3 \sin^2 \theta dr d\theta = 4\rho \int_0^{\pi/4} \sin^2 \theta \cdot \left[ \frac{r^4}{4} \right]_0^{a\sqrt{\cos 2\theta}} d\theta$$

$$= \rho a^2 \int_0^{\pi/4} \cos^2 2\theta \cdot \sin^2 \theta d\theta = \rho a^4 \int_0^{\pi/2} \cos^2 \phi \cdot \sin^2 \frac{\phi}{2} \cdot \frac{d\phi}{2} \quad [\text{Put } 2\theta = \phi, d\theta = d\phi/2]$$

$$= \frac{\rho a^4}{4} \int_0^{\pi/2} (\cos^2 \phi - \cos^3 \phi) d\phi = \frac{\rho a^4}{48} (3\pi - 8) = \frac{Ma^2}{48} (3\pi - 8). \quad [\text{By (ii)}]$$




Fig. 7.39




Fig. 7.40

\*Named after a Swiss geometrer Jacob Steiner (1796–1863) who was a professor at Berlin University.



**Example 7.38.** Find the moment of inertia of a hollow sphere about a diameter, its external and internal radii being 5 metres and 4 metres.

**Solution.** Let  $\rho$  be the density of the given hollow sphere. Then the M.I. about the diameter, i.e.,  $x$ -axis is

$$I_x = \iiint_V \rho(y^2 + z^2) dx dy dz$$

Changing to polar spherical coordinates, we get

$$\begin{aligned} I_x &= \int_0^{2\pi} \int_0^\pi \int_4^5 \rho \{ (r \sin \theta \sin \phi)^2 + (r \cos \theta)^2 \} r^2 \sin \theta dr d\theta d\phi \\ &= \rho \left\{ \int_0^{2\pi} \sin^2 \phi d\phi \cdot \int_0^\pi \sin^3 \theta d\theta \left[ \frac{r^5}{5} \right]_4^5 + \int_0^{2\pi} d\phi \int_0^\pi \cos^2 \theta \sin \theta d\theta \left[ \frac{r^5}{5} \right]_4^5 \right\} \\ &= \frac{8\pi\rho}{15} (5^5 - 4^5) = 1120.5 \text{ m.} \end{aligned}$$

**Example 7.39.** A solid body of density  $\rho$  is in the shape of the solid formed by revolution of the centroid  $r = a(1 + \cos \theta)$  about the initial line. Show that its moment of inertia about a straight line through the pole perpendicular to the initial line is  $\frac{352}{105} \pi \rho a^5$ . (U.P.T.U., 2001)

**Solution.** An elementary area  $rd\theta dr$ , when revolved about  $OX$  generates a circular ring of radius  $LP = r \sin \theta$  (Fig. 7.41).

M.I. of this ring about a diameter parallel to  $OY$

$$= (2\pi r \sin \theta) (rd\theta dr) \rho \cdot \frac{(r \sin \theta)^2}{2}.$$

[ $\therefore$  M.I. of a ring about a diameter =  $Ma^2/2$ .]

Now using Steiner's theorem, we have M.I. of the ring about  $OY$  = M.I. of the ring about a diameter  $LP$  parallel to  $OY$  + Mass of the ring  $(OL)^2 (r \cos \theta)^2$

$$= 2\pi\rho r^4 \sin^3 \theta d\theta dr + 2\pi r \sin \theta (rd\theta dr) (r \cos \theta)^2$$

Hence M.I. of the solid generated by revolution about  $OY$

$$\begin{aligned} &= \pi\rho \int_0^\pi \int_0^{r=a(1+\cos\theta)} (r^4 \sin^3 \theta + 2r^4 \sin \theta \cos^2 \theta) d\theta dr \\ &= \pi\rho \int_0^\pi (\sin^3 \theta + 2 \sin \theta \cos^2 \theta) d\theta \int_0^{a(1+\cos\theta)} r^4 dr \\ &= \frac{\pi\rho a^5}{5} \int_0^\pi \sin \theta (1 + \cos^2 \theta) (1 + \cos \theta)^5 d\theta \\ &= \frac{\pi\rho a^5}{5} \int_0^{\pi/2} \sin 2\phi (1 + \cos^2 2\phi) (1 + \cos 2\phi)^5 2d\phi \\ &= \frac{\pi\rho a^5}{5} \int_0^{\pi/2} 2 \sin \phi \cos \phi \{1 + (2 \cos^2 \phi - 1)^2\} (2 \cos^2 \phi)^5 2d\phi \\ &= \frac{256 \pi\rho a^5}{5} \int_0^{\pi/2} (\cos^{11} \phi - 2 \cos^{13} \phi + 2 \cos^{15} \phi) \sin \phi d\phi \\ &= \frac{256 \pi\rho a^5}{5} \left[ -\frac{\cos^{12} \phi}{12} + \frac{2 \cos^{14} \phi}{14} - \frac{2 \cos^{16} \phi}{16} \right]_0^{\pi/2} = \frac{352 \pi\rho a^5}{105}. \end{aligned}$$




Fig. 7.41

[Put  $\theta = 2\phi$ ]

**Example 7.40.** A hemisphere of radius  $R$  has a cylindrical hole of radius  $a$  drilled through it, the axis of the hole being along the radius normal to the plane face of the hemisphere. Find its radius of gyration about a diameter of this face.

**Solution.** M.I. of the given solid about x-axis

$$= \iiint \rho(y^2 + z^2) dx dy dz$$

The limits of integration for  $z$  are from 0 to  $z = \sqrt{R^2 - x^2 - y^2}$  found from the equation of the sphere  $x^2 + y^2 + z^2 = R^2$ . The limits for  $x$  and  $y$  are to be such as to cover the shaded area  $A$  in the  $xy$ -plane between the concentric circles of radii  $a$  and  $R$  (Fig. 7.42).

Thus the required M.I. about x-axis

$$\begin{aligned} &= \rho \iint_A \int_0^{\sqrt{R^2 - x^2 - y^2}} (y^2 + z^2) dz dx dy \\ &= \rho \iint_A \left[ y^2 z + z^3 / 3 \right]_0^{\sqrt{R^2 - x^2 - y^2}} dx dy = \rho \iint_A \left[ y^2 (R^2 - x^2 - y^2)^{1/2} + \frac{1}{3} (R^2 - x^2 - y^2)^{3/2} \right] dx dy. \end{aligned}$$

Now changing to polar coordinates, we have  $x = r \cos \theta$ ,  $y = r \sin \theta$  and  $dx dy = r dr d\theta$ .

Also to cover the area  $A$ ,  $r$  varies from  $a$  to  $R$  and  $\theta$  varies from 0 to  $2\pi$ .

Hence the required M.I. about x-axis

$$\begin{aligned} &= \rho \int_a^R \int_0^{2\pi} \left[ r^2 \sin^2 \theta \cdot (R^2 - r^2)^{1/2} + \frac{1}{3} (R^2 - r^2)^{3/2} \right] r d\theta dr \\ &= \rho \int_a^R \int_0^{2\pi} \left[ \frac{1}{2} r^2 (1 - \cos 2\theta) + \frac{1}{3} (R^2 - r^2) \right] d\theta \cdot r (R^2 - r^2)^{1/2} dr \\ &= \rho \int_a^R \left[ \frac{r^2}{2} \left( \theta - \frac{\sin 2\theta}{2} \right) + \frac{1}{3} (R^2 - r^2) \theta \right]_0^{2\pi} \cdot r (R^2 - r^2)^{1/2} dr \\ &= \rho \int_a^R 2\pi \left( \frac{r^2}{2} + \frac{R^2 - r^2}{3} \right) \cdot r (R^2 - r^2)^{1/2} dr \\ &= \frac{\pi \rho}{3} \int_a^R (2R^2 + r^2)(R^2 - r^2)^{1/2} \cdot r dr && [\text{Put } r^2 = t \text{ and } r dr = dt/2] \\ &= \frac{\pi \rho}{6} \int_{a^2}^{R^2} (2R^2 + t)(R^2 - t)^{1/2} dt && [\text{Integrate by parts}] \\ &= \frac{\pi \rho}{9} \left[ (2R^2 + a^2)(R^2 - a^2)^{3/2} + \frac{2}{5} (R^2 - a^2)^{5/2} \right] = \frac{2\pi \rho}{3} (R^2 - a^2)^{3/2} \times \frac{1}{10} (4R^2 + a^2) \\ &\quad \left[ \because \text{Mass} = \rho \int_0^{2\pi} \int_a^R \int_0^{\sqrt{R^2 - r^2}} dz \cdot r dr \cdot d\theta = \frac{2\pi \rho}{3} (R^2 - a^2)^{3/2} \right] \end{aligned}$$

Hence, the radius of gyration  $= [(4R^2 + a^2)/10]^{1/2}$ .




Fig. 7.42

### 7.13 (1) PRODUCT OF INERTIA

If a particle of mass  $m$  of a body be at distances  $x$  and  $y$  from two given perpendicular lines, then  $\sum mxy$  is called the *product of inertia* of the body about the given lines.

Consider an elementary mass  $\delta x \delta y \delta z$  enclosing the point  $P(x, y, z)$  of solid of volume  $V$ . Then the product of inertia (P.I.) of this element about the axes of  $x$  and  $y$  is  $\rho \delta x \delta y \delta z xy$ .

$$\therefore \text{P.I. of the solid about } x \text{ and } y\text{-axes, i.e., } P_{xy} = \iiint_V \rho xy dx dy dz$$

$$\text{Similarly, } P_{yz} = \iiint_V \rho yz dx dy dz \text{ and } P_{zx} = \iiint_V \rho zx dx dy dz.$$

In particular, for a plane lamina of surface density  $\rho$  and covering a region  $A$  in the  $xy$ -plane,

$$P_{xy} = \iint_A \rho xy dx dy \text{ whereas } P_{yz} = P_{zx} = 0. \quad [\because z = 0]$$



**(2) Principal axes.** The principal axes of a lamina at a given point are that pair of axes in its plane through the given point, about which the product of inertia of the lamina vanishes.

Let  $P(x, y)$  be a point of the plane area  $A$  referred to rectangular axes  $OX, OY$ . Let  $(x', y')$  be the coordinates of  $P$  referred to another pair of rectangular axes  $OX', OY'$  in the same plane and inclined at an angle  $\theta$  to the first pair (Fig. 7.43).




Fig. 7.43

$$\begin{aligned} \text{Then} \quad x' &= x \cos \theta + y \sin \theta \\ y' &= y \cos \theta - x \sin \theta \end{aligned}$$

If  $I_x, I_y$  be the moments of inertia of the area  $A$  about  $OX$  and  $OY$  and  $P_{xy}$  be its product of inertia about these axes, then

$$I_x = \iint_A \rho y^2 dA, I_y = \iint_A \rho x^2 dA, P_{xy} = \iint_A \rho xy dA.$$

$\therefore$  the product of inertia  $P'_{xy}$  about  $OX'$  and  $OY'$  is given by

$$\begin{aligned} P'_{xy} &= \iint_A \rho x'y' dA = \iint_A \rho (x \cos \theta + y \sin \theta)(y \cos \theta - x \sin \theta) dA \\ &= \sin \theta \cos \theta \iint_A \rho (y^2 - x^2) dA + (\cos^2 \theta - \sin^2 \theta) \iint_A \rho xy dA \\ &= \frac{1}{2} \sin 2\theta \cdot (I_x - I_y) + \cos 2\theta P_{xy}. \end{aligned}$$

Now  $OX', OY'$  will be the principal axes of the area  $A$  if  $P'_{xy}$  vanishes.

$$\text{i.e., If } \frac{1}{2} \sin 2\theta (I_x - I_y) + \cos 2\theta P_{xy} = 0$$

$$\text{i.e., If } \tan 2\theta = 2P_{xy}/(I_y - I_x).$$

This gives two values of  $\theta$  differing by  $\pi/2$ .

**Example 7.41.** Show that the principal axes at the node of a half-loop of the lemniscate  $r^2 = a^2 \cos 2\theta$  are inclined to the initial line at angles

$$\frac{1}{2} \tan^{-1} \frac{1}{2} \text{ and } \frac{\pi}{2} + \frac{1}{2} \tan^{-1} \frac{1}{2}.$$

**Solution.** Let the element of mass at  $P(r, \theta)$  be  $\rho r d\theta dr$ .

$$\text{Then } I_x = \rho \int_0^{\pi/4} \int_0^{a\sqrt{\cos 2\theta}} r^2 \sin^2 \theta \cdot r d\theta dr$$

[See Fig. 7.40]

$$= \frac{\rho a^4}{4} \int_0^{\pi/4} \sin^2 \theta \cos^2 2\theta d\theta = \frac{\rho a^4}{16} \left( \frac{\pi}{4} - \frac{2}{3} \right)$$

$$I_y = \rho \int_0^{\pi/4} \int_0^{a\sqrt{\cos 2\theta}} r^2 \cos^2 \theta \cdot r d\theta dr = \frac{\rho a^4}{16} \left( \frac{\pi}{4} + \frac{2}{3} \right)$$

and

$$P_{xy} = \rho \int_0^{\pi/4} \int_0^{a\sqrt{\cos 2\theta}} r^2 \sin \theta \cos \theta \cdot r d\theta dr = \frac{\rho a^4}{48}.$$

Hence the required direction of the principal axes at  $O$  are given by

$$\tan 2\theta = \frac{2P_{xy}}{I_y - I_x} = \frac{\rho a^4 / 24}{(\rho a^4 / 16) \times (4/3)} = \frac{1}{2}$$

$$\text{or by } \theta = \frac{1}{2} \tan^{-1} \frac{1}{2} \text{ and } \frac{\pi}{2} + \frac{1}{2} \tan^{-1} \frac{1}{2}.$$

### PROBLEMS 7.7

- Using double integrals, find the moment of inertia about the  $x$ -axis of the area enclosed by the lines  $x = 0, y = 0, (x/a) + (y/b) = 1$ . (P.T.U., 2005)
- Find the moment of inertia of a circular plate about a tangent.
- Find the moment of inertia of the area  $y = \sin x$  from  $x = 0$  to  $x = 2\pi$  about  $OX$ .

4. Find the moment of inertia of a quadrant of the ellipse  $(x/a)^2 + (y/b)^2 = 1$  of mass  $M$  about the  $x$ -axis, if the density at a point is proportional to  $xy$ .
5. Find the moment of inertia about the initial line of the cardioid  $r = a(1 + \cos \theta)$ .
6. Find the moment of inertia of a uniform spherical ball of mass  $M$  and radius  $R$  about a diameter.
7. Find the moment of inertia of a solid right circular cylinder about (i) its axis (P.T.U., 2006)  
(ii) a diameter of the base.
8. Find the M.I. of a solid right circular cone having base-radius  $r$  and height  $h$ , about (i) its axis, (ii) an axis through the vertex and perpendicular to its axis, (iii) a diameter of its base.
9. Find the moment of inertia of a hollow sphere about a diameter, its external and internal radii being 51 metres and 49 metres.
10. Find the moment of inertia about  $z$ -axis of a homogeneous tetrahedron bounded by the planes  $x = 0, y = 0, z = x + y$  and  $z = 1$ .
11. Find the moment of inertia of an octant of the ellipsoid  $(x/a)^2 + (y/b)^2 + (z/c)^2 = 1$ , about the  $x$ -axis.
12. Find the product of inertia of a quadrant of the ellipse  $(x/a)^2 + (y/b)^2 = 1$ , about the coordinate axes.
13. Show that the principal axes at the origin of the triangle enclosed by  $x = 0, y = 0, (x/a) + (y/b) = 1$  are inclined to the  $x$ -axis at angles  $\alpha$  and  $\alpha + \pi/2$ , where  $\alpha = \frac{1}{2} \tan^{-1} [ab/(a^2 - b^2)]$  (U.P.T.U., 2002)
14. The lengths  $AB$  and  $AD$  of the sides of a rectangle  $ABCD$  are  $2a$  and  $2b$ . Show that the inclination to  $AB$  of one of the principal axes at  $A$  is  $\frac{1}{2} \tan^{-1} \left\{ \frac{3ab}{2(a^2 - b^2)} \right\}$ .

## 7.14 BETA FUNCTION

The beta function is defined as

$$\beta(m, n) = \int_0^1 x^{m-1} (1-x)^{n-1} dx \quad \begin{cases} m > 0 \\ n > 0 \end{cases} \quad \dots(1)$$

Putting  $x = 1 - y$  in (1), we get  $\beta(m, n) = - \int_1^0 (1-y)^{m-1} y^{n-1} dy$

$$= \int_0^1 y^{n-1} (1-y)^{m-1} dy = \beta(n, m)$$

Thus  $\beta(m, n) = \beta(n, m)$  ...(2)

Putting  $x = \sin^2 \theta$  so that  $dx = 2 \sin \theta \cos \theta d\theta$ , (1) becomes

$$\begin{aligned} \beta(m, n) &= \int_0^{\pi/2} (\sin^2 \theta)^{m-1} (\cos^2 \theta)^{n-1} 2 \sin \theta \cos \theta d\theta \\ &= 2 \int_0^{\pi/2} \sin^{2m-1} \theta \cos^{2n-1} \theta d\theta \end{aligned} \quad \dots(3)$$

which is another form of  $\beta(m, n)$ .

This function is also *Euler's integral of the first kind*.\*

## 7.15 (1) GAMMA FUNCTION

The gamma function is defined as

$$\Gamma(n) = \int_0^\infty e^{-x} x^{n-1} dx \quad (n > 0) \quad \dots(i)$$

This integral is also known as *Euler's integral of the second kinds*. It defines a function of  $n$  for positive values of  $n$ .

\*After an enormously creative Swiss mathematician *Leonhard Euler (1707-1783)*. He studied under *John Bernoulli* and became a professor of mathematics in St. Petersburg, Russia. Even after becoming totally blind in 1771, he contributed to almost all branches of mathematics.



In particular,  $\Gamma(1) = \int_0^{\infty} e^{-x} dx = \left| -e^{-x} \right|_0^{\infty} = 1.$  ... (ii)

**(2) Reduction formula for  $\Gamma(n)$ .**

Since  $\Gamma(n+1) = \int_0^{\infty} e^{-x} x^n dx$  [Integrating by parts] =  $\left| -x^n e^{-x} \right|_0^{\infty} + n \int_0^{\infty} e^{-x} x^{n-1} dx$

$\therefore \Gamma(n+1) = n\Gamma(n)$  ... (iii)

which is the reduction formula for  $\Gamma(n)$ . From this formula, it is clear that if  $\Gamma(n)$  is known throughout a unit interval say :  $1 < n \leq 2$ , then the values of  $\Gamma(n)$  throughout the next unit interval  $2 < n \leq 3$  are found, from which the values of  $\Gamma(n)$  for  $3 < n \leq 4$  are determined and so on. In this way, the values of  $\Gamma(n)$  for all positive values of  $n > 1$  may be found by successive application of (iii).

Also using (iii) in the form

$$\Gamma(n) = \frac{\Gamma(n+1)}{n} \quad \dots (iv)$$

We can define  $\Gamma(n)$  for values of  $n$  for which the definition (1) fails. It gives the value of  $\Gamma(n)$  for  $0 < n \leq 1$  in terms of the values of  $\Gamma(n)$  for  $1 < n \leq 2$ . Thus we can define  $\Gamma(n)$  for all  $n < 0$  provided its value for  $1 < n \leq 2$  is known. Also if  $-1 < n < 0$ , (4) gives  $\Gamma(n)$  in terms of its values for  $0 < n < 1$ . Then we may find,  $\Gamma(n)$  for  $-2 < n < -1$  and so on.

Thus (i) and (iv) together give a complete definition of  $\Gamma(n)$  for all values of  $n$  except when  $n$  is zero or a negative integer and its graph is as shown in Fig. 7.44. The values of  $\Gamma(n)$  for  $1 < n \leq 2$  are given in (Table I-Appendix 2) from which the values of  $\Gamma(n)$  for values of  $n$  outside the interval  $1 < n \leq 2$  ( $n \neq 0, -1, -2, -3, \dots$ ) may be found.




Fig. 7.44

**(3) Value of  $\Gamma(n)$  in terms of factorial.**

Using  $\Gamma(n+1) = n\Gamma(n)$  successively, we get

$$\Gamma(2) = 1 \times \Gamma(1) = 1 !$$

$$\Gamma(3) = 2 \times \Gamma(2) = 2 \times 1 = 2 !$$

$$\Gamma(4) = 3 \times \Gamma(3) = 3 \times 2 ! = 3 !$$

.....

In general  $\Gamma(n+1) = n !$  provided  $n$  is a positive integer

... (v)

Taking  $n = 0$ , it defines  $0 ! = \Gamma(1) = 1$ .

**(4) Value of  $\Gamma(\frac{1}{2})$ . We have**

$$\Gamma\left(\frac{1}{2}\right) = \int_0^{\infty} e^{-x} x^{-1/2} dx$$

[Put  $x = y^2$  so that  $dx = 2y dy$ ]

$$= 2 \int_0^{\infty} e^{-y^2} dy \text{ which is also } = 2 \int_0^{\infty} e^{-x^2} dx$$

$$\therefore \left[ \Gamma\left(\frac{1}{2}\right) \right]^2 = 4 \int_0^{\infty} \int_0^{\infty} e^{-(x^2+y^2)} dx dy = 4 \int_0^{\pi/2} \int_0^{\infty} e^{-r^2} r dr d\theta$$

$$= 4 \cdot \frac{\pi}{2} \int_0^{\infty} e^{-r^2} r dr = 2\pi \left[ \left( -\frac{1}{2} \right) e^{-r^2} \right]_0^{\infty} = \pi$$

whence  $\Gamma\left(\frac{1}{2}\right) = \sqrt{\pi} = 1.772$

... (vi) (V.T.U., 2006)

**7.16 RELATION BETWEEN BETA AND GAMMA FUNCTIONS**

$$\beta(m, n) = \frac{\Gamma(m) \Gamma(n)}{\Gamma(m+n)}$$

We have

$$\Gamma(m) = \int_0^{\infty} e^{-t} t^{m-1} dt$$

[Put  $t = x^2$  so that  $dt = 2x dx$ ]

$$= 2 \int_0^{\infty} e^{-x^2} x^{2m-1} dx \quad \dots(2)$$

Similarly,  $\Gamma(n) = 2 \int_0^{\infty} e^{-y^2} y^{2n-1} dy$

$$\therefore \Gamma(m)\Gamma(n) = 4 \int_0^{\infty} e^{-x^2} x^{2m-1} dx \int_0^{\infty} e^{-y^2} y^{2n-1} dy$$

$$= 4 \int_0^{\infty} \int_0^{\infty} e^{-(x^2+y^2)} x^{2m-1} y^{2n-1} dx dy \quad \dots(3) \quad [\because \text{the limits of integration are constant.}]$$

Now change to polar coordinates by writing  $x = r \cos \theta$ ,  $y = r \sin \theta$  and  $dx dy = r d\theta dr$ . To cover the region in (3) which is the entire first quadrant,  $r$  varies from 0 to  $\infty$  and  $\theta$  from 0 to  $\pi/2$ . Thus (3) becomes

$$\Gamma(m)\Gamma(n) = 4 \int_0^{\pi/2} \int_0^{\infty} e^{-r^2} r^{2(m+n)-1} \cos^{2m-1} \theta \sin^{2n-1} \theta d\theta dr$$

$$= \left[ 2 \int_0^{\pi/2} \cos^{2m-1} \theta \sin^{2n-1} \theta d\theta \right] \times \left[ 2 \int_0^{\infty} e^{-r^2} r^{2(m+n)-1} dr \right] \quad \dots(4)$$

But by (2),  $2 \int_0^{\infty} e^{-r^2} r^{2(m+n)-1} dr = \Gamma(m+n)$

and by (3) of § 7.14,  $2 \int_0^{\pi/2} \cos^{2m-1} \theta \sin^{2n-1} \theta d\theta = \beta(m, n)$ .

Thus (4) gives  $\Gamma(m)\Gamma(n) = \beta(m, n) \Gamma(m+n)$  (U.T.U., 2010 ; Bhopal, 2009 ; V.T.U., 2008 S)  
whence follows (1) which is extremely useful for evaluating definite integrals in terms of gamma functions.

**Cor.** Rule to evaluate  $\int_0^{\pi/2} \sin^p x \cos^q x dx$ .

$$\int_0^{\pi/2} \sin^p x \cos^q x dx = \frac{1}{2} \beta\left(\frac{p+1}{2}, \frac{q+1}{2}\right) \quad [\text{By (3) of § 7.14}]$$

$$= \frac{\Gamma\left(\frac{p+1}{2}\right) \Gamma\left(\frac{q+1}{2}\right)}{2\Gamma\left(\frac{p+q+2}{2}\right)} \quad \dots(5)$$

In particular, when  $q = 0$ , and  $p = n$ , we have

$$\int_0^{\pi/2} \sin^n x dx = \frac{\Gamma\left(\frac{n+1}{2}\right)}{\Gamma\left(\frac{n+2}{2}\right)} \cdot \frac{\sqrt{\pi}}{2}$$

Similarly,

$$\int_0^{\pi/2} \cos^n x dx = \frac{\Gamma\left(\frac{n+1}{2}\right)}{\Gamma\left(\frac{n+2}{2}\right)} \cdot \frac{\sqrt{\pi}}{2} \quad \dots(6)$$

**Example 7.42.** Show that

(a)  $\Gamma(n) = \int_0^1 \left( \log \frac{1}{y} \right)^{n-1} dy \quad (n > 0).$  (J.N.T.U., 2003 ; Madras, 2003 S)

(b)  $\beta(p, q) = \int_0^{\infty} \frac{y^{q-1}}{(1+y)^{p+q}} dy$  (V.T.U., 2003 ; Gauhati, 1999)

$$= \int_0^1 \frac{x^{p-1} + x^{q-1}}{(1+x)^{p+q}} dx. \quad (V.T.U., 2008 ; Osmania, 2003 ; Rohtak, 2003)$$

**Solution.** (a)  $\Gamma(n) = \int_0^{\infty} x^{n-1} e^{-x} dx \quad (n > 0)$



$$= \int_1^0 \left( \log \frac{1}{y} \right)^{n-1} y \left( -\frac{1}{y} dy \right) = \int_0^1 \left( \log \frac{1}{y} \right)^{n-1} dy.$$

$$\left| \begin{array}{l} \text{Put } y = e^{-x} \\ \text{i.e., } x = \log(1/y) \\ \text{so that } dx = -(1/y) dy \end{array} \right.$$

$$(b) \quad \beta(p, q) = \int_0^1 x^{p-1} (1-x)^{q-1} dx$$

$$= \int_{\infty}^0 \frac{1}{(1+y)^{p-1}} \left( \frac{y}{1+y} \right)^{q-1} \frac{-1}{(1+y)^2} dy$$

$$\left| \begin{array}{l} \text{Put } x = \frac{1}{1+y} \text{ i.e., } y = \frac{1}{x} - 1 \\ \text{so that } dx = \frac{-1}{(1+y)^2} dy \end{array} \right.$$

$$= \int_0^{\infty} \frac{y^{q-1}}{(1+y)^{p+q}} dy = \int_0^1 \frac{y^{q-1}}{(1+y)^{p+q}} dy + \int_1^{\infty} \frac{y^{q-1}}{(1+y)^{p+q}} dy$$

Now substituting  $y = 1/z$  in the second integral, we get

$$\int_1^{\infty} \frac{y^{q-1}}{(1+y)^{p+q}} dy = \int_1^0 \frac{1}{z^{q-1}} \cdot \frac{1}{(1+1/z)^{p+q}} \left( -\frac{1}{z^2} \right) dz = \int_0^1 \frac{z^{p-1}}{(1+z)^{p+q}} dz.$$

$$\text{Hence, } \beta(p, q) = \int_0^1 \frac{y^{q-1}}{(1+y)^{p+q}} dy + \int_0^1 \frac{z^{p-1}}{(1+z)^{p+q}} dz = \int_0^1 \frac{x^{p-1} + x^{q-1}}{(1+x)^{p+q}} dx.$$

**Example 7.43.** Express the following integrals in terms of gamma functions :

$$(a) \int_0^1 \frac{dx}{\sqrt{1-x^4}}$$

$$(b) \int_0^{\pi/2} \sqrt{\tan \theta} d\theta.$$

(Madras, 2006)

$$(c) \int_0^{\infty} \frac{x^c}{e^x} dx \quad (\text{U.P.T.U., 2006})$$

$$(d) \int_0^{\infty} a^{-bx^2} dx.$$

$$(e) \int_0^1 x^5 [\log(1/x)]^3 dx$$

(Madras, 2000)

**Solution.** (a)  $\int_0^1 \frac{dx}{\sqrt{1-x^4}}$

$$\left| \begin{array}{l} \text{Put } x^2 = \sin \theta, \text{ i.e., } x = \sin^{1/2} \theta \\ \text{so that } dx = 1/2 \sin^{-1/2} \theta \cos \theta d\theta \end{array} \right.$$

$$= \int_0^{\pi/2} \frac{1}{2} \cdot \frac{\sin^{-1/2} \theta \cdot \cos \theta d\theta}{\sqrt{1-\sin^2 \theta}} = \frac{1}{2} \int_0^{\pi/2} \sin^{-1/2} \theta d\theta = \frac{1}{2} \cdot \frac{\sqrt{\pi}}{2} \cdot \frac{\Gamma\left(-\frac{1}{2}+1\right)}{\Gamma\left(-\frac{1}{2}+2\right)} = \frac{\sqrt{\pi}}{4} \frac{\Gamma\left(\frac{1}{4}\right)}{\Gamma\left(\frac{3}{4}\right)}$$

$$(b) \int_0^{\pi/2} \sqrt{\tan \theta} d\theta = \int_0^{\pi/2} \sin^{1/2} \theta \cos^{-1/2} \theta d\theta$$

$$= \frac{\Gamma\left(\frac{1}{2}+1\right) \Gamma\left(-\frac{1}{2}+1\right)}{2\Gamma\left(\frac{1}{2}-\frac{1}{2}+2\right)} = \frac{\Gamma\left(\frac{3}{4}\right) \Gamma\left(\frac{1}{4}\right)}{2\Gamma(1)} = \frac{1}{2} \Gamma\left(\frac{1}{4}\right) \Gamma\left(\frac{3}{4}\right)$$

$$(c) \int_0^{\infty} \frac{x^c}{e^x} dx = \int_0^{\infty} \frac{x^c}{e^{x \log e}} dx$$

$$[\because c^x = e^{\log c^x} = e^{x \log c}]$$

$$= \int_0^{\infty} e^{-x \log c} x^c dx$$

$$[\text{Put } x \log c = t \text{ so that } dx = dt/\log c]$$

$$= \int_0^{\infty} e^{-t} \left( \frac{t}{\log c} \right)^c \frac{dt}{\log c} = \frac{1}{(\log c)^{c+1}} \int_0^{\infty} t^c e^{-t} dt = \Gamma(c+1)/(\log c)^{c+1}$$

$$(d) \int_0^{\infty} a^{-bx^2} dx = \int_0^{\infty} e^{-bx^2 \log a} dx \quad \left[ \begin{array}{l} \text{Put } (b \log a) x^2 = t \\ \text{so that } dx = dt/2\sqrt{(b \log a)} \end{array} \right]$$

$$= \frac{1}{2\sqrt{(b \log a)}} \int_0^{\infty} e^{-t} t^{-1/2} dt = \frac{\Gamma\left(\frac{1}{2}\right)}{2\sqrt{(b \log a)}} = \frac{\sqrt{\pi}}{2\sqrt{(b \log a)}}$$

$$(e) \int_0^1 x^4 [\log(1/x)]^3 dx = \frac{1}{625} \int_0^{\infty} e^{-t} \cdot t^3 dt \quad \left[ \begin{array}{l} \text{Put } x = e^{-t/5} \text{ so that } \log(1/x) = t/5 \\ dx = -\frac{1}{5} e^{-t/5} dt \end{array} \right]$$

$$= \frac{\Gamma(4)}{625} = \frac{6}{625}.$$

**Example 7.44.** Evaluate  $\int_0^{\infty} e^{-ax} x^{m-1} \sin bx dx$  in terms of Gamma function. (U.P.T.U., 2003)

**Solution.** We have  $\Gamma(m) = \int_0^{\infty} e^{-x} x^{m-1} dx$  [Put  $x = ay$ ,  $dx = ady$ ]

$$= \int_0^{\infty} e^{-ay} a^m y^{m-1} dy \quad \text{or} \quad \int_0^{\infty} e^{-ay} y^{m-1} dy = \Gamma(m)/a^m. \quad \dots(i)$$

Then

$$I = \int_0^{\infty} e^{-ax} x^{m-1} \sin bx dx = \int_0^{\infty} e^{-ax} x^{m-1} (\text{Imaginary part of } e^{ibx}) dx$$

$$= \text{I.P. of } \int_0^{\infty} e^{-(a-ib)x} x^{m-1} dx$$

$$= \text{I.P. of } \{\Gamma(m)/(a-ib)^m\} \quad [\text{By (i)}]$$

$$= \text{I.P. of } \{\Gamma(m)/(r^m (\cos \theta - i \sin \theta)^m)\} \quad \text{where } a = r \cos \theta, b = r \sin \theta$$

$$= \text{I.P. of } \Gamma(m)/(r^m (\cos m\theta - i \sin m\theta)) \quad (\text{Using Demoivre's theorem §19.5})$$

$$= \text{I.P. of } \left\{ \frac{\Gamma(m) \cdot (\cos m\theta + i \sin m\theta)}{r^m (\cos m\theta + i \sin m\theta) (\cos m\theta - i \sin m\theta)} \right\}$$

$$= \frac{\Gamma(m)}{r^m} \sin m\theta \quad \text{where } r = \sqrt{(a^2 + b^2)}, \theta = \tan^{-1} b/a.$$

**Example 7.45.** Prove that  $\int_0^1 \frac{x^2 dx}{\sqrt{(1-x^4)}} \times \int_0^1 \frac{dx}{\sqrt{(1+x^4)}} = \frac{\pi}{4\sqrt{2}}$ .

$$\text{Solution. } \int_0^1 \frac{x^2 dx}{\sqrt{(1-x^4)}} \times \int_0^{\pi/2} \frac{\sin \theta}{\cos \theta} \cdot \frac{\cos \theta}{2\sqrt{(\sin \theta)}} d\theta \quad \left[ \text{Putting } x^2 = \sin \theta, dx = \frac{\cos \theta d\theta}{2\sqrt{(\sin \theta)}} \right]$$

$$= \frac{1}{2} \int_0^{\pi/2} \sqrt{(\sin \theta)} d\theta = \frac{1}{4} \beta\left(\frac{3}{4}, \frac{1}{2}\right) = \frac{1}{4} \frac{\Gamma(3/4) \Gamma(1/2)}{\Gamma(5/4)} = \frac{\Gamma(3/4) \Gamma(1/2)}{\Gamma(1/4)}$$

$$\int_0^1 \frac{dx}{\sqrt{(1+x^4)}} = \int_0^{\pi/4} \frac{\sec^2 \theta d\theta}{2\sqrt{(\tan \theta) \sec \theta}} \quad \left[ \text{Putting } x^2 = \tan \theta, dx = \frac{\sec^2 \theta d\theta}{2\sqrt{(\tan \theta)}} \right]$$

$$= \frac{1}{\sqrt{2}} \int_0^{\pi/4} \frac{d\theta}{\sqrt{(\sin 2\theta)}} = \frac{1}{2\sqrt{2}} \int_0^{\pi/2} \sin^{-1/2} \phi d\phi \quad \left[ \text{Putting } 2\theta = \phi, d\theta = \frac{1}{2} d\phi \right]$$

$$= \frac{1}{4\sqrt{2}} \beta\left(\frac{1}{4}, \frac{1}{2}\right) = \frac{1}{4\sqrt{2}} \frac{\Gamma(1/4) \Gamma(1/2)}{\Gamma(3/4)}$$



$$\therefore \int_0^1 \frac{x^2 dx}{\sqrt{(1-x^4)}} \times \int_0^1 \frac{dx}{\sqrt{(1+x^4)}} = \frac{1}{4\sqrt{2}} \left\{ \Gamma\left(\frac{1}{2}\right) \right\}^2 = \frac{\pi}{4\sqrt{2}}.$$

**Example 7.46.** Prove that (i)  $\beta(m, 1/2) = 2^{2m-1} \beta(m, m)$  (V.T.U., 2004)

$$(ii) \Gamma(m) \Gamma\left(m + \frac{1}{2}\right) = \frac{\sqrt{\pi}}{2^{2m-1}} \Gamma(2m) \quad (\text{Duplication Formula})$$

(V.T.U., 2010 ; Kerala, M.E., 2005 ; Madras, 2003 S)

**Solution.** (i) We know that  $\beta(m, n) = 2 \int_0^{\pi/2} \sin^{2m-1} \theta \cos^{2n-1} \theta d\theta$  ... (1)

Putting  $n = \frac{1}{2}$ , we have  $\beta\left(m, \frac{1}{2}\right) = 2 \int_0^{\pi/2} \sin^{2m-1} \theta d\theta$  ... (2)

Again putting  $n = m$  in (i), we get  $\beta(m, m) = 2 \int_0^{\pi/2} (\sin \theta \cos \theta)^{2m-1} d\theta$

$$\begin{aligned} &= \frac{1}{2^{2m-2}} \int_0^{\pi/2} \sin^{2m-1} 2\theta d\theta \\ &= \frac{1}{2^{2m-1}} \int_0^{\pi} \sin^{2m-1} \phi d\phi, \text{ putting } 2\theta = \phi \\ &= \frac{1}{2^{2m-1}} \cdot 2 \int_0^{\pi/2} \sin^{2m-1} \phi d\phi \end{aligned}$$

or  $2^{2m-1} \beta(m, m) = 2 \int_0^{\pi/2} \sin^{2m-1} \theta d\theta = \beta\left(m, \frac{1}{2}\right)$  [by (2)]

(ii) Rewriting the above result in terms of  $\Gamma$  functions, we get

$$2^{2m-1} \frac{\Gamma(m) \Gamma(m)}{\Gamma(m+m)} = \frac{\Gamma(m) \Gamma\left(\frac{1}{2}\right)}{\Gamma\left(m + \frac{1}{2}\right)} \quad \left[ \because \Gamma\left(\frac{1}{2}\right) = \sqrt{\pi} \right]$$

or  $\Gamma(m) \Gamma\left(m + \frac{1}{2}\right) = \frac{\sqrt{\pi} \Gamma(2m)}{2^{2m-1}}.$

**Example 7.47.** Prove that

(a)  $\iint_D x^{l-1} y^{m-1} dx dy = \frac{\Gamma(l) \Gamma(m)}{\Gamma(l+m+1)} h^{l+m}$  where  $D$  is the domain  $x \geq 0, y \geq 0$  and  $x+y \leq h$ .

(U.P.T.U., 2005)

(b)  $\iiint_V x^{l-1} y^{m-1} z^{n-1} dx dy dz = \frac{\Gamma(l) \Gamma(m) \Gamma(n)}{\Gamma(l+m+n+1)}$

where  $V$  is the region  $x \geq 0, y \geq 0, z \geq 0$  and  $x+y+z \leq 1$ . This important result is known as **Dirichlet's integral\***.

**Solution.** (a) Putting  $x/h = X$  and  $y/h = Y$ , we see that the given integral

$$\begin{aligned} &= \iint_{D'} (hX)^{l-1} (hY)^{m-1} h^2 dXdY \text{ where } D' \text{ is the domain } X \geq 0, Y \geq 0 \text{ and } X+Y \leq 1. \\ &= h^{l+m} \int_0^1 \int_0^{1-X} X^{l-1} Y^{m-1} dY dX = h^{l+m} \int_0^1 X^{l-1} \left[ \frac{Y^m}{m} \right]_0^{1-X} dX \\ &= \frac{h^{l+m}}{m} \int_0^1 X^{l-1} (1-X)^m dX = \frac{h^{l+m}}{m} \beta(l, m+1) = \frac{h^{l+m}}{m} \cdot \frac{\Gamma(l) \Gamma(m+1)}{\Gamma(l+m+1)} \end{aligned}$$

\*Named after a German mathematician *Peter Gustav Lejeune Dirichlet* (1805–1859) who studied under Cauchy and succeeded Gauss at Gottingen. He is known for his contributions to Fourier series and number theory.

$$= h^{l+m} \frac{\Gamma(l) \Gamma(m)}{\Gamma(l+m+1)} \quad \dots(i) [\because \Gamma(m+1)/m = \Gamma(m)]$$

(b) Taking  $y + z \leq 1 - x (= h)$ : say), the triple integral

$$\begin{aligned} &= \int_0^1 \int_0^{1-x} \int_0^{1-x-y} x^{l-1} y^{m-1} z^{n-1} dz dy dx \\ &= \int_0^1 x^{l-1} \left[ \int_0^h \int_0^{h-y} y^{m-1} z^{n-1} dz dy \right] dx = \int_0^1 x^{l-1} \frac{\Gamma(m) \Gamma(n)}{\Gamma(m+n+1)} h^{m+n} dx \quad \dots [\text{By (i)}] \\ &= \frac{\Gamma(m) \Gamma(n)}{\Gamma(m+n+1)} \int_0^1 x^{l-1} (1-x)^{m+n} dx = \frac{\Gamma(m) \Gamma(n)}{\Gamma(m+n+1)} \beta(l, m+n+1) \\ &= \frac{\Gamma(m) \Gamma(n)}{\Gamma(m+n+1)} \cdot \frac{\Gamma(l) \Gamma(m+n+1)}{\Gamma(l+m+n+1)} = \frac{\Gamma(l) \Gamma(m) \Gamma(n)}{\Gamma(l+m+n+1)}. \end{aligned}$$

**Example 7.48.** Evaluate the integral  $\iiint x^{l-1} y^{m-1} z^{n-1} dx dy dz$  where  $x, y, z$  are all positive with condition,  $(x/a)^p + (y/b)^q + (z/c)^r \leq 1$ . (U.P.T.U., 2005 S)

**Solution.** Put  $(x/a)^p = u$ , i.e.,  $x = au^{1/p}$  so that  $dx = \frac{a}{p} u^{1/p-1} du$

$(y/b)^q = v$ , i.e.,  $y = bv^{1/q}$  so that  $dy = \frac{b}{q} v^{1/q-1} dv$

and  $(z/c)^r = w$ , i.e.,  $z = cw^{1/r}$  so that  $dz = \frac{c}{r} w^{1/r-1} dw$

$$\begin{aligned} \text{Then } &\iiint x^{l-1} y^{m-1} z^{n-1} dx dy dz \\ &= \iiint (au^{1/p})^{l-1} (bv^{1/q})^{m-1} (cw^{1/r})^{n-1} \left(\frac{a}{p}\right) u^{1/p-1} \left(\frac{b}{q}\right) v^{1/q-1} \left(\frac{c}{r}\right) w^{1/r-1} du dv dw \\ &= \frac{a^l b^m c^n}{pqr} \iiint u^{l/p-1} v^{m/q-1} w^{n/r-1} du dv dw \text{ where } u + v + w \leq 1. \\ &= \frac{a^l b^m c^n}{pqr} \frac{\Gamma(l/p) \Gamma(m/q) \Gamma(n/r)}{\Gamma(l/p + m/q + n/r + 1)} \quad [\text{By Dirichlet's integral}] \end{aligned}$$

**Example 7.49.** The plane  $\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$  meets the axes in A, B and C. Apply Dirichlet's integral to find the volume of the tetrahedron OABC. Also find its mass if the density at any point is  $kxyz$ . (U.P.T.U., 2004)

**Solution.** Put  $x/a = u, y/b = v, z/c = w$  then the tetrahedron OABC has  $u \geq 0, v \geq 0, w \geq 0$  and  $u + v + w \leq 1$ .

$$\begin{aligned} \therefore \text{ volume of this tetrahedron} &= \iiint_D dx dy dz \\ &= \iiint_D abc du dv dw \quad \left[ \begin{array}{l} a dx = a du, dy = b dv, dz = c dw \\ \text{for } D' = u \geq 0, v \geq 0, w \geq 0 \text{ \& } u + v + w \leq 1. \end{array} \right. \\ &= abc \iiint_D u^{1-1} v^{1-1} w^{1-1} du dv dw \\ &= abc \frac{\Gamma(1) \Gamma(1) \Gamma(1)}{\Gamma(1+1+1+1)} = \frac{abc}{6} \quad [\text{By Dirichlet's integral}] \end{aligned}$$

$$\begin{aligned} \text{Mass} &= \iiint kxyz dx dy dz = \iiint k(au)(bv)(cw) abc du dv dw \\ &= ka^2 b^2 c^2 \iiint u^{2-1} v^{2-1} w^{2-1} du dv dw \\ &= ka^2 b^2 c^2 \frac{\Gamma(2) \Gamma(2) \Gamma(2)}{\Gamma(2+2+2+1)} ka^2 b^2 c^2 \cdot \frac{1}{6!} = \frac{k}{720} a^2 b^2 c^2. \end{aligned}$$



PROBLEMS 7.8

1. Compute :

(i)  $\Gamma(3.5)$  (Assam, 1998)

(ii)  $\Gamma(4.5)$

(iii)  $\Gamma\left(\frac{1}{4}\right) \Gamma\left(\frac{3}{4}\right)$  (S.V.T.U., 2009) (iv)  $\beta(2.5, 1.5)$  (v)  $\beta\left(\frac{9}{2}, \frac{7}{2}\right)$ . (Andhra, 2000)

2. Express the following integrals in terms of gamma functions :

(i)  $\int_0^\infty e^{-x^2} dx$  (ii)  $\int_0^\infty x^{p-1} e^{-kx} dx (k > 0)$  (Delhi, 2002 ; V.T.U., 2000)

(iii)  $\int_0^\infty \sqrt{x} e^{-x^3} dx$  (J.N.T.U., 2003) (iv)  $\int_0^\infty \frac{dx}{x^{p+1} (x-1)^q} (-p < q < 1)$

3. Show that :

(i)  $\int_0^\infty \frac{x^4}{4^x} dx = \frac{\Gamma(5)}{(\log 4)^5}$  (Marathwada, 2008)

(ii)  $\int_0^{\pi/2} \sqrt{\cot \theta} d\theta = \frac{1}{2} \Gamma\left(\frac{1}{4}\right) \Gamma\left(\frac{3}{4}\right)$  (Osmania, 2003 S ; V.T.U., 2001)

(iii)  $\int_0^{\pi/2} [\sqrt{\tan \theta} + \sqrt{\sec \theta}] d\theta = \frac{1}{2} \Gamma\left(\frac{1}{4}\right) \left\{ \Gamma\left(\frac{3}{4}\right) + \sqrt{\pi} \Gamma\left(\frac{3}{4}\right) \right\}$  (J.N.T.U., 2000)

(iv)  $\int_0^{\pi/2} \sqrt{\sin \theta} d\theta \times \int_0^{\pi/2} \frac{d\theta}{\sqrt{\sin \theta}} = \pi$ . (V.T.U., 2007)

4. Given  $\int_0^\infty \frac{x^{n-1}}{1+x} dx = \frac{\pi}{\sin n\pi}$ , show that  $\Gamma(n) \Gamma(1-n) = \frac{\pi}{\sin n\pi}$ . (S.V.T.U., 2008)

Hence evaluate  $\int_0^\infty \frac{dy}{1+y^4}$ . (V.T.U., 2006 ; J.N.T.U., 2005)

5. Prove that :

(i)  $\int_0^1 \frac{xdx}{\sqrt{(1-x^5)}} = \frac{1}{5} \beta\left(\frac{2}{5}, \frac{1}{2}\right)$  (Raipur, 2006) (ii)  $\int_0^1 \frac{dx}{\sqrt{(1+x^4)}} = \frac{1}{4\sqrt{2}} \beta\left(\frac{1}{4}, \frac{1}{2}\right)$ . (V.T.U., 2003)

(iii)  $\int_0^1 x^3 (1-\sqrt{x})^5 dx = 2\beta(8, 6)$ . (Marathwada, 2008 ; J.N.T.U., 2006)

6. Show that (i)  $\int_a^b (x-a)^{m-1} (b-x)^{n-1} dx = (b-a)^{m+n-1} \beta(m, n)$  (P.T.U., 2010 ; Mumbai, 2005)

(ii)  $\int_0^\infty \frac{x^{m-1}}{(a+bx)^{m+n}} dx = \frac{1}{a^n b^m} \beta(m, n)$  (Nagpur, 2009) (iii)  $\int_0^\infty \frac{x^{10}-x^{18}}{(1+x)^{30}} dx = 0$  (Mumbai, 2005)

(iv)  $\int_0^1 \frac{(1-x^4)^{3/4}}{(1+x^4)^2} dx = \frac{1}{2^{9/2}} \beta\left(\frac{7}{4}, \frac{1}{4}\right)$  (Mumbai, 2007)

7. Prove that  $\int_0^1 x^m (\log x)^n dx = \frac{(-1)^n n!}{(m+1)^{n+1}}$ , where  $n$  is a positive integer and  $m > -1$ . (S.V.T.U., 2006)

Hence evaluate  $\int_0^1 x (\log x)^3 dx$ . (Nagpur, 2009)

8. Show that  $\int_0^1 y^{q-1} \left(\log \frac{1}{y}\right)^{p-1} dy = \frac{\Gamma(p)}{q^p}$ , where  $p > 0, q > 0$ . (Rohtak, 2006 S)

9. Express  $\int_0^1 x^m (1-x^n)^p dx$  in terms of gamma functions (Marathwada, 2008)

Hence evaluate : (i)  $\int_0^1 x(1-x^3)^{10} dx$ . (Bhopal, 2008) (ii)  $\int_0^1 \frac{dx}{\sqrt{(1-x^n)}}$  (Anna, 2005)



10. Prove that  $\int_0^\infty \frac{dx}{(e^x + e^{-x})^n} = \frac{1}{4} \beta\left(\frac{n}{2}, \frac{n}{2}\right)$  and hence evaluate  $\int_0^\infty \operatorname{sech}^8 x \, dx$ .

11. Prove that  $\beta\left(n + \frac{1}{2}, n + \frac{1}{2}\right) = \frac{\Gamma(n + 1/2)\sqrt{\pi}}{2^{2n} \Gamma(n + 1)}$ . Hence show that  $2^n \Gamma(n + 1/2) = 1 \cdot 3 \cdot 5 \dots (2n - 1) \sqrt{\pi}$

(Mumbai, 2007)

12. Prove that :

$$(i) \frac{\beta(m+1, n)}{m} = \frac{\beta(m, n+1)}{n} = \frac{\beta(m, n)}{m+n}$$

$$(ii) \beta(n, n) = \frac{\sqrt{\pi} \Gamma(n)}{2^{2n-1} \Gamma\left(n + \frac{1}{2}\right)}$$

$$(iii) \Gamma\left(n + \frac{1}{2}\right) = \frac{\Gamma(2n+1) \sqrt{\pi}}{2^{2n} \Gamma(n+1)}$$

$$(iv) \beta(m+1) + \beta(m, n+1) = \beta(m, n)$$

(Bhopal, 2008 ; J.N.T.U., 2006 ; Madras, 2003)

13. Show that  $\iint x^{m-1} y^{n-1} dx dy$  over the positive quadrant of the ellipse

$$\left(\frac{x}{a}\right)^2 + \left(\frac{y}{b}\right)^2 = 1 \text{ is } \frac{a^m b^n}{2n} \beta\left(\frac{m}{2}, \frac{n}{2} + 1\right).$$

14. Show that the area in the first quadrant enclosed by the curve  $(x/a)^\alpha + (y/b)^\beta = 1$ ,  $\alpha > 0$ ,  $\beta > 0$ , is given by

$$\frac{ab}{\alpha + \beta} \frac{\Gamma(1/\alpha) \Gamma(1/\beta)}{\Gamma(1/\alpha + 1/\beta)}.$$

15. Find the mass of an octant of the ellipsoid  $(x/a)^2 + (y/b)^2 + (z/c)^2 = 1$ , the density at any point being  $\rho = kxyz$ .

(U.P.T.U., 2002)

## 7.17 (1) ELLIPTIC INTEGRALS

In Applied Mathematics, we often come across integrals of the form  $\int_0^1 e^{-x^2} dx$  or  $\int_0^1 \sin x^2 dx$  which cannot be evaluated by any of the standard methods of integration. In such cases, we may find the value to any desired degree of accuracy by expanding their integrands as power series. An important class of such integrals is the *elliptic integrals*.

**Def.** The integral  $F(k, \phi) = \int_0^\phi \frac{dx}{\sqrt{1 - k^2 \sin^2 x}} \quad (k^2 < 1)$  ... (i)

which is a function of the two variables  $k$  and  $\phi$ , is called the *elliptic integral of the first kind with modulus  $k$  and amplitude  $\phi$* .

The integral  $E(k, \phi) = \int_0^\phi \sqrt{1 - k^2 \sin^2 x} \, dx \quad (k^2 < 1)$  ... (ii)

is called the *elliptic integral of the second kind with modulus  $k$  and amplitude  $\phi$* .

The name *elliptic integral* arose from its original application in finding the length of an elliptic arc (Fig.7.45). For instance, consider the ellipse

$$x = a \cos \phi, \quad y = b \sin \phi, \quad (a < b)$$

Then length of its arc

$$\begin{aligned} AP &= \int_0^\phi \sqrt{\left(\frac{dx}{d\phi}\right)^2 + \left(\frac{dy}{d\phi}\right)^2} d\phi = \int_0^\phi \sqrt{[(-a \sin \phi)^2 + (b \cos \phi)^2]} d\phi \\ &= \int_0^\phi \sqrt{[b^2 + (a^2 - b^2) \sin^2 \phi]} d\phi = b \int_0^\phi \sqrt{1 - \left(1 - \frac{a^2}{b^2}\right) \sin^2 \phi} d\phi \\ &= bE(k, \phi) \text{ for } k^2 = 1 - a^2/b^2 \leq 1. \end{aligned}$$




Fig. 7.45



Also the perimeter of the ellipse

$$= 4b \int_0^{\pi/2} \sqrt{1 - k^2 \sin^2 \phi} \, d\phi = 4bE(k, \pi/2).$$

This particular integral with upper limit  $\phi = \pi/2$  is called the *complete elliptic integral of the second kind* and is denoted by  $E(k)$ .

Thus 
$$E(k) = \int_0^{\pi/2} (1 - k^2 \sin^2 \phi) \, d\phi \quad (k^2 < 1) \quad \dots(iii)$$

Similarly, the *complete elliptic integral of first kind* is

$$K(k) = \int_0^{\pi/2} \frac{d\phi}{\sqrt{1 - k^2 \sin^2 \phi}} \quad (k^2 < 1) \quad \dots(iv)$$

To evaluate it, we expand the integral in the form

$$(1 - k^2 \sin^2 \phi)^{-1/2} = 1 + \frac{k^2}{2} \sin^2 \phi + \frac{3k^4}{4} \sin^4 \phi + \dots$$

This series can be shown to be uniformly convergent for all  $k$ , and may, therefore, be integrated term by term [See § 9.19-II]. Then we have

$$\begin{aligned} K(k) &= \int_0^{\pi/2} \left( 1 + \frac{k^2}{2} \sin^2 \phi + \frac{3k^4}{8} \sin^4 \phi + \frac{5k^6}{16} \sin^6 \phi + \dots \right) d\phi \\ &= \frac{\pi}{2} \left[ 1 + \left( \frac{1}{2} \right)^2 k^2 + \left( \frac{1.3}{2.4} \right)^2 k^4 + \left( \frac{1.3.5}{2.4.6} \right)^2 k^6 + \dots \right] \quad \dots(v) \end{aligned}$$

This series may be used to compute  $K$  for various values of  $k$ . In particular, if  $k = \sin 10^\circ$ ; we have

$$K = \frac{\pi}{2} (1 + 0.00754 + 0.00012 + \dots) = 1.5828 \quad \dots(vi)$$

In this way tables of the elliptic integrals are constructed. Values of  $F(k, \phi)$  and  $E(k, \phi)$  are readily available for  $0 \leq \phi \leq \pi/2$ ,  $0 < k < 1$ . (See Peirce's short tables).

**Example 7.50.** Express  $\int_0^{\pi/6} \frac{dx}{\sqrt{(\sin x)}}$  in terms of elliptic integral.

**Solution.** Put  $\cos x = \cos^2 \phi$  and  $dx = \frac{2 \cos \phi \, d\phi}{\sqrt{1 + \cos^2 \phi}}$

$$\begin{aligned} \text{Then } I &= \int_0^{\pi/2} \frac{2 \cos^2 \phi}{\sqrt{1 + \cos^2 \phi}} \, d\phi = 2 \int_0^{\pi/2} \frac{(1 + \cos^2 \phi) - 1}{\sqrt{1 + \cos^2 \phi}} \, d\phi \\ &= 2 \left\{ \int_0^{\pi/2} \sqrt{1 + \cos^2 \phi} \, d\phi - \int_0^{\pi/2} \frac{d\phi}{\sqrt{1 + \cos^2 \phi}} \right\} = 2 \left\{ \int_0^{\pi/2} \sqrt{2 - \sin^2 \phi} \, d\phi - \int_0^{\pi/2} \frac{d\phi}{\sqrt{2 - \sin^2 \phi}} \right\} \\ &= 2\sqrt{2} \int_0^{\pi/2} \sqrt{1 - 1/2 \sin^2 \phi} \, d\phi - \sqrt{2} \int_0^{\pi/2} \frac{d\phi}{\sqrt{1 - 1/2 \sin^2 \phi}} = 2\sqrt{2} E\left(\frac{1}{\sqrt{2}}\right) - \sqrt{2} K\left(\frac{1}{\sqrt{2}}\right) \end{aligned}$$

(2) **Elliptic functions.** By putting  $\sin x = t$  and  $\sin \phi = z$ , (i) becomes

$$u = \int_0^z \frac{dt}{\sqrt{[(1 - t^2)(1 - k^2 t^2)]}} \quad (k^2 < 1) \quad \dots(vii)$$

This is known as *Jacobi's form of the elliptic integral of first kind*\* whereas (i) is the *Legendre's form*†.

If  $k = 0$ , (vii) gives  $u = \sin^{-1} z$ . By analogy, we denote (vii)  $sn^{-1} z$  for a fixed non-zero value of  $k$ . This leads to the functions  $sn \, u = z = \sin \phi$  and  $cn \, u = \cos \phi$  which are called the *Jacobi's elliptic functions*.

\* See footnote p. 215.

† A French mathematician *Adrien Marie Legendre* (1752–1833) who made important contributions to number theory, special functions, calculus of variations and elliptic integrals.

The elliptic functions  $sn\ u$  and  $cn\ u$  are periodic with a period depending on  $k$  and an amplitude equal to unity. These behave somewhat like  $\sin u$  and  $\cos u$ . For instance

$$sn(0) = 0, cn(1) = 1 \quad \text{and} \quad sn(-u) = -sn(u), cn(-u) = cn(u).$$

**Example 7.51.** Show that  $\int_0^{a/2} \frac{dx}{\sqrt{(2ax - x^2)}\sqrt{(a^2 - x^2)}} = \frac{2}{3a} K\left(\frac{1}{3}\right)$ .

**Solution.** Putting  $x = \frac{a}{2}(1 - \sin \theta)$ ,  $dx = -\frac{a}{2} \cos \theta d\theta$ ,

$$2ax - x^2 = \frac{a^2}{4}(1 - \sin \theta)(3 + \sin \theta) \text{ and } a^2 - x^2 = \frac{a^2}{4}(1 + \sin \theta)(3 - \sin \theta)$$

Also when  $x = 0$ ,  $\theta = \pi/2$ ; when  $x = a/2$ ,  $\theta = 0$ .

Thus the given integral

$$= \frac{4}{a^2} \int_{\pi/2}^0 \frac{-(a/2) \cos \theta d\theta}{\sqrt{[(1 - \sin^2 \theta)(2 - \sin^2 \theta)]}} = \frac{2}{3a} \int_0^{\pi/2} \frac{d\theta}{\sqrt{[(1 - (1/3)^2 \sin^2 \theta)]}} = \frac{2}{3a} K\left(\frac{1}{3}\right).$$

## 7.18 (1) ERROR FUNCTION OR PROBABILITY INTEGRAL

The error function or the probability integral is defined as

$$erf(x) = \frac{2}{\sqrt{\pi}} \int_0^x e^{-t^2} dt.$$

This integral arises in the solution of certain partial differential equations of applied mathematics and occupies an important position in the probability theory.

The complementary error function  $erfc(x)$  is defined as  $erfc(x) = 1 - erf(x)$ .

(2) **Properties :** (i)  $erf(-x) = -erf(x)$ ; (ii)  $erf(0) = 0$

$$(iii) erf(\infty) = \frac{2}{\sqrt{\pi}} \int_0^{\infty} e^{-t^2} dt = \frac{2}{\sqrt{\pi}} \frac{\sqrt{\pi}}{2} = 1 \quad [\text{By (iii), p. 289}]$$

This proves that the total area under the Normal or Gaussian error function curve is unity – § 26.16.

### PROBLEMS 7.9

1. By means of the substitution  $k \sin x = \sin z$ , show that

$$(i) \int_0^{\pi} \frac{dx}{\sqrt{(1 - k^2 \sin^2 x)}} = \frac{1}{k} F\left(\frac{1}{k}, \phi'\right);$$

$$(ii) \int_0^{\phi} \sqrt{(1 - k^2 \sin^2 x)} dx = \left(\frac{1}{k} - k\right) F\left(\frac{1}{k}, \phi'\right) + kE\left(\frac{1}{k}, \phi'\right)$$

where  $k > 1$  and  $\phi' = \sin^{-1}(k \sin \phi)$ .

Express the following integrals in terms of elliptic integrals :

2.  $\int_0^{\pi/2} \frac{dx}{\sqrt{(1 + 3 \sin^2 x)}} \quad (\text{Kerala, M.E., 2005})$

3.  $\int_0^{\pi/2} \frac{dx}{\sqrt{(2 - \cos x)}}.$

4.  $\int_0^{\pi/2} \sqrt{(\cos x)} dx.$

5. Expand  $erf(x)$  in ascending powers of  $x$ . Hence evaluate  $erf(0)$ .

(P.T.U., 2009 S)

6. Compute (i)  $erf(0.3)$ , (ii)  $erf(0.5)$ , correct to three decimal places.

7. Show that (i)  $erf(x) + erf(-x) = 0$  (ii)  $erfc(x) + erfc(-x) = 2$

8. Prove that

$$(i) \frac{d}{dx} [erf(ax)] = \frac{2a}{\sqrt{\pi}} e^{-a^2 x^2} \quad (\text{Osmania, 2003}) \quad (ii) \frac{d}{dx} [erfc(ax)] = -\frac{2a}{\sqrt{\pi}} e^{-a^2 x^2}.$$

9. Prove that  $\int_0^{\infty} e^{-x^2 - 2ax} dx = \frac{\sqrt{\pi}}{2} e^{a^2} [1 - erf(0)]$



**7.19 OBJECTIVE TYPE OF QUESTIONS**

**PROBLEMS 7.10**

Fill up the blanks or choose the correct answer from the following problems :

1.  $\int_0^2 \int_0^x (x+y) dx dy = \dots\dots$
2.  $\int_0^1 \int_0^{1-x} dx dy \dots\dots$
3.  $\int_0^1 e^{-x^2} dx = \dots\dots$
4.  $\beta\left(\frac{1}{2}, \frac{1}{2}\right) = \dots\dots$  (V.T.U., 2010)
5.  $\Gamma(3.5) = \dots\dots$
6. The surface area of the sphere  $x^2 + y^2 + z^2 + 2x - 4y + 8z - 2 = 0$  is  $\dots\dots$
7.  $\int_0^2 \int_1^3 \int_1^2 xy^2z dz dy dx = \dots\dots$
8. If  $u = x + y$  and  $v = x - 2y$ , then the area-element  $dx dy$  is replaced by  $\dots\dots du dv$ .
9. In terms of Beta function  $\int_0^{\pi/2} \sin^7 \theta \sqrt{\cos \theta} d\theta = \dots\dots$
10. The value of  $\beta(2, 1) + \beta(1, 2)$  is  $\dots\dots$
11.  $\int_0^1 \int_1^2 xy dy dx = \dots\dots$
12. Volume bounded by  $x \geq 0, y \geq 0, z \geq 0$  and  $x^2 + y^2 + z^2 = 1$  as a triple integral  $\dots\dots$
13. Value of  $\int_0^1 \int_0^{x^2} xe^y dy dx$  is equal to  
(a)  $e/2$  (b)  $e - 1$  (c)  $1 - e$  (d)  $e/2 - 1$ . (Bhopal, 2008)
14.  $\iint x^2 y^3 dx dy$  over the rectangle  $0 \leq x \leq 1$  and  $0 \leq y \leq 3$  is  $\dots\dots$
15.  $\int_0^\pi \int_0^{a \sin \theta} r dr d\theta = \dots\dots$
16.  $\int_{x=0}^{x=3} \int_{y=0}^{y=1/x} ye^{xy} dx dy = \dots\dots$
17.  $\int_0^{\pi/2} \int_0^\infty \frac{r dr d\theta}{(r^2 + a^2)} = \dots\dots$
18.  $\int_0^1 \int_x^{\sqrt{x}} (x^2 + y^2) dx dy = \dots\dots$
19. To change cartesian coordinates  $(x, y, z)$  to spherical polar coordinate  $(r, \theta, \phi)$ ;  $dx dy dz$  is replaced by  $\dots\dots$
20.  $\int_0^2 \int_0^{x^2} e^{y/x} dy dx = \dots\dots$
21.  $\iint (x+y)^2 dx dy$  over the area bounded by the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ , is  $\dots\dots$
22.  $\int_0^1 \int_0^{\sqrt{1+x^2}} \frac{dx dy}{1+x^2+y^2} = \dots\dots$
23.  $\iint xy(x+y) dx dy$  over the area between  $y + x^2$  and  $y = x$ , is  $\dots\dots$
24. Value of  $\int_0^1 \int_x^x xy dx dy$  is  
(a) zero (b)  $-1/24$  (c)  $1/24$  (d) 24. (V.T.U., 2010)
25.  $\iint dx dy$  over the area bounded by  $x = 0, y = 0, x^2 + y^2 = 1$  and  $5y = 3$ , is  $\dots\dots$
26.  $\iint_R y dx dy$  where  $R$  is the region bounded by the parabolas  $y^2 = 4x$  and  $x^2 = 4y$ , is  $\dots\dots$
27.  $\iint (x^2 + y^2) dx dy$  in the positive quadrant for which  $x + y \leq 1$ , is  $\dots\dots$
28. Area between the parabolas  $y^2 = 4x$  and  $x^2 = 4y$  is  $\dots\dots$
29. Changing the order of integration in  $\int_{-a}^a \int_0^{\sqrt{a^2-y^2}} f(x, y) dx dy = \dots\dots$
30.  $\lceil(1/4)\rceil(3/4) = \dots\dots$  (V.T.U., 2011)
31.  $\beta(5/2, 7/2) = \dots\dots$
32.  $\int_0^\infty \int_0^x xe^{-x^2/y} dy dx = \dots\dots$
33. On changing to polar coordinates  $\int_0^{2a} \int_0^{\sqrt{2ax-x^2}} dx dy$  becomes  $\dots\dots$

34. A square lamina is immersed in the liquid with one vertex in the surface and the diagonal of length vertical. Its centre of pressure is at a depth .....
35. The centroid of the area enclosed by the parabola  $y^2 = 4x$ ,  $x$ -axis and its latus-rectum is .....
36. The moment of inertia of a uniform spherical ball of mass 10 gm and radius 2 cm about a diameter is .....
37. M.I. of a solid right circular cone (base-radius  $r$  and height  $h$ ) about its axis is .....
38.  $\operatorname{erf}_c(-x) - \operatorname{erf}(x) = \dots\dots\dots$       39.  $\int_0^1 \frac{x-1}{\log x} dx = \dots\dots\dots$       40.  $\Gamma\left(\frac{3}{2}\right) = \dots\dots\dots$
41. Value of  $\int_0^a \int_0^b \int_0^c x^2 y^2 z^2 dx dy dz$  is  
 (a)  $\frac{abc}{3}$       (b)  $\frac{a^2 b^2 c^2}{27}$       (c)  $\frac{a^3 b^3 c^3}{27}$       (d)  $\frac{a^2 b^2 c^2}{9}$
42. The integral  $\int_0^1 \int_0^{\sqrt{1-x^2}} (x+y) dy dx$  after changing the order of integration.  
 (a)  $\int_0^2 \int_0^{\sqrt{1-y^2}} (x+y) dx dy$       (b)  $\int_0^1 \int_0^{\sqrt{1-y^2}} (x+y) dx dy$   
 (c)  $\int_0^1 \int_0^{\sqrt{1+y^2}} (x+y) dx dy$       (d)  $\int_0^{-1} \int_0^{\sqrt{1-y^2}} (x+y) dx dy.$  (V.T.U., 2011)