

BLS Rules, LLC Meridian, ID 83642 BLS HCP	1	2	3	4	2	9	7	8	6	10
1 and 2 rescuer Adult BLS				-						
Verifies scene safety.	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ
Assessment: Checks response, breathing, pulse 5 - 10 seconds and activates										
emergency response system	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ
Cycle 1 of CPR										
Locates correct hand placement lower half of sternum	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ
Delivers first set of compressions (30 at 15 to 18 seconds)	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ
Compression depth at least 2"	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ
Complete chest recoil after each compression	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ
Gives 2 breaths at 1 second each with mask making chest visibly rise.										
Returns to compressions w/in 10 seconds		Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ
Delivers 2nd Cycle of CPR(Repeat steps in Cycle 1)										
Gives 2 breaths at 1 second each with mask making chest visibly rise.										
Returns to compressions w/in 10 seconds	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ
AED Arrives, Rescuer 2 could take over compressions										
Rescuer follows AED prompts (power, pad placement, Clear and analyze, clear and										
shock)		Χ	Х	Χ	Χ	Х	Х	Χ	Χ	Х
Resume compressions (if 2-rescuer, student tells instructor to resume)	Χ	Х	Χ	Χ	Х	Х	Χ	Χ	Χ	Χ
Passed	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х
Needs Remediation										
	1								į	l

verify that this information is accurate and tru	thful and that it may be confirmed as being taught in accordance with AHA	Guidelines.
Signature	Date:	

	-\/\/\	
7		

BLS HCP										
BLS Rules, LLC Meridian, ID 83642	1	2	3	4	5	9	7	8	6	10
1 and 2 Rescuer Infant BLS Skills										
Verifies scene safety.	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ
Assessment: Checks response, breathing, pulse 5 - 10 seconds and activates										
emergency response system	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ
Cycle 1 of CPR										
Locates correct finger placement in center of chest below nipple line	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ
Delivers first set of compressions (30 at 15 to 18 seconds)	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ
Compression depth at 1.5"	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ
Complete chest recoil after each compression	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ
Gives 2 breaths at 1 second each with mask making chest visibly rise.										
Returns to compressions w/in 10 seconds		Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ
Delivers 2nd CPR Cycle (Repeat steps in Cycle 1)										
Rescuer 2 arrives with BVM and begins venitlation while rescuer 1 con't compressions										
w 2-thumb encircling hands technique)		Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ
Cycle 3 of CPR										
Rescuer 1 gives 15 compressions w 2-thumb encircling hands technique	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ
Rescuer 1 gives 15 compressions in 7 - 9 seconds	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ
Rescuer 1 compresses 1.5 inches depth	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ
Complete chest recoil after each compression	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ
Cycle 4 of CPR										
Rescuer 1 gives 2 breaths with BVM at 1 second each with visible chest rise										
and resumes compressions within 10 seconds	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ	Χ
Passed	Χ	Χ	X	X	Χ	X	Χ	X	X	X
Needs Remediation										

I verify that this information is accurate and truthful and the	nat it may be confirmed as being taught in accordance with AHA Guidelines
Signature	Date: