Reasonable Suspicion Process Quick Reference

The reasonable suspicion process consists of

five essential elements.

They may not necessarily occur in this order, but it's important that they all be completed as part of your process.

OBSERVATION

Personally witness unusual/uncharacteristic behavior exhibited by a driver you supervise.

CO<u>NFIRMATION</u>

Confirm that the driver's

appearance, behavior, speech, and/or odors

are consistent with signs and symptoms of

alcohol misuse and/or drug use (see reverse side).

If you are unsure you can confirm your observations, enlist the assistance of another trained supervisor within your company.

DO<u>CUMENTATION</u>

Specifically and clearly state (in writing) your observations of the driver's

appearance, behavior, speech, and/or odors.

CO<u>NFRONTATION</u>

Discuss with the driver—in a confidential setting—your suspicion. Describe exactly what you observed and why it has prompted you to order reasonable suspicion alcohol and drug tests.

Immediately remove the driver from all safety-sensitive functions.

TE<u>STING</u>

Direct the driver to undergo testing.

Though not required by the regulations, if the test is conducted off-site, the driver should be transported to the testing facility by a person of the same gender. Never allow a driver you reasonably suspect of alcohol misuse or drug use to drive himself or herself to the testing facility.

Signs & symptoms of	Some things you should look for who	n dotormining	
alcohol misuse:	Some things you should look for when determining reasonable suspicion:		
diconol misuse:			
▲ Dulled mental processes			
▲ Constricted pupils			
▲ Sleepy or stuporous condition			
▲ Slowed reactions			
▲ Slurred speech			
Poor reflex actionsImpaired vision			
 Inparied vision Inability to divide attention 			
▲ Lowered inhibitions			
Signs & symptoms of		Some things you should look for when determining	
marijuana use:	reasonable suspicion:		
▲ Reddened eyes	Very bloodshot eyes		
✓ Slowed speech	 Muscular tremors or involuntary quivering Short attention span 		
▲ Chronic fatigue or apathy	 Disoriented behavior 		
▲ Anxiety	✓ Impaired time and distance perception		
▲ Coughing, sore throat			
Signs & symptoms of	Some things you should look for whe	n determining	
cocaine use:	reasonable suspicion:		
	▲ Teeth grinding or body tremors		
Anxiety or agitation	▲ Dilated pupils or difficulty focusing the eyes		
▲ Dilated pupils	▲ Sniffles or runny nose		
 Difficulty concentrating Fatigue or insomnia 	▲ Distorted thinking		
▲ Sweating	Uncontrolled talkativeness Future and excitable behavior		
✓ Dry mouth	Extremely excitable behavior		
Signs & symptoms of	Some things you should look for when determining		
amphetamine or meth use:	reasonable suspicion:	3	
	▲ Dilated pupils	Exaggerated reflexes	
Anxiety or restlessness	▲ Jaw grinding or frequent teeth clenching	▲ Loss of coordination	
 Profuse sweating or body tremors Difficulty for uning many 	▲ Dizziness or fainting	▲ Hallucinations	
 Difficulty focusing eyes Exaggerated reflexes 	Repetitive, bizarre behavior	▲ Signs of fatigue	
Dizziness, fainting or nausea	Depression, anxiety or mood swings	Physical collapse	
	✓ Distorted thinking		
Signs & symptoms of	Some things you should look for whe	Some things you should look for when determining	
Opioid use:	reasonable suspicion:		
opioid use:	reasonable suspicion: ▲ Constricted pupils or droopy eyelids		
Opioid use: ▲ Mood swings	 reasonable suspicion: Constricted pupils or droopy eyelids Dry mouth 		
opioid use:	 reasonable suspicion: Constricted pupils or droopy eyelids Dry mouth Low, raspy speech 		
Opioid use: ▲ Mood swings ▲ Impaired vision	 reasonable suspicion: Constricted pupils or droopy eyelids Dry mouth 		
 Opioid use: Mood swings Impaired vision Drowsiness Apathy or impaired coordination 	 reasonable suspicion: Constricted pupils or droopy eyelids Dry mouth Low, raspy speech Depressed reflexes Poor coordination 	adetermining	
 Opioid use: Mood swings Impaired vision Drowsiness Apathy or impaired coordination Signs & symptoms of	 reasonable suspicion: Constricted pupils or droopy eyelids Dry mouth Low, raspy speech Depressed reflexes Poor coordination Some things you should look for when 	n determining	
 Opioid use: Mood swings Impaired vision Drowsiness Apathy or impaired coordination Signs & symptoms of PCP use:	 reasonable suspicion: Constricted pupils or droopy eyelids Dry mouth Low, raspy speech Depressed reflexes Poor coordination 	-	
 Opioid use: Mood swings Impaired vision Drowsiness Apathy or impaired coordination Signs & symptoms of PCP use: Dizziness or diminished coordination 	 reasonable suspicion: Constricted pupils or droopy eyelids Dry mouth Low, raspy speech Depressed reflexes Poor coordination Some things you should look for when reasonable suspicion: 	-	
 Opioid use: Mood swings Impaired vision Drowsiness Apathy or impaired coordination Signs & symptoms of PCP use: Dizziness or diminished coordination Profuse sweating or convulsions 	 reasonable suspicion: Constricted pupils or droopy eyelids Dry mouth Low, raspy speech Depressed reflexes Poor coordination Some things you should look for when reasonable suspicion: Involuntary eye movements, confusion or log Disorientation to time and environment Thick, slurred speech 	-	
 Opioid use: Mood swings Impaired vision Drowsiness Apathy or impaired coordination Signs & symptoms of PCP use: Dizziness or diminished coordination Profuse sweating or convulsions Extreme mood swings 	 reasonable suspicion: Constricted pupils or droopy eyelids Dry mouth Low, raspy speech Depressed reflexes Poor coordination Some things you should look for when reasonable suspicion: Involuntary eye movements, confusion or loging the providence of the	-	
 Opioid use: Mood swings Impaired vision Drowsiness Apathy or impaired coordination Signs & symptoms of PCP use: Dizziness or diminished coordination Profuse sweating or convulsions 	 reasonable suspicion: Constricted pupils or droopy eyelids Dry mouth Low, raspy speech Depressed reflexes Poor coordination Some things you should look for when reasonable suspicion: Involuntary eye movements, confusion or log Disorientation to time and environment Thick, slurred speech 	-	