

Network Radiotherapy protocol for adjuvant and radical treatment of Vulva

SECTION 1: Treatment aims and options

Adjuvant treatment of vulva following surgery. Consider external beam and/or chemotherapy dependent on tumour stage and clinical risk factors.

Radical treatment of vulva. Consider external beam and/or chemotherapy dependent on tumour stage and clinical factors.

Chemotherapy

- Weekly concurrent Cisplatin 40mg/m²

Radiotherapy

Intent and indications	Regime, technique and RCR
<i>Adjuvant radiotherapy of Vulva</i> <i>External beam +/- chemotherapy</i>	<ul style="list-style-type: none"> 45Gy in 25# over 5 weeks Consider SIB for positive margins, extra nodal spread up to 54-64Gy / EQD2 up to 60 Gy based on histological features Prescribed to median dose (D50%) PTV SCC Cat1 +/- Concurrent Cisplatin chemotherapy
<i>Radical radiotherapy of Vulva</i> <i>External beam +/- chemotherapy</i>	<ul style="list-style-type: none"> 45Gy in 25# over 5 weeks Consider SIB / Phase 2 Boost to overall dose of 60-70Gy to macroscopic primary disease and involved nodes Prescribed to median dose (D50%) PTV SCC Cat1 +/- Concurrent Cisplatin chemotherapy