

This is a part paper delivered by Dr. Shirahata to the Functional Water Symposium in Tokyo. It does not represent the opinions of management of Healing Water Machines nor do we publish it to hold out that it is a curative or therapeutic recommendation. Please always consult your doctor for specialized medical advice.

REDUCED WATER FOR PREVENTION OF DISEASES

Dr. Sanetaka Shirahata

Graduate school of Genetic Resources Technology, Kyushu University,

6-10-1 Hakozaki, Higashi-ku, Fukuoka 812-8581, Japan.

It has long been established that reactive oxygen species (ROS) cause many types of damage to biomolecules and cellular structures, that, in turn result in the development of a variety of pathologic states such as diabetes, cancer and aging. Reduced water is defined as anti-oxidative water produced by reduction of water.

Electrolyzed reduced water (ERW) has been demonstrated to be hydrogen-rich water and can scavenge ROS in vitro (Shirahata et al., 1997). The reduction of proton in water to active hydrogen (atomic hydrogen, hydrogen radical) that can scavenge ROS is very easily caused by a weak current, compared to oxidation of hydroxyl ion to oxygen molecule. Activation of water by magnetic field, collision, minerals etc. will also produce reduced water containing active hydrogen and/or hydrogen molecule.

Several natural waters such as Hita Tenryosui water drawn from deep underground in Hita city in Japan, Nordenau water in Germany and Tlacote water in Mexico are known to alleviate various diseases. We have developed a sensitive method by which we can detect active hydrogen existing in reduced water, and have demonstrated that not only ERW but also natural reduced waters described above contain active hydrogen and scavenge ROS in cultured cells. ROS is known to cause reduction of glucose uptake by inhibiting the insulin-signaling pathway in cultured cells. Reduced water scavenged intracellular ROS and stimulated glucose uptake in the presence or absence of insulin in both rat L6 skeletal muscle cells and mouse 3T3/L1 adipocytes. This insulin-like activity of reduced water was inhibited by wortmannin that is specific inhibitor of PI-3 kinase, a key molecule in insulin signaling pathways.

Reduced water protected insulin-responsive cells from sugar toxicity and improved the damaged sugar tolerance of type 2 diabetes model mice, suggesting that reduced water may improve insulin-independent diabetes mellitus.

Cancer cells are generally exposed to high oxidative stress. Reduced water causes impaired tumor phenotypes of human cancer cells, such as reduced growth rate, morphological changes, reduced colony formation ability in soft agar, passage number-

dependent telomere shortening, reduced binding abilities of telomere binding proteins and suppressed metastasis.

Reduced water suppressed the growth of cancer cells transplanted into mice, demonstrating their anticancer effects in vivo. Reduced water will be applicable to not only medicine but also food industries, agriculture, and manufacturing industries.

Shirahata, S. et al.: Electrolyzed reduced water scavengers active oxygen species and protects DNA from oxidative damage. Biochem. Biophys. Res. Commun., 234, 269-274, 1997.