
A simple smartphone app transmits the study data to the cloud. As soon as the study is completed, the clinician can review the automated study results.

The raw data is downloaded and auto-scored differentiating obstructive and central events, providing AHI, RDI and ODI based upon True Sleep Time and Sleep Staging. Both the AHI and RDI derived from the PAT were clinically validated with an 89% correlation to PSG.² The PAT signal is an approved measure in the 2017 AASM HSAT Clinical Practice Guidelines for Adults with OSA.3





ZOLL itamar



www.itamar-medical.com

- 2- Yalamanchali S, Farajian V, Hamilton C, Pott TR, Samuelson CG, Friedman M. Diagnosis of obstructive sleep apnea by peripheral arterial tonometry: meta-analysis. JAMA Otolaryngol. Head Neck Surg. December 2013;139[12]:1343-1350
- 3- Clinical Practice Guideline for Diagnostic Testing for Adult Obstructive Sleep Apnea: An American Academy of Sleep Medicine Clinical
- 4- Hedner J. et al. A Novel Adaptive Wrist Actigraphy Algorithm for Sleep-Wake Assessment in Sleep Apnea Patients. SLEEP, Vol. 27,
- 5- Hedner J. et al. Sleep Staging Based on Automimcal Signals: A Multi-Center Validation Study. JCSM. Journal of Sleep Medicine,
- 6- Comparison of AHI using recording time versus sleep time Schutte Rodin et al., J Sleep Abs supl 2014, p. A373





WatchPAT*ONE •



First and Only Disposable HSAT





First and Only Disposable HSAT

- Same WatchPAT Simplicity, Accuracy and Reliability
- NO Return Shipment, NO Cleaning, Downloading or Charging, NO Infection Risk
- Immediate access to sleep data for interpretation
- Data interpretation anytime, anywhere with Cloud based solution

WatchPAT **ONE is designed for practices with:


- Limited access to capital
- Limited back office support
- Patients with access limitations to clinic
- Desire to grow HSAT program or facing significant backlog

WatchPAT ONE measures 7 channels (PAT signal, heart rate, oximetry, actigraphy, body position, snoring and chest motion) via three points of contact. It's 3 points of contact delivers 98% study success rate without compromising diagnostic capability.¹

Improved Accuracy with Sleep Architecture and True Sleep Time

Sleep Architecture


WatchPAT's clinically validated Sleep Architecture provides information on sleep staging, sleep efficiency and sleep and REM latency.⁴⁻⁵ It uses the PAT amplitude and pulse rate to differentiate between non-REM and REM thereby creating a Sleep Architecture (Light, Deep, REM) WatchPAT also provides the added value of detecting REM related sleep apnea with REM and non-REM AHI.



True Sleep Time

The WatchPAT uses its advanced actigraphy to differentiate between wake and sleep periods to calculate True Sleep Time. It calculates AHI and RDI using the patient's True

Sleep Time rather than the recorded time used in most commercially available HSAT's. WatchPAT's True Sleep Time reduces the risk of misdiagnosis and misclassification that has been reported to be 20% with HSAT's using total recording time.⁶



WatchPAT Automated Report and Clinical Parameters



Fully reliable. Fully disposable.

Single use to facilitate practice scalability

No limit to the number of tests that can be ordered and patients that can be tested at any given time

Immediate, highly accurate results to expedite diagnoses

Looks further than AHI to enhance diagnosis, including

- ✓ Sleep architecture
- ✓ True sleep time
- ✓ Central Sleep Apnea
- ✓ Arrhythmia Detection Feature

✓ Cheyne-Stokes respiration

- ✓ REM sleep apnea
- ✓ Sleep latency
- ✓ AHI severity scale
- ✓ Body position statistics
- √ Heart rate
- ✓ Snoring