

Common Treatable Benign Skin Conditions

Many treatable skin conditions affect people of all Fitzpatrick Skin Types. The skin conditions listed below become more common based on the following factors: as we age, increased exposure to the sun, and certain health conditions. They affect men and women. Between 40% - 70% of the population is affected by at least one of these conditions. These conditions can develop as single or multiple growths.

Treatment* has a Typical heal time of 4 – 7+ days.

Acrochordons (Skin Tags)

Small, soft, benign neoplasms commonly occur in skin folds (neck, armpits, groin). They may be flesh-colored or dark brown and range in size from 1mm to 5cm.



Dermatosis Papulosa Nigra (DPN)

A prevalent skin condition in higher Fitzpatrick skin types and almost always misclassified or described as moles. DPN is an overaccumulation of pigment resulting in benign growths.



Seborrheic Keratosis

Are common noncancerous skin growth affecting over 80 million Americans. Although benign, it can significantly impact people's physical and emotional well-being due to its appearance and can itch, bleed, or become inflamed. SK is the most frequent reason for Dermatology visits.



Cherry Angiomas

Common skin growths that can develop on any area of the body. They're also known as senile angiomas or Campbell de Morgan spots. Cherry Angiomas are often linked to pregnancy, chemical exposure, certain medical conditions, and climate.



Book your appointment today.

www.spalifesanctuary | 267.895.8319 | IG: SL.Sanctuary | FB & YouTube: Spa Life Sanctuary

*Results May Vary

