

Growth: Common or European buckthorn, *Rhamnus catharticus*, is a multi-stemmed small tree to large shrub growing 10-25 ft. tall with an upright, oval form. Glossy or smooth buckthorn (also known as European alder), *Frangula alnus*, or *Rhamnus frangula*, is a large shrub to small tree with an upright, oval form growing 10-18 ft. tall. Glossy buckthorn has several common cultivars, commonly sold prior to the regulation of both buckthorns are Restricted Invasive Species.

Leaves: Common buckthorn leaves are mostly subopposite or opposite on the stems, and are simple, rounded to egg-shaped with finely toothed margins. The leaves are very dark, dull to glossy green with 3-5 pairs of leaf veins that are sickle-shaped (curve along the margins of the leaf). Glossy buckthorn leaves alternate on stems, and are simple and oval shaped with toothless margins. Leaves are dark green and glossy, with 8-9 pairs of leaf veins that radiate out from along the central midvein. Undersides of leaves are moderately hairy to smooth. The 'Columnaris' (tall hedge buckthorn) cultivar grows 10-12 ft. tall with a narrow,

Flowers: Common buckthorn flowers are produced in May and are dioecious (i.e., male and female flowers are produced on separate plants). Flowers form in clusters and are not showy, small and yellowish-green with four petals. Glossy buckthorn flowers are produced in late May-June and sporadically in the summer. Flowers are arranged in lateral clusters, and are not showy, but creamy-green with five petals.

Fruit and Seed: Both common and glossy buckthorn have berry-like drupes that are about 1/4 in. in diameter and arranged in large clusters. Common buckthorn fruits are green changing to black in fall and each fruit contains 3-4 grooved seeds. Fruits are retained long into winter. Glossy buckthorn produces less fruit than common buckthorn. Glossy buckthorn fruit is red-brown, changing to black in late summer to early fall and the fruit falls rapidly when ripe. Each fruit contains 2-3 ungrooved seeds. Both species produce fruit annually, and fruit can serve as a food source for birds. However, fruit of both species is poisonous to humans.

Stems & Bark: Common buckthorn has upright, columnar form and low branching. The 'Asplenifolia' (fernleaf buckthorn) cultivar has finely textured, narrow, fernlike leaves and grows 6-10 ft. tall. Buckthorns leaf out very early in spring and retain their leaves late into autumn. Leaves do not change color before being shed in the fall.

Threats

- displaces native understory vegetation;
- forms an impenetrable understory layer;
- destroys wildlife habitat;
- causes long-term decline of forests by shading out other woody and herbaceous plants.

Identification



Stems & Bark: Common buckthorn has upright, columnar form and low branching. The 'Asplenifolia' (fernleaf buckthorn) cultivar has finely textured, narrow, fernlike leaves and grows 6-10 ft. tall. Buckthorns leaf out very early in spring and retain their leaves late into autumn. Leaves do not change color before being shed in the fall.

Stems & Bark: Common buckthorn has elongated, dark buds that are closely pressed against the twig. The terminal bud is often modified into a spine that is longer than the buds. Older bark resembles cherry or plum bark; it is gray to blackish-brown, and smooth, becoming roughened, with prominent, light-colored, horizontal lenticels. Glossy buckthorn has naked, hairy buds. The twigs lack the terminal spine that is typical on common buckthorn twigs. Both common and glossy buckthorns have closely-spaced, prominent leaf scars that give twigs a warty or bumpy silhouette. A cut branch of either buckthorn reveals yellow sapwood and a pinkish to orange-colored heartwood.

Stems & Bark: Common buckthorn has upright, columnar form and low branching. The 'Asplenifolia' (fernleaf buckthorn) cultivar has finely textured, narrow, fernlike leaves and grows 6-10 ft. tall. Buckthorns leaf out very early in spring and retain their leaves late into autumn. Leaves do not change color before being shed in the fall.

Stems & Bark: Common buckthorn has elongated, dark buds that are closely pressed against the twig. The terminal bud is often modified into a spine that is longer than the buds. Older bark resembles cherry or plum bark; it is gray to blackish-brown, and smooth, becoming roughened, with prominent, light-colored, horizontal lenticels. Glossy buckthorn has naked, hairy buds. The twigs lack the terminal spine that is typical on common buckthorn twigs. Both common and glossy buckthorns have closely-spaced, prominent leaf scars that give twigs a warty or bumpy silhouette. A cut branch of either buckthorn reveals yellow sapwood and a pinkish to orange-colored heartwood.

Stems & Bark: Common buckthorn has upright, columnar form and low branching. The 'Asplenifolia' (fernleaf buckthorn) cultivar has finely textured, narrow, fernlike leaves and grows 6-10 ft. tall. Buckthorns leaf out very early in spring and retain their leaves late into autumn. Leaves do not change color before being shed in the fall.



Common and Glossy BUCKTHORN

(*Rhamnus catharticus* and *Rhamnus frangula*)

A serious threat to Wisconsin woodlands

Printed References

Converse, C. K. 1984. Element Stewardship Abstract for *Rhamnus cathartica* and *Rhamnus frangula* (syn. *Frangula alnus*). The Nature Conservancy, Arlington, VA.

Glass, S. 1994. Experiment finds less herbicide needed to control buckthorn. *Restoration and Mgt. Notes*. vol. 12(1):93.

Hoffman, R. and K. Kearns (eds.). 1997. Wisconsin manual of control recommendations for ecologically invasive plants. Bureau of Endangered Resources, Wisconsin Dept. of Nat. Res., Madison, WI.

McClain, B. 1996. *Rhamnus frangula*: Smooth or glossy buckthorn. p. 65. In: J. M. Randall and J. Marinelli (eds.). *Invasive plants: Weeds of the global garden*. Brooklyn Botanic Garden Publications, Brooklyn, NY.

Samuels, A. 1996. *Rhamnus cathartica*: Common or European buckthorn. p. 64. In: J. M. Randall and J. Marinelli (eds.). *Invasive plants: Weeds of the global garden*. Brooklyn Botanic Garden Publications, Brooklyn, NY.

Websites

- <http://dnr.wi.gov/topic/invasives/fact/glossybuckthorn.html>
Wisconsin Department of Natural Resources summary on glossy buckthorn
- www.dnr.state.mn.us/invasives/terrestrialplants/woody/buckthorn
Minnesota Department of Natural Resources summary of buckthorn
- <http://wssa.net/2012/01/european-buckthorn/>
Weed Science Society of America-European Buckthorn
- www.botany.wisc.edu/wisflora
Checklist of Wisconsin vascular plants



Scan here to read more about Common Buckthorn!

<http://dnr.wi.gov/u/?q=134>

UW Extension County Offices

- | | |
|---------------------------|---------------------------|
| Adams: 608-339-4237 | Marathon: 715-261-1230 |
| Ashland: 715-682-7017 | Marinette: 715-732-7510 |
| Barron: 715-537-6250 | Marquette: 608-297-3141 |
| Bayfield: 715-373-6104 | Menominee: 715-799-4654 |
| Brown: 920-391-4610 | Milwaukee: 414-615-0550 |
| Buffalo: 608-685-6256 | Monroe: 608-269-8722 |
| Burnett: 715-349-2151 | Oconto: 920-834-6845 |
| Calumet: 920-849-1450 | Oneida: 715-365-2750 |
| Chippewa: 715-726-7950 | Outagamie: 920-832-5121 |
| Clark: 715-743-5121 | Ozaukee: 262-284-8288 |
| Columbia: 608-742-9680 | Pepin: 715-672-5214 |
| Crawford: 608-326-0223 | Pierce: 715-273-6781 |
| Dane: 608-224-3700 | Polk: 715-485-8600 |
| Dodge: 920-386-3790 | Portage: 715-346-1316 |
| Door: 920-746-2260 | Price: 715-339-2555 |
| Douglas: 715-395-1363 | Racine: 262-767-2929 |
| Dunn: 715-232-1636 | Richland: 608-647-6148 |
| Eau Claire: 715-839-4712 | Rock: 608-757-5066 |
| Florence: 715-528-4480 | Rusk: 715-532-2151 |
| Fond du Lac: 920-929-3170 | St. Croix: 715-531-1930 |
| Forest: 715-478-7797 | Sauk: 608-355-3250 |
| Grant: 608-723-2125 | Sawyer: 715-634-4839 |
| Green: 608-328-9440 | Shawano: 715-526-6136 |
| Green Lake: 920-294-4032 | Sheboygan: 920-459-5900 |
| Iowa: 608-930-9850 | Taylor: 715-748-3327 |
| Iron: 715-561-2695 | Trempealeau: 715-538-2311 |
| Jackson: 715-284-4257 | Vernon: 608-637-5276 |
| Jefferson: 920-674-7295 | Vilas: 715-479-3648 |
| Juneau: 608-847-9329 | Walworth: 262-741-4951 |
| Kenosha: 262-857-1945 | Washburn: 715-635-4444 |
| Kewaunee: 920-388-7141 | Washington: 262-335-4477 |
| La Crosse: 608-785-9593 | Waukesha: 262-548-7770 |
| Lafayette: 608-776-4820 | Waupaca: 715-258-6230 |
| Langlade: 715-627-6236 | Waushara: 920-787-0416 |
| Lincoln: 715-539-1072 | Winnebago: 920-232-1970 |
| Manitowoc: 920-683-4169 | Wood: 715-421-8440 |



This brochure was originally produced by Dr. Laura Jull, UW-Madison Woody Ornamental Extension Specialist, with layout by Jeffrey J. Strobel, UW-Extension Environmental Resources Center. It was revised and reprinted by Bernadette Williams and Kelly Kearns, WI DNR, with layout by Stephanie Miller.

This brochure was originally produced by the Urban Horticulture-Wisconsin Program of the USDA CSREES, grant number 2002-45060-01394. Originally reviewed by: Elizabeth Czarapata, author of *Invasive Plants of the Upper Midwest*; Paul Hartman, Brown County UW-Extension; Brian Hudelson, UW-Madison Plant Pathology; Kelly Kearns, Wisconsin DNR; Sharon Morrissey, Milwaukee County UW-Extension; Joe Neal, NC State University; and Bill Schmitt, UW-Madison Horticulture, and Brock Woods, Wisconsin DNR/UW-Extension.

Additional buckthorn brochures are available from your county Extension office and local Wisconsin DNR Service Center (PUB-FR-216 2019).

History

Common and glossy buckthorns were introduced into North America as ornamentals. They were planted in hedgerows and shelter-belts during the 1800s. Both species were regulated as Restricted Invasive Species in 2009 in Wisconsin. However, the glossy buckthorn cultivars 'Aspeniflora' and 'Fineline' (Ron Williams) are exempt and may be sold.

Distribution and Habitat

Although common and glossy buckthorns are native to Europe and northwestern Asia, they have readily naturalized in the northeastern and north central third of the U.S.A. and Canada. Common buckthorn invades woodlands, savannas, prairies, abandoned fields, and roadsides, forming dense thickets. Glossy buckthorn invades bogs, marshes, river banks, fens, wetlands, and pond margins, as well as dry sites such as forests, roadsides, and prairies. Both species are capable of growing in full sun and dense shade, and are quite adaptable to adverse habitats.

Other Problems

Common and glossy buckthorns are alternate hosts for the fungus (*Puccinia coronata*) that causes oat rust disease, and have been recently cited as an overwintering site for the newly introduced soybean aphid (*Aphis glycines*).

Spread

Common and glossy buckthorns are fast-growing, woody perennials. If buckthorn is not controlled, it can quickly spread to nearby areas. Forest understories can become so dense with buckthorn that native species of wildflowers and other perennials cannot compete and eventually disappear. The primary means of spread of both common and glossy buckthorn is by seeds that are eaten by birds during the harsh winter months. Common buckthorn fruit causes a severe laxative effect in animals, thereby allowing for distribution of seeds in their excrement. Buckthorns growing in full sun produce seeds a few years after establishment. In shaded habitats, fruit production may be delayed for 10-20 years. Seedlings can readily grow beneath parent buckthorn plants, as well as at the edges of forests and fields. Seedlings can become established and grow in full sun to shade. Buckthorns grow fast, thus quickly forming dense thickets. Buckthorn stumps re-sprout vigorously after plants have been cut.

Control Methods

Control of buckthorn is best achieved with early and frequent identification, and removal of isolated plants before they begin to produce seeds. Once established, buckthorn soon shades out existing vegetation and prevents establishment of native understory flora. With large infestations of buckthorn, the larger, seed-producing plants should be removed first.

Hand removal: Buckthorn seedlings can be removed by hand, if the stems are under 0.5 in. in diameter. Removal is easier when the soil is moist. Larger plants (0.5-1.5 in. diameter) can be dug or pulled using some type of mechanical device such as a Weed Wrench™. Be sure to tamp disturbed soil resulting from hand pulling to discourage re-infestation by buckthorn.

Flooding: In wetlands with lowered water tables, restoring the water to its historical levels will often kill glossy buckthorn.

Fire: Prescribed burns in early spring or fall may kill seedlings, but will only kill the tops of larger plants, which can easily re-sprout. Use of fire is best left for fire-adapted plant communities, such as prairies, savannas, and oak woodlands. Fire should not be used if it could adversely affect a plant community. Annual or biannual burning may be needed for several years to control buckthorn. Consult with a restoration specialist and include people with prescribed fire training before attempting a burn. Follow local and state fire codes and ordinances and procure local permits that may be required for a burn.

Cutting: Large plants are hard to remove by hand. These plants should be cut or girdled at the base. Remember that both common and glossy buckthorn can readily re-sprout from cut stumps, thus necessitating herbicide treatments.

Herbicides: Herbicides are an effective option for buckthorn control. Glyphosate (e.g., RoundUp Pro®, Touchdown®, Rodeo® for use

Control Methods, cont'd.

near waterways]], triclopyr (e.g., Garlon 4®, Ortho Brush-B-Gon®), and 2,4-D with triclopyr (e.g., Crossbow®) are readily available and can provide excellent control when used properly. Dyes can be added to herbicide solutions to help identify treated areas. An advantage of using triclopyr-containing herbicides (or triclopyr + 2,4-D) is that these herbicides, unlike glyphosate-containing herbicides, do not kill grasses. The presence of dense layer of grass can potentially reduce germination of, and compete with, new buckthorn seedlings. When using herbicides always read and follow all label instructions for the product that you select to insure that you use the product in the safest and most effective manner possible. If you have questions, contact your county Extension agent for advice. In general, the best time to treat buckthorn is in mid to late autumn and early winter. At this time, sap is flowing toward the roots and this allows for maximum herbicide absorption. Foliar applications should be done after a frost, as both common and glossy buckthorns retain their leaves and continue growing much later into the fall than other plants. Spring is the least effective time for treating cut stumps (see below), because sap is flowing away from the roots.



There are three common methods for applying herbicides for buckthorn control: cut stump, basal bark, and foliar sprays. Once again, check the label of the herbicide that you choose for the legal application rates and methods. Larger buckthorn plants (greater than 6" in diameter) are best treated using the cut stump method. As the name of this technique implies, buckthorn plants should be cut and the stumps treated immediately with a herbicide solution. Following the label, use the higher rate, often 20-25% (by volume) to translocate to the roots and prevent re-sprouting. Herbicide solutions are typically applied using a low-pressure hand sprayer, a spray bottle, wick applicator, or sponge paintbrush. Spray treatments should be directed to the vascular tissue of the cut stem located just inside of the bark. Follow-up treatments may be necessary for the next several years if plants re-sprout. Smaller buckthorn plants (less than 6" in diameter) can be treated using the cut stump method or the basal bark method. These plants need not be cut before herbicide is applied. Apply the rate indicated on the label, typically near 12.5% (by volume) in an oil-based solution to the base of the plant, wetting the bark from the soil line up to about 12-15 in. above the soil line. Triclopyr is typically used for basal bark treatment. Do not use oil-based formulations at temperatures above 80° F. If you have a large number of buckthorn seedlings, foliar sprays of glyphosate, triclopyr, or triclopyr + 2,4-D may be the most effective means of control.

Foliar herbicides can be applied using a backpack sprayer. Foliar applications tend to require a lower concentration of herbicide than stump applications. Dilute sprays are typically 1-3% (by volume). Foliar sprays are more likely to damage or kill non-target vegetation.

Biological Control: Biological control agents are not available to control either common or glossy buckthorn.

Education: One of the best ways to insure adequate control of buckthorn is education.

Tell your neighbors about buckthorn. A neighbor's buckthorn can produce large amounts of seed that can be disseminated into your yard, and the surrounding neighborhood. Encourage your neighbors to remove their buckthorn and monitor their yards for seedlings in the future.



© Photo by Elizabeth J. Czarapata