

Brightside Specialty Pharmacy

431 N. Tustin Ave., STE C, Santa Ana, CA 92705

Phone 714-707-5115

Date :8/10/2022

Patient Education for AMOXICILLIN 500 MG CAPSULE



Must complete full course of therapy Space doses evenly Do not use if allergic to penicillins Tell doctor what medicines you are taking Physician must know patient's complete medical history Call doctor if rash or severe diarrhea occur. Oral or vaginal yeast infections may occur due to changes in flora.

Must use for full length of treatment Space doses evenly throughout the day Do not take if you are allergic to penicillin Tell doctor what medicines you are taking Tell doctor your complete medical history Call DR if rash or severe diarrhea occur. Watch for oral thrush or vaginal yeast infections.

.....Education Section.....

IMPORTANT:

HOW TO USE THIS INFORMATION: This is a summary and

does NOT have all possible information about this product. This information does not assure that this product is safe, effective, or appropriate for you. This information is not individual medical advice and does not substitute for the advice of your health care professional. Always ask your health care professional for complete information about this product and your specific health needs.

AMOXICILLIN - ORAL

(a-MOX-i-SIL-in)

COMMON BRAND NAME(S):

Amoxil

USES:

Amoxicillin is used to treat a wide variety of bacterial

infections. This medication is a penicillin-type antibiotic. It works by stopping the growth of bacteria.

This antibiotic treats only bacterial infections. It will not work for viral infections (such as common cold, flu). Using any antibiotic when it is not needed can cause it to not work for future infections.

Amoxicillin is also used with other medications to treat stomach/intestinal ulcers caused by the bacteria *H. pylori* and to prevent the ulcers from returning.

OTHER USES:

This section contains uses of this drug that are not

listed in the approved US professional labeling for the drug but that may be prescribed by your health care professional. Use this drug for a condition that is listed in this section only if it has been so prescribed by your health care professional.

This drug may also be used before medical/dental procedures by people with certain heart conditions (such as heart valve disease) to reduce the risk of a serious heart infection.

HOW TO USE:

Take this medication by mouth with or without food

as directed by your doctor, usually every 8 or 12 hours. The dosage is based on your medical condition and response to treatment.

Drink plenty of fluids while using this medication unless your doctor tells you otherwise.

For the best effect, take this antibiotic at evenly spaced times. To help you remember, take this medication at the same time(s) every day.

Continue to take this medication until the full prescribed amount is finished, even if symptoms disappear after a few days. Stopping the medication too early may allow bacteria to continue to grow, which may result in a return of the infection.

Tell your doctor if your condition lasts or gets worse.

SIDE EFFECTS:

Nausea, vomiting, or diarrhea may occur. If any of

these effects last or get worse, tell your doctor or pharmacist promptly.

Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects.

Use of this medication for prolonged or repeated periods may result in oral thrush or a new vaginal yeast infection (oral or vaginal fungal infection). Contact your doctor if you notice white patches in your mouth, a change in vaginal discharge or other new symptoms.

Tell your doctor right away if you have any serious side effects, including: signs of liver disease (such as nausea/vomiting that doesn't stop, loss of appetite, stomach/abdominal pain, yellowing eyes/skin, dark urine), easy bruising or bleeding.

This medication may rarely cause a severe intestinal condition due to a bacteria called *C. difficile*. This condition may occur during treatment or weeks to months after treatment has stopped.

Tell your doctor right away if you develop:

diarrhea that doesn't

stop, abdominal or stomach pain/cramping, blood/mucus in your stool.

If you have these symptoms, do not use anti-diarrhea or opioid products because they may make symptoms worse.

A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: fever that doesn't go away, new or worsening lymph node swelling, rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing.

Amoxicillin can commonly cause a mild rash that is usually not serious. However, you may not be able to tell it apart from a rare rash that could be a sign of a severe allergic reaction. Get medical help right away if you develop any rash.

This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist.

In the US -

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch.

In Canada - Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345.

PRECAUTIONS:

Before taking amoxicillin, tell your doctor or

pharmacist if you are allergic to it; or to penicillin or cephalosporin antibiotics; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details.

Before using this medication, tell your doctor or pharmacist your medical history, especially of: kidney disease, a certain type of viral infection (infectious mononucleosis).

Amoxicillin may cause live bacterial vaccines (such as typhoid vaccine) to not work well. Tell your health care professional that you are using amoxicillin before having any immunizations/vaccinations.

Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products).

During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor.

Amoxicillin passes into breast milk. Consult your doctor before breast-feeding.

DRUG INTERACTIONS:

Drug interactions may change how your

medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval.

A product that may interact with this drug is: methotrexate.

Amoxicillin may cause false positive results with certain diabetic urine testing products (cupric sulfate-type). This drug may also affect the results of certain lab tests. Make sure

laboratory personnel and your doctors know you use this drug.

OVERDOSE:

If someone has overdosed and has serious symptoms such

as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Symptoms of overdose may include: diarrhea that doesn't stop, severe vomiting, a severe decrease in the amount of urine, or seizures.

NOTES:

Do not share this medication with others.

This medication has been prescribed for your current condition only. Do not use it later for another infection unless your doctor tells you to.

With prolonged treatment, laboratory and/or medical tests (such as kidney and liver function, complete blood counts) may be performed periodically to monitor your progress or check for side effects. Consult your doctor for more details.

MISSED DOSE:

If you miss a dose, use it as soon as you remember.

If it is near the time of the next dose, skip the missed dose. Use your next dose at the regular time. Do not double the dose to catch up.

STORAGE:

Store at room temperature away from light and moisture.

Do not store in the bathroom. Keep all medications away from children and pets.

Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

Information last revised May 2022.

Copyright(c) 2022 First Databank, Inc.