

# Brightside Specialty Pharmacy

431 N. Tustin Ave., STE C, Santa Ana, CA 92705

Phone 714-707-5115

Date :8/10/2022

## Patient Education for

### ASPIRIN EC 81 MG TABLET



Take with food to lessen gastrointestinal upset Aspirin smells like vinegar when it has expired Not recommended for use in children < 12 yrs Call doctor if symptoms of GI bleeding Physician must know patient's complete medical history Not recommended during pregnancy or breast feeding Do not use this medicine longer than label or doctor directs. Swallow whole with 8 oz. water. Remain erect for 10 minutes.

Best to take with food to lessen stomach upset Do not take this medicine if it smells like vinegar Do not use in kids less than 12 years Call Dr if you have black stools or stomach pain Tell doctor your complete medical history Do not take while breast feeding or when pregnant Do not use longer than label or doctor directs. Swallow whole with water. Do not lie down for 10min

#### .....Education Section.....

#### **IMPORTANT:**

HOW TO USE THIS INFORMATION: This is a summary and

does NOT have all possible information about this product. This information does not assure that this product is safe, effective, or appropriate for you. This information is not individual medical advice and does not substitute for the advice of your health care professional. Always ask your health care professional for complete information about this product and your specific health needs.

ASPIRIN - ORAL

(AS-pir-in)

#### **COMMON BRAND NAME(S):**

Bayer Aspirin, Easprin, Ecotrin

#### **USES:**

Aspirin is used to reduce fever and relieve mild to

moderate pain from conditions such as muscle aches, toothaches, common cold, and headaches. It may also be used to reduce pain and swelling in conditions such as arthritis. Aspirin is known as a salicylate and a nonsteroidal anti-inflammatory drug (NSAID). It works by blocking a certain natural substance in your body to reduce pain and swelling. Consult your doctor before treating a child younger than 12 years.

Your doctor may direct you to take a low dose of aspirin to prevent blood clots. This effect reduces the risk of stroke and heart attack. If you have recently had surgery on clogged arteries (such as bypass surgery, carotid endarterectomy, coronary stent), your doctor may direct you to use aspirin in low doses as a "blood thinner" to prevent blood clots.

#### **HOW TO USE:**

If you are taking this medication for

self-treatment, follow all directions on the product package. If you have any questions, ask your doctor or pharmacist. If your doctor has directed you to take this medication, take it exactly as prescribed.

Take this medication by mouth. Drink a full glass of water (8 ounces/240 milliliters) with it unless your doctor tells you

otherwise. Do not lie down for at least 10 minutes after you have taken this drug. If stomach upset occurs while you are taking this medication, you may take it with food or milk.

Swallow enteric-coated tablets whole. Do not crush or chew enteric-coated tablets. Doing so can increase stomach upset.

Do not crush or chew extended-release tablets or capsules. Doing so can release all of the drug at once, increasing the risk of side effects. Also, do not split extended-release tablets unless they have a score line and your doctor or pharmacist tells you to do so. Swallow the whole or split tablet without crushing or chewing.

The dosage and length of treatment are based on your medical condition and response to treatment. Read the product label to find recommendations on how many tablets you can take in a 24-hour period and how long you may self-treat before seeking medical advice. Do not take more medication or take it for longer than recommended unless directed by your doctor. Use the smallest effective dose. Consult your doctor or pharmacist if you have any questions.

If you are taking this medication for self-treatment of headache, seek immediate medical attention if you also have trouble speaking, weakness on one side of the body, or sudden vision changes. Before using this drug, consult a doctor or pharmacist if you have headaches caused by head injury, coughing, or bending, or if you have a headache with vomiting that is severe or doesn't stop, fever, and stiff neck.

If you are taking this medication as needed (not on a regular schedule), remember that pain medications work best if they are used as the first signs of pain occur. If you wait until the pain has worsened, the medicine may not work as well. Aspirin with a special coating (enteric coating) or slow release may take longer to stop pain because it is absorbed more slowly. Ask your doctor or pharmacist to help select the best type of aspirin for you.

You should not take this medication for self-treatment of pain for longer than 10 days. You should not use this drug to self-treat a fever that lasts longer than 3 days. In these cases, consult a doctor because you may have a more serious condition. Tell your doctor promptly if you develop ringing in the ears or difficulty hearing.

If your condition lasts or gets worse (such as new or unusual

symptoms, redness/swelling of the painful area, pain/fever that does not go away or gets worse) or if you think you may have a serious medical problem, tell your doctor promptly.

**SIDE EFFECTS:**

Upset stomach and heartburn may occur. If either

of these effects last or get worse, tell your doctor or pharmacist promptly.

If your doctor has directed you to use this medication, remember that your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects.

Tell your doctor right away if you have any serious side effects, including: nausea/vomiting that is severe or doesn't stop, easy bruising/bleeding, difficulty hearing, ringing in the ears, signs of kidney problems (such as change in the amount of urine), unexplained tiredness, dizziness, dark urine, yellowing eyes/skin.

This drug may rarely cause serious bleeding from the stomach/intestine or other areas of the body. If you notice any of the following very serious side effects, get medical help right away: stomach/abdominal pain that is severe or doesn't go away, black/tarry stools, vomit that looks like coffee grounds, trouble speaking, weakness on one side of the body, sudden vision changes or severe headache.

A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: fever, swollen lymph nodes, rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing.

This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist.

In the US -

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

In Canada - Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345.

**PRECAUTIONS:**

Before taking aspirin, tell your doctor or

pharmacist if you are allergic to it; or to other salicylates (such as choline salicylate); or to other pain relievers or fever reducers (NSAIDs such as ibuprofen, naproxen); or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details.

This medication should not be used if you have certain medical conditions. Before using this medicine, consult your doctor or pharmacist if you have: bleeding/blood-clotting disorders (such as hemophilia, vitamin K deficiency, low platelet count).

If you have any of the following health problems, consult your doctor or pharmacist before using this medication: kidney disease, liver disease, diabetes, stomach problems (such as ulcers, heartburn, stomach pain), aspirin-sensitive asthma (a

history of worsening breathing with runny/stuffy nose after taking aspirin or other NSAIDs), growths in the nose (nasal polyps), gout, certain enzyme deficiencies (pyruvate kinase or G6PD deficiency).

This medicine may cause stomach bleeding. Daily use of alcohol and tobacco, especially when combined with this product, may increase your risk for this side effect. Limit alcoholic beverages, and stop smoking. Check with your doctor or pharmacist for more information.

Before having surgery, tell your doctor or dentist that you are taking this medication.

Children and teenagers less than 18 years old should not take aspirin if they have chickenpox, flu, or any undiagnosed illness or if they have recently received a vaccine. In these cases, taking aspirin increases the risk of Reye's syndrome, a rare but serious illness. Tell your doctor promptly if you see changes in behavior with nausea and vomiting. This may be an early sign of Reye's syndrome.

Older adults may be more sensitive to the side effects of this drug, especially stomach/intestinal bleeding and ulcers.

Aspirin is not recommended for use to treat pain or fever during pregnancy. Before using this medication, women of childbearing age should talk with their doctor(s) about the benefits and risks. Tell your doctor if you are pregnant or if you plan to become pregnant. This medication may harm an unborn baby and cause problems with normal labor/delivery. It is not recommended for use in pregnancy from 20 weeks until delivery. If your doctor decides that you need to use this medication between 20 and 30 weeks of pregnancy, you should use the lowest effective dose for the shortest possible time. In some cases, low-dose aspirin (usually 81-162 milligrams a day) may be used safely during pregnancy to prevent certain conditions. Talk to your doctor for more details.

Aspirin passes into breast milk. When used in large amounts (such as to treat pain or fever), it may harm a nursing infant and breast-feeding while using this drug is not recommended. However, low-dose aspirin for heart attack or stroke prevention may be used if directed by your doctor. Consult your doctor before breast-feeding.

**DRUG INTERACTIONS:**

Drug interactions may change how your

medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval.

Some products that may interact with this drug include: mifepristone, acetazolamide, "blood thinners" (such as warfarin, heparin), corticosteroids (such as prednisone), dichlorphenamide, methotrexate, valproic acid, herbal medications (such as ginkgo biloba).

Before using this product, consult your doctor if you have recently received certain live vaccines (such as varicella vaccine, live flu vaccine).

Check all prescription and nonprescription medicine labels

carefully since many medications contain pain relievers/fever reducers known as NSAIDs (non-steroidal anti-inflammatory drugs such as ibuprofen, ketorolac, naproxen). To prevent an overdose of aspirin, read the labels carefully before taking other pain relievers or cold products to make sure they do not contain aspirin. Ask your pharmacist about using these products safely.

Daily use of NSAIDs (such as ibuprofen) may decrease aspirin's ability to prevent heart attack/stroke. If you are taking low-dose aspirin for prevention of heart attack/stroke, consult your doctor or pharmacist for more details and to discuss other possible treatments (such as acetaminophen) for your pain/fever.

This medication may interfere with certain laboratory tests (including certain urine sugar tests), possibly causing false test results. Make sure laboratory personnel and all your doctors know you are taking this medication.

**OVERDOSE:**

If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Symptoms of overdose may include: burning pain in the throat/stomach, confusion, mental/mood changes, fainting, weakness, ringing in the ears, fever, rapid breathing, change in the amount of urine, seizures, loss of consciousness.

**NOTES:**

If you use this medication regularly or at high doses, laboratory and/or medical tests (such as liver and kidney function tests, blood count, salicylate level) may be performed to monitor your progress or check for side effects. Consult your doctor for more details.

There are many different aspirin products. Some have special coatings and some are long-acting. Ask your doctor or pharmacist to recommend the best product for you.

**MISSED DOSE:**

If your doctor directs you to take this drug on a regular schedule (not just "as needed") and you miss a dose, take it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up.

**STORAGE:**

Store at room temperature away from moisture and light. Different brands of this medication may have different storage needs. Check the product package for instructions on how to store your brand, or ask your pharmacist. Do not store in the bathroom. Do not use any aspirin product that has a strong vinegar-like smell. Keep all medications away from children and pets.

Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

Information last revised March 2022.  
Copyright(c) 2022 First Databank, Inc.