

Brightside Specialty Pharmacy

431 N. Tustin Ave., STE C, Santa Ana, CA 92705

Phone 714-707-5115

Date :8/10/2022

Patient Education for CARISOPRODOL 350 MG TABLET

May be habit forming. Physician must know patient's complete medical history. Not recommended in the 1st trimester of pregnancy. Avoid taking with other CNS depressant drugs or alcohol. May cause drowsiness or dizziness. Use caution driving. Check with doctor before increasing dose or frequency. Call doctor if no improvement in 3 weeks.

May be habit forming. Tell doctor your complete medical history. Tell doctor if you think you are pregnant. Avoid alcohol/other drugs that make you sleepy. May make you drowsy or dizzy. Drive with caution. Call Dr before increasing dose or frequency. Call doctor if you are not better in 3 weeks.

.....Education Section.....

IMPORTANT:

HOW TO USE THIS INFORMATION: This is a summary and

does NOT have all possible information about this product. This information does not assure that this product is safe, effective, or appropriate for you. This information is not individual medical advice and does not substitute for the advice of your health care professional. Always ask your health care professional for complete information about this product and your specific health needs.

CARISOPRODOL - ORAL

(kar-iss-oh-PRO-dole)

COMMON BRAND NAME(S):

Soma

USES:

Carisoprodol is used short-term to treat muscle pain and

discomfort. It is usually used along with rest, physical therapy, and other treatments. It works by helping to relax the muscles.

HOW TO USE:

Take this medication by mouth with or without food

as directed by your doctor. Do not increase your dose or use this drug more often or for longer than prescribed. Your condition will not improve any faster, and your risk of side effects will increase.

The dosage is based on your medical condition and response to treatment. This medication should only be used short-term (for 3 weeks or less) unless directed by your doctor.

If you suddenly stop using this medication, you may have withdrawal symptoms (such as stomach cramps, trouble sleeping, headache, nausea). To help prevent withdrawal, your doctor may lower your dose slowly. Withdrawal is more likely if you have used carisoprodol for a long time or in high doses. Tell your doctor or pharmacist right away if you have withdrawal.

Though it helps many people, this medication may sometimes cause addiction. This risk may be higher if you have a substance use disorder (such as overuse of or addiction to drugs/alcohol).

Take this medication exactly as prescribed to lower the risk of

addiction. Ask your doctor or pharmacist for more details.

Tell your doctor if your condition lasts after 2 to 3 weeks or if it gets worse.

SIDE EFFECTS:

Dizziness, drowsiness, or headache may occur. If

any of these effects last or get worse, tell your doctor or pharmacist promptly.

Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects.

Tell your doctor right away if you have any serious side effects, including: confusion.

A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing.

This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist.

In the US -

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch.

In Canada - Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345.

PRECAUTIONS:

Before taking carisoprodol, tell your doctor or

pharmacist if you are allergic to it; or to meprobamate, tybamate, or mebutamate; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details.

Before using this medication, tell your doctor or pharmacist your medical history, especially of: a certain blood disorder (acute intermittent porphyria), kidney disease, liver disease,

seizure, personal or family history of a substance use disorder (such as overuse of or addiction to drugs/alcohol).

This drug may make you dizzy or drowsy. Alcohol or marijuana (cannabis) can make you more dizzy or drowsy. Do not drive, use machinery, or do anything that needs alertness until you can do it safely. Avoid alcoholic beverages. Talk to your doctor if you are using marijuana (cannabis).

Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products).

Older adults may be more sensitive to the side effects of this drug, especially drowsiness, or confusion. These side effects can increase the risk of falling.

During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor.

This medication passes into breast milk and may have undesirable effects on a nursing infant. Consult your doctor before breast-feeding.

DRUG INTERACTIONS:

Drug interactions may change how your

medications work or increase your risk for serious side effects.

This document does not contain all possible drug interactions.

Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval.

Tell your doctor or pharmacist if you are taking other products that cause drowsiness such as opioid pain or cough relievers (such as codeine, hydrocodone), alcohol, marijuana (cannabis), drugs for sleep or anxiety (such as alprazolam, lorazepam, zolpidem), other muscle relaxants (such as cyclobenzaprine, methocarbamol), or antihistamines (such as cetirizine, diphenhydramine).

Check the labels on all your medicines (such as allergy or cough-and-cold products) because they may contain ingredients that cause drowsiness. Ask your pharmacist about using those products safely.

OVERDOSE:

If someone has overdosed and has serious symptoms such

as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Symptoms of overdose may include: severe drowsiness/dizziness, seizures, slow/shallow breathing, mental/mood changes (such as confusion, hallucinations), inability to move your legs/arms, shaky/unsteady movement, vision changes (such as blurred vision).

NOTES:

Do not share this medication with others. Sharing it is

against the law.

This medication has been prescribed for your current condition only. Do not use it later for another condition unless your doctor directs you to do so. A different medication may be necessary in that case.

MISSED DOSE:

If you miss a dose, take it as soon as you

remember. If it is near the time of the next dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up.

STORAGE:

Store at room temperature away from light and moisture.

Do not store in the bathroom. Keep all medications away from children and pets.

Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

Information last revised March 2022.

Copyright(c) 2022 First Databank, Inc.