

Brightside Specialty Pharmacy

431 N. Tustin Ave., STE C, Santa Ana, CA 92705

Phone 714-707-5115

Date :8/10/2022

Patient Education for FLUTICASONE PROP 50 MCG SPRAY



If available, dispense patient package insert and explain to pt Blow nose gently before dose Physician must know patient's complete medical history Compliance important even during symptom-free periods Take at the same time daily Shake gently Prime before use. Instruct patient on proper number of sprays. May take several days for full benefit of therapy.

If available, read the patient package insert Blow nose gently before dose Tell doctor your complete medical history Take doses regularly, even when feeling well Take at the same time every day Shake gently Test spray before first use; look for fine mist May take several days for full benefit.

.....Education Section.....

IMPORTANT:

HOW TO USE THIS INFORMATION: This is a summary and

does NOT have all possible information about this product. This information does not assure that this product is safe, effective, or appropriate for you. This information is not individual medical advice and does not substitute for the advice of your health care professional. Always ask your health care professional for complete information about this product and your specific health needs.

FLUTICASONE SPRAY - NASAL

(floo-TIK-a-son)

COMMON BRAND NAME(S):

Flonase

USES:

Fluticasone is used to relieve seasonal and year-round

allergic and non-allergic nasal symptoms, such as stuffy/runny nose, itching, and sneezing. It can also help relieve allergy eye symptoms such as itchy, watery eyes. This medication belongs to a class of drugs known as corticosteroids. It works in your nose to block the effects of substances that cause allergies (such as pollen, pet dander, dust mites, mold) and to reduce swelling.

HOW TO USE:

If you are using the over-the-counter product to

self-treat, read and follow all directions on the product package before using this medication. If your doctor has prescribed this medication, read the Patient Information Leaflet before you start using fluticasone and each time you get a refill. If you have any questions, ask your doctor or pharmacist.

Use this medication in the nose as directed by your doctor or the product package, usually once or twice a day. Do not spray in your eyes.

Gently blow your nose before using this drug. Shake the container gently before each use. Follow the instructions on how to properly prime the bottle if you are using it for the first time or if you have not used it for a week or longer.

The dosage is based on your age, medical condition, and

response to treatment. Do not increase your dose or use this drug more often or for longer than directed. Your condition will not improve any faster, and your risk of side effects will increase. You may be directed to start with a higher dose of this drug for the first several days until you have begun to feel better, then decrease your dose. Children may need to use this drug for a shorter amount of time to lower the risk of side effects. If a child is using the over-the-counter product, read the package information to see how long he/she should use it and when you should check with the doctor.

Use this medication regularly to get the most benefit from it. To help you remember, use it at the same time(s) each day.

Clean the applicator regularly. Keep track of the number of sprays used from the container. Discard the container after you have used the number of sprays on the package label.

This medication does not work right away. You may feel an effect as soon as 12 hours after starting treatment, but it may take several days before you get the full benefit. If your condition does not improve after 1 week, or if it worsens, stop using this medication and consult your doctor or pharmacist. If you think you may have a serious medical problem, get medical help right away.

SIDE EFFECTS:

Nasal dryness/irritation, nausea, or vomiting may

occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly.

If your doctor has directed you to use this medication, remember that your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects.

Tell your doctor right away if you have any serious side effects, including: sore throat that doesn't go away, eye pain, facial pain, signs of nasal damage (such as pain, severe nosebleeds, a constant whistling sound from your nose), white patches in your nose/back of your throat, pain when swallowing.

Rarely, it is possible that corticosteroids given in the nose will be absorbed into the bloodstream. This can lead to side effects of too much corticosteroid. These side effects are more likely in children and people who use this medication for a long

time and in high doses. Tell your doctor right away if any of the following side effects occur: unusual/extreme tiredness, weight loss, headache, swelling ankles/feet, increased thirst/urination, vision problems.

A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing.

This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist.

In the US - Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch.

In Canada - Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345.

PRECAUTIONS:

Before using fluticasone, tell your doctor or

pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details.

Before using this medication, tell your doctor or pharmacist your medical history, especially of: recent nose problems (such as injury, ulcers, surgery), infections (including tuberculosis, herpes eye infection), certain eye problems (glaucoma, cataracts), liver disease.

Avoid contact with people who have infections that may spread to others (such as chickenpox, measles, flu). Consult your doctor if you have been exposed to an infection or for more details.

Rarely, using corticosteroid medications for a long time can make it more difficult for your body to respond to physical stress. Before having surgery or emergency treatment, or if you get a serious illness/injury, tell your doctor or dentist that you are using this medication or have used this medication within the past few months.

Though it is unlikely, this medication may temporarily slow down a child's growth if used for a long time. See the doctor regularly so your child's height can be checked.

During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor.

It is unknown if this drug passes into breast milk. Consult your doctor before breast-feeding.

DRUG INTERACTIONS:

Drug interactions may change how your

medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval.

Other medications can affect the removal of fluticasone from

your body, which may affect how fluticasone works. Examples include HIV protease inhibitors (such as lopinavir, ritonavir), among others.

OVERDOSE:

If someone has overdosed and has serious symptoms such

as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center.

NOTES:

Do not share this medication with others.

If you use this medication for a long time, laboratory and/or medical tests (such as nose exams, height measurement in children) may be performed to monitor your progress and check for side effects.

Avoid substances (such as pollen, pet dander, dust mites, mold, and smoke) that can worsen allergy symptoms.

MISSED DOSE:

If you are using this product on a regular schedule

and miss a dose, use it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Use your next dose at the regular time. Do not double the dose to catch up.

STORAGE:

Different brands of this medication have different

storage needs. Check the product package for instructions on how to store your brand, or ask your pharmacist. Keep all medications away from children and pets.

Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

Information last revised December 2021.

Copyright(c) 2022 First Databank, Inc.