

Brightside Specialty Pharmacy

431 N. Tustin Ave., STE C, Santa Ana, CA 92705

Phone 714-707-5115

Date :8/10/2022

Patient Education for

GLIPIZIDE 5 MG TABLET



Call Dr. if signs of hypoglycemia and know to self-treat acutely May cause photosensitive reactions Use sunscreen Use with caution in pregnancy Monitor blood glucose to check for efficacy and avoid hypoglycemia Diet, exercise, and weight control are important for therapy success Many drug-drug interactions possible with this drug May cause drowsiness or dizziness Use caution driving. Reduce or limit drinking alcohol

Call doctor if you feel faint or sweaty/confused Skin may be more prone to sunburn; Use sunscreen Tell doctor if you think you are pregnant Check blood glucose levels regularly Diet and exercise will improve therapy response Review all drugs you are taking with your doctor May make you drowsy or dizzy. Drive with caution Reduce or limit drinking alcohol

.....Education Section.....

IMPORTANT:

HOW TO USE THIS INFORMATION: This is a summary and

does NOT have all possible information about this product. This information does not assure that this product is safe, effective, or appropriate for you. This information is not individual medical advice and does not substitute for the advice of your health care professional. Always ask your health care professional for complete information about this product and your specific health needs.

GLIPIZIDE TABLET - ORAL

(GLIP-i-zide)

COMMON BRAND NAME(S):

Glucotrol

USES:

Glipizide is used with a proper diet and exercise program

to control high blood sugar in people with type 2 diabetes. It may also be used with other diabetes medications. Controlling high blood sugar helps prevent kidney damage, blindness, nerve problems, loss of limbs, and sexual function problems. Proper control of diabetes may also lessen your risk of a heart attack or stroke. Glipizide belongs to the class of drugs known as sulfonylureas. It lowers blood sugar by causing the release of your body's natural insulin.

HOW TO USE:

Take this medication by mouth 30 minutes before

breakfast or the first meal of the day as directed by your doctor, usually once daily. Some patients, especially those taking higher doses, may be directed to take this drug twice a day. The dosage is based on your medical condition and response to treatment.

To reduce your risk of side effects, your doctor may direct you to start this medication at a low dose and gradually increase your dose. Follow your doctor's instructions carefully.

If you are already taking another diabetes drug (such as chlorpropamide), follow your doctor's directions carefully for stopping the old drug and starting glipizide.

Colesevelam can decrease the absorption of glipizide. If you are taking colesevelam, take glipizide at least 4 hours before taking colesevelam.

Use this medication regularly to get the most benefit from it. To help you remember, take it at the same time(s) each day.

Tell your doctor if your condition does not improve or if it worsens (your blood sugar is too high or too low).

SIDE EFFECTS:

Nausea, vomiting, loss of appetite, diarrhea,

constipation, upset stomach, headache, and weight gain may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly.

Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects.

Tell your doctor right away if you have any serious side effects, including: signs of infection (such as sore throat that doesn't go away, fever), easy bleeding/bruising, stomach pain, yellowing eyes/skin, dark urine, unusual tiredness/weakness, unusual/sudden weight gain, mental/mood changes, swelling hands/feet, seizures.

This medication can cause low blood sugar (hypoglycemia). This may occur if you do not consume enough calories from food or if you do unusually heavy exercise. Symptoms of low blood sugar include sudden sweating, shaking, fast heartbeat, hunger, blurred vision, dizziness, or tingling hands/feet. It is a good habit to carry glucose tablets or gel to treat low blood sugar. If you don't have these reliable forms of glucose, rapidly raise your blood sugar by eating a quick source of sugar such as table sugar, honey, or candy, or drink fruit juice or non-diet soda. Tell your doctor right away about the reaction and the use of this product. To help prevent low blood sugar, eat meals on a regular schedule, and do not skip meals. Check with your doctor or pharmacist to find out what you should do if you miss a meal.

Symptoms of high blood sugar (hyperglycemia) include thirst, increased urination, confusion, drowsiness, flushing, rapid breathing, and fruity breath odor. If these symptoms occur, tell your doctor right away. Your dosage may need to be increased.

A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing.

This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist.

In the US -

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch.

In Canada - Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345.

PRECAUTIONS:

Before taking glipizide, tell your doctor or

pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details.

Before using this medication, tell your doctor or pharmacist your medical history, especially of: liver disease, kidney disease, thyroid disease, certain hormonal conditions (adrenal/pituitary insufficiency, syndrome of inappropriate secretion of antidiuretic hormone-SIADH), electrolyte imbalance (hyponatremia).

You may experience blurred vision, dizziness, or drowsiness due to extremely low or high blood sugar. Do not drive, use machinery, or do any activity that requires alertness or clear vision until you are sure you can perform such activities safely.

Limit alcohol while taking this medication because it can increase your risk of developing low blood sugar. Alcohol can rarely interact with glipizide and cause a serious reaction (disulfiram-like reaction) with symptoms such as facial flushing, nausea, vomiting, dizziness, or stomach pain. Consult your doctor or pharmacist about the safe use of alcohol.

It may be harder to control your blood sugar when your body is stressed (such as due to fever, infection, injury, or surgery). Consult your doctor because this may require a change in your treatment plan, medications, or blood sugar testing.

This medication may make you more sensitive to the sun. Limit your time in the sun. Avoid tanning booths and sunlamps. Use sunscreen and wear protective clothing when outdoors. Tell your doctor right away if you get sunburned or have skin blisters/redness.

Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products).

Older adults may be more sensitive to the side effects of this drug, especially low blood sugar.

During pregnancy, this medication should be used only when clearly needed. Pregnancy may cause or worsen diabetes. Discuss a plan with your doctor for managing your blood sugar while pregnant. Your doctor may substitute insulin for this drug during pregnancy. If glipizide is used, it may be switched to insulin at

least 2 weeks before the expected delivery date because of glipizide's risk of causing low blood sugar in your newborn. Discuss the risks and benefits with your doctor.

It is unknown if this medication passes into breast milk.

However, similar drugs pass into breast milk. Consult your doctor before breast-feeding.

DRUG INTERACTIONS:

See also How to Use section.

Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval.

Many drugs can affect your blood sugar, making it harder to control. Before you start, stop, or change any medication, talk with your doctor or pharmacist about how the medication may affect your blood sugar. Check your blood sugar regularly as directed and share the results with your doctor. Tell your doctor right away if you have symptoms of high or low blood sugar. (See also Side Effects section.) Your doctor may need to adjust your diabetes medication, exercise program, or diet.

Beta-blocker medications (including metoprolol, propranolol, glaucoma eye drops such as timolol) may prevent the fast/pounding heartbeat you would usually feel when your blood sugar falls too low (hypoglycemia). Other symptoms of low blood sugar such as dizziness, hunger, or sweating are unaffected by these drugs.

Check the labels on all your medicines (such as cough-and-cold products) because they may contain ingredients that could affect your blood sugar. Ask your pharmacist about using those products safely.

OVERDOSE:

If someone has overdosed and has serious symptoms such

as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Symptoms of overdose may include: shakiness, fast heartbeat, sweating, loss of consciousness.

NOTES:

Do not share this medication with others.

Attend a diabetes education program to learn more about how to manage your diabetes with medications, diet, exercise, and regular medical exams.

Learn the symptoms of high and low blood sugar and how to treat low blood sugar. Check your blood sugar regularly as directed.

Keep all regular medical and laboratory appointments. Laboratory and/or medical tests (such as liver and kidney function tests, fasting blood glucose, hemoglobin A1c, complete blood counts) should be performed periodically to monitor your progress or check for side effects.

MISSED DOSE:

If you miss a dose, take it as soon as you

remember. If it is near the time of the next dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up.

STORAGE:

Store at room temperature away from light and moisture.

Do not store in the bathroom. Keep all medications away from children and pets.

Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

MEDICAL ALERT:

Your condition can cause complications in a

medical emergency. For information about enrolling in MedicAlert, call 1-888-633-4298 (US) or 1-800-668-1507 (Canada).

Information last revised May 2022.

Copyright(c) 2022 First Databank, Inc.