

Brightside Specialty Pharmacy

431 N. Tustin Ave., STE C, Santa Ana, CA 92705

Phone 714-707-5115

Date :8/10/2022

Patient Education for HYDROXYZINE HCL 50 MG TABLET



Call doctor if no improvement in condition May cause blurred vision Decreases salivation; Use sugarless hard candy, gum or ice for relief Physician must know patient's complete medical history Avoid taking with other CNS depressant drugs or alcohol May cause drowsiness or dizziness Use caution driving. Check with doctor before increasing dose or frequency. May cause paradoxical excitability in patients, particularly children

Call doctor if you are not getting better May cause blurred vision May cause dry mouth; Use candy or gum for relief Tell doctor your complete medical history Avoid alcohol/other drugs that make you sleepy May make you drowsy or dizzy. Drive with caution Call Dr before increasing dose or frequency. May cause excitability in some patients & children.

.....Education Section.....

IMPORTANT:

HOW TO USE THIS INFORMATION: This is a summary and

does NOT have all possible information about this product. This information does not assure that this product is safe, effective, or appropriate for you. This information is not individual medical advice and does not substitute for the advice of your health care professional. Always ask your health care professional for complete information about this product and your specific health needs.

HYDROXYZINE HYDROCHLORIDE - ORAL

(hye-DROX-i-zeen HYE-droe-KLOR-ide)

COMMON BRAND NAME(S):

Atarax

USES:

Hydroxyzine is used to treat itching caused by allergies.

It is an antihistamine and works by blocking a certain natural substance (histamine) that your body makes during an allergic reaction. Hydroxyzine may also be used short-term to treat anxiety or to help you feel sleepy/relaxed before and after surgery.

HOW TO USE:

Take this medication by mouth with or without food

as directed by your doctor, usually three or four times daily. If you are using the liquid form of this medication, carefully measure the dose using a special measuring device/spoon. Do not use a household spoon because you may not get the correct dose.

The dosage is based on your age, medical condition, and response to treatment. In children, the dosage may also be based on weight. Do not increase your dose or take this medication more often than directed.

Tell your doctor if your condition does not improve or if it worsens.

SIDE EFFECTS:

Drowsiness, dizziness, blurred vision,

constipation, or dry mouth may occur. If any of these effects

last or get worse, tell your doctor or pharmacist promptly.

To relieve dry mouth, suck (sugarless) hard candy or ice chips, chew (sugarless) gum, drink water, or use a saliva substitute.

Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects.

Tell your doctor right away if you have any serious side effects, including: mental/mood changes (such as restlessness, confusion, hallucinations), shaking (tremor), difficulty urinating.

Get medical help right away if you have any very serious side effects, including: seizures, fast/irregular heartbeat, severe dizziness, fainting.

A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing.

This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist.

In the US -

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch.

In Canada - Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345.

PRECAUTIONS:

Before taking hydroxyzine, tell your doctor or

pharmacist if you are allergic to it; or to cetirizine; or to levocetirizine; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details.

Before using this medication, tell your doctor or pharmacist your medical history, especially of: breathing problems (such as

emphysema, asthma), high pressure in the eye (glaucoma), high blood pressure, kidney problems, liver problems, seizures, stomach/intestine problems (such as ulcer, blockage), overactive thyroid (hyperthyroidism), difficulty urinating (for example, due to enlarged prostate).

Hydroxyzine may cause a condition that affects the heart rhythm (QT prolongation). QT prolongation can rarely cause serious (rarely fatal) fast/irregular heartbeat and other symptoms (such as severe dizziness, fainting) that need medical attention right away.

The risk of QT prolongation may be increased if you have certain medical conditions or are taking other drugs that may cause QT prolongation. Before taking hydroxyzine, tell your doctor or pharmacist of all the drugs you take and if you have any of the following conditions: certain heart problems (heart failure, slow heartbeat, QT prolongation in the EKG), family history of certain heart problems (QT prolongation in the EKG, sudden cardiac death).

Low levels of potassium or magnesium in the blood may also increase your risk of QT prolongation. This risk may increase if you use certain drugs (such as diuretics/"water pills") or if you have conditions such as severe sweating, diarrhea, or vomiting. Talk to your doctor about taking hydroxyzine safely.

This drug may make you dizzy or drowsy or blur your vision. Alcohol or marijuana (cannabis) can make you more dizzy or drowsy. Do not drive, use machinery, or do anything that needs alertness or clear vision until you can do it safely. Avoid alcoholic beverages. Talk to your doctor if you are using marijuana (cannabis).

Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products).

Liquid products may contain sugar and/or alcohol. Caution is advised if you have diabetes, liver disease, or any other condition that requires you to limit/avoid these substances in your diet. Ask your doctor or pharmacist about using this product safely.

Children may be more sensitive to the side effects of this drug. This drug can often cause excitement in young children instead of drowsiness.

Older adults may be more sensitive to the side effects of this drug, especially drowsiness, confusion, constipation, trouble urinating or QT prolongation (see above). Drowsiness and confusion can increase the risk of falling.

During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor.

It is unknown if this drug passes into breast milk. Consult your doctor before breast-feeding.

DRUG INTERACTIONS:

Drug interactions may change how your

medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval.

Tell your doctor or pharmacist if you are taking other products that cause drowsiness such as opioid pain or cough relievers (such as codeine, hydrocodone), alcohol, marijuana (cannabis), drugs for sleep or anxiety (such as alprazolam, lorazepam, zolpidem), muscle relaxants (such as carisoprodol, cyclobenzaprine), or other antihistamines (such as diphenhydramine, promethazine).

Check the labels on all your medicines (such as allergy or cough-and-cold products) because they may contain ingredients that cause drowsiness. Ask your pharmacist about using those products safely.

Do not use with any other antihistamines applied to the skin (such as diphenhydramine cream, ointment, spray) because increased side effects may occur.

Hydroxyzine is very similar to cetirizine and levocetirizine. Do not use these medications while using hydroxyzine.

This medication may interfere with certain laboratory tests (including allergy skin testing, urine corticosteroids level), possibly causing false test results. Make sure laboratory personnel and all your doctors know you use this drug.

OVERDOSE:

If someone has overdosed and has serious symptoms such

as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Symptoms of overdose may include: severe drowsiness, seizures. In children, mental/mood changes (such as restlessness, irritability) may occur before drowsiness.

NOTES:

Do not share this medication with others.

MISSED DOSE:

If you miss a dose, take it as soon as you

remember. If it is near the time of the next dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up.

STORAGE:

Store at room temperature away from light and moisture.

Do not store in the bathroom. Do not freeze liquid forms of this medication. Keep all medications away from children and pets.

Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

Information last revised February 2022.

Copyright(c) 2022 First Databank, Inc.