

# Brightside Specialty Pharmacy

431 N. Tustin Ave., STE C, Santa Ana, CA 92705

Phone 714-707-5115

Date :8/10/2022

## Patient Education for

### OLANZAPINE 20 MG TABLET



May cause orthostatic hypotension Shake legs out before standing Call doctor if signs of tardive dyskinesia occur or get worse Rare:neuroleptic malignant syndrom prolonged fever & muscle rigidity Dispense Medication Guide and explain to patient before use May cause drowsiness or dizziness Use caution driving. Not recommended for dementia- related behavioral problems. May cause hyperglycemia, weight gain, and hyperlipidemia. May cause problems in newborns. Use with caution in 3rd trimester

Caution; Be careful not to stand up too quickly Call doctor if unusual muscle movements occur Call Dr. for unexplained fever & severe cramping Before use, important to read the Medication Guide May make you drowsy or dizzy. Drive with caution Tell doctor your complete medical history Have lab tests to monitor blood chemistry changes. Discuss risk vs benefits of use during pregnancy.

#### Education Section

#### IMPORTANT:

HOW TO USE THIS INFORMATION: This is a summary and

does NOT have all possible information about this product. This information does not assure that this product is safe, effective, or appropriate for you. This information is not individual medical advice and does not substitute for the advice of your health care professional. Always ask your health care professional for complete information about this product and your specific health needs.

OLANZAPINE - ORAL

(oh-LAN-za-peen)

#### COMMON BRAND NAME(S):

Zyprexa

#### WARNING:

There may be a slightly increased risk of serious,

possibly fatal side effects (such as stroke, heart failure, fast/irregular heartbeat, pneumonia) when this medication is used by older adults with dementia. This medication is not approved for the treatment of dementia-related behavior problems. Discuss the risks and benefits of this medication, as well as other effective and possibly safer treatments for dementia-related behavior problems, with the doctor.

If you are using olanzapine in combination with other medication to treat depression, also carefully read the drug information for the other medication.

#### USES:

Olanzapine is used to treat certain mental/mood conditions

(such as schizophrenia, bipolar disorder). It may also be used in combination with other medication to treat depression. This medication can help to decrease hallucinations and help you to think more clearly and positively about yourself, feel less agitated, and take a more active part in everyday life.

Olanzapine belongs to a class of drugs called atypical antipsychotics. It works by helping to restore the balance of certain natural substances in the brain.

Talk to the doctor about the risks and benefits of treatment

(especially when used by teenagers). See also Precautions section.

#### OTHER USES:

This section contains uses of this drug that are not

listed in the approved professional labeling for the drug but that may be prescribed by your health care professional. Use this drug for a condition that is listed in this section only if it has been so prescribed by your health care professional.

This drug may also be used to reduce the risk of nausea and vomiting caused by cancer drug treatment (chemotherapy).

#### HOW TO USE:

Read the Medication Guide provided by your

pharmacist before you start taking olanzapine and each time you get a refill. If you have any questions, ask your doctor or pharmacist.

Take this medication by mouth with or without food as directed by your doctor, usually once daily.

The dosage is based on your medical condition and response to treatment. To reduce your risk of side effects, your doctor may direct you to start this medication at a low dose and gradually increase your dose. Follow your doctor's instructions carefully.

Take this medication regularly to get the most benefit from it. To help you remember, take it at the same time each day. Keep taking this medication even if you feel well. Do not stop taking this medication without consulting your doctor.

Tell your doctor if your condition lasts or gets worse.

#### SIDE EFFECTS:

Drowsiness, dizziness, lightheadedness, stomach

upset, dry mouth, constipation, increased appetite, or weight gain may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly.

Dizziness and lightheadedness can increase the risk of falling. Get up slowly when rising from a sitting or lying position.

Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not

have serious side effects.

Tell your doctor right away if you have any serious side effects, including: difficulty swallowing, shaking (tremor), slow heartbeat, fainting, mental/mood changes (such as confusion, restlessness), numbness/tingling of arms/legs, yellowing eyes/skin, severe stomach/abdominal pain, trouble urinating, interrupted breathing during sleep.

This drug may rarely make your blood sugar rise, which can cause or worsen diabetes. Tell your doctor right away if you have symptoms of high blood sugar such as increased thirst/urination. If you already have diabetes, check your blood sugar regularly as directed and share the results with your doctor. Your doctor may need to adjust your diabetes medication, exercise program, or diet.

This drug may also cause significant weight gain and a rise in your blood cholesterol (or triglyceride) levels, especially in teenagers. These effects, along with diabetes, may increase your risk for developing heart disease. Discuss the risks and benefits of treatment with your doctor. (See also Notes section.)

Olanzapine may rarely cause a condition known as tardive dyskinesia. In some cases, this condition may be permanent. Tell your doctor right away if you develop any unusual/uncontrolled movements (especially of the face, lips, mouth, tongue, arms or legs).

This medication may increase a certain natural substance (prolactin) made by your body. For females, this increase in prolactin may result in unwanted breast milk, missed/stopped periods, or difficulty becoming pregnant. For males, it may result in decreased sexual ability, inability to produce sperm, or enlarged breasts. If you develop any of these symptoms, tell your doctor right away.

Get medical help right away if you have any very serious side effects, including: seizures.

This medication may rarely cause a very serious condition called neuroleptic malignant syndrome (NMS). Get medical help right away if you have any of the following symptoms: fever, muscle stiffness/pain/tenderness/weakness, severe tiredness, severe confusion, sweating, fast/irregular heartbeat, dark urine, signs of kidney problems (such as change in the amount of urine).

A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: fever, swollen lymph nodes, rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing.

This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist.

In the US -

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

In Canada - Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345.

#### **PRECAUTIONS:**

Before taking olanzapine, tell your doctor or

pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details.

Before using this medication, tell your doctor or pharmacist your medical history, especially of: liver problems, seizures, difficulty swallowing, low white blood cell count, dementia, difficulty urinating (for example, due to enlarged prostate), glaucoma (narrow angle), stomach/intestinal problems (such as blockage, chronic constipation, paralytic ileus), smoking, personal or family history of diabetes, heart disease, high cholesterol/triglyceride levels, breathing trouble during sleep (sleep apnea).

This drug may make you dizzy or drowsy. Alcohol or marijuana (cannabis) can make you more dizzy or drowsy. Do not drive, use machinery, or do anything that needs alertness until you can do it safely. Avoid alcoholic beverages. Talk to your doctor if you are using marijuana (cannabis).

Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products).

This medication may make you sweat less, making you more likely to get heat stroke. Avoid doing things that may cause you to overheat, such as hard work or exercise in hot weather, or using hot tubs. When the weather is hot, drink a lot of fluids and dress lightly. If you overheat, quickly look for a place to cool down and rest. Get medical help right away if you have a fever that does not go away, mental/mood changes, headache, or dizziness.

Teenagers may be more sensitive to the side effects of this drug, especially weight gain, and also increased amounts of cholesterol, triglycerides, and prolactin. See also Side Effects section for more details.

Older adults may be more sensitive to the side effects of this drug, especially drowsiness, constipation, trouble urinating, confusion, dizziness, and lightheadedness. Drowsiness, confusion, dizziness, and lightheadedness can increase the risk of falling.

During pregnancy, this medication should be used only when clearly needed. Babies born to mothers who have used this drug during the last 3 months of pregnancy may rarely develop symptoms including muscle stiffness or shakiness, drowsiness, feeding/breathing difficulties, or constant crying. If you notice any of these symptoms in your newborn especially during their first month, tell the doctor right away.

Since untreated mental/mood problems (such as schizophrenia, bipolar disorder, depression) can be a serious condition, do not stop taking this medication unless directed by your doctor. If you are planning pregnancy, become pregnant, or think you may be pregnant, immediately discuss with your doctor the benefits and risks of using this medication during pregnancy.

This drug passes into breast milk and may have undesirable effects on a nursing infant. Tell the doctor right away if your baby develops symptoms such as muscle stiffness or shakiness, unusual sleepiness, or difficulty feeding. Consult your doctor before breast-feeding.

#### **DRUG INTERACTIONS:**

Drug interactions may change how your

medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval.

A product that may interact with this drug is: metoclopramide.

Tell your doctor or pharmacist if you are taking other products that cause drowsiness such as opioid pain or cough relievers (such as codeine, hydrocodone), alcohol, marijuana (cannabis), drugs for sleep or anxiety (such as alprazolam, lorazepam, zolpidem), muscle relaxants (such as carisoprodol, cyclobenzaprine), or antihistamines (such as cetirizine, diphenhydramine).

Check the labels on all your medicines (such as allergy or cough-and-cold products) because they may contain ingredients that cause drowsiness. Ask your pharmacist about using those products safely.

**OVERDOSE:**

If someone has overdosed and has serious symptoms such

as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Symptoms of overdose may include: severe drowsiness/dizziness, fast/irregular heartbeat, unusual/uncontrolled movements, seizures.

**NOTES:**

Do not share this medication with others.

Laboratory and/or medical tests (such as blood sugar, weight, blood pressure, blood cholesterol/triglyceride levels, liver function tests) should be performed periodically to monitor your progress or check for side effects. Consult your doctor for more details.

**MISSED DOSE:**

If you miss a dose, take it as soon as you

remember. If it is near the time of the next dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up.

**STORAGE:**

Store at room temperature away from light and moisture.

Do not store in the bathroom. Keep all medications away from children and pets.

Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

Information last revised April 2022.

Copyright(c) 2022 First Databank, Inc.